



Australian Government

Department of the Environment,
Water, Heritage and the Arts



Threat abatement plan
for the biological effects,
including lethal toxic ingestion,
caused by cane toads (Draft)

© Commonwealth of Australia 2009

This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation. Apart from any use as permitted under the *Copyright Act 1968*, all other rights are reserved. Requests and inquiries concerning reproduction and rights should be addressed to Commonwealth Copyright Administration, Attorney General's Department, Robert Garran Offices, National Circuit, Barton ACT 2600 or posted at www.ag.gov.au/cca.

Disclaimer

The contents of this document have been compiled using a range of source materials and is valid as at March 2010. The Australian Government is not liable for any loss or damage that may be occasioned directly or indirectly through the use of or reliance on the contents of the document.

**Threat abatement plan for the biological effects, including
lethal toxic ingestion, caused by cane toads
(Draft)**

MARCH 2010

SUMMARY	3
<i>Rationale</i>	3
<i>Objectives for the threat abatement plan</i>	3
<i>Implementation of the threat abatement plan</i>	4
1. INTRODUCTION	4
<i>1.1 Threat abatement plans</i>	4
<i>1.2 Threat abatement plan for cane toads</i>	5
1.2.1 The threat	5
History	5
Ecological impacts	7
1.2.2 Managing the threat	7
1.2.3 Involvement of stakeholders	10
<i>1.3 Definition of priority native species and ecological communities</i>	10
2. OBJECTIVES AND ACTIONS	11
<i>Objective 1 – Identify priority native species and ecological communities at risk from the impact of cane toads</i>	11
Ecological communities	11
Species	12
Recommended actions and priorities	15
Performance indicators	15
<i>Objective 2 - Reduce the impact on populations of native species and ecological communities</i>	16
Performance indicators	18
<i>Objective 3 – Communicate information about cane toads, their impacts and this TAP</i>	19
Performance indicators	20
3. DURATION, IMPLEMENTATION AND EVALUATION OF THE PLAN	20
<i>3.1 Duration of the plan</i>	20
<i>3. 2 Cost of the plan</i>	21
<i>3.3 Implementing the plan</i>	22
<i>3.4 Evaluating and reviewing the plan</i>	22
REFERENCES	23
APPENDICES	26

Draft cane toad threat abatement plan

Summary

Rationale

Since the introduction of cane toads (*Bufo marinus*, now revised to *Rhinella marina*) to Australia in 1935, the ecological impact of this animal has aroused considerable concern. Cane toads use potent steroid-derived toxins as chemical defences. The active constituents of these differ from the toxins found in native frogs. All life stages of the cane toad (eggs, tadpoles, metamorphs and adults) are toxic, although toxin types and content change markedly during a toad's lifespan. Toxin levels (and thus, danger to native vertebrate predators) are high in eggs, decrease through tadpole life, are lowest at around the time of metamorphosis, and increase rapidly thereafter. It is difficult to tease apart the effects of cane toads from other threatening processes operating on native species and ecological communities (Shine 2009a). However, there is no scientific evidence that cane toads have caused species extinction. The direct pathway of lethal toxic ingestion of cane toads is the most important cane toad impact.

In 2005, the biological effect of the cane toad was listed as a key threatening process under the *Environment Protection and Biodiversity Conservation Act 1999* (Threatened Species Scientific Committee 2005).

Since 1986, the Australian Government has directed at least \$11 million dollars to development of a broad-scale means to control cane toads without success. Community action to manually remove cane toads from the landscape has also been funded. Neither of these endeavours have prevented the continued spread of the pest or significantly limited its impact on Australia's biodiversity. Recognising that it is not currently possible to contain or eradicate cane toads across the nation, a new approach to dealing with their negative impacts is needed. This involves identifying and reducing impacts on key natural assets affected by cane toads, an approach that requires national coordination.

This threat abatement plan (TAP) provides a national strategy to guide investment and effort by the Australian Government, jurisdictions, research organisations and non-government organisations in abating the impacts of cane toads across their known and anticipated range.

Objectives for the threat abatement plan

This TAP has three objectives:

- to identify priority native species and ecological communities at risk from the impact of cane toads
- to reduce the impact of cane toads on populations of priority native species and ecological communities
- to communicate information about cane toads, their impacts and this TAP.

Implementation of the threat abatement plan

This TAP will be implemented by the Australian Government in conjunction with a broad range of stakeholders.

1. Introduction

This threat abatement plan (TAP) has been developed to address the listed key threatening process *The biological effects, including lethal toxic ingestion, caused by Cane Toads (Bufo marinus)* (see listing advice, Threatened Species Scientific Committee 2005) in a feasible, effective and efficient manner. The TAP binds the Australian Government and its agencies in Australia's response to the impact of cane toads and identifies the research, management and other actions needed to address the impacts of this species on Australia's biodiversity.

This plan should be read in conjunction with the two publications: *The ecological impact of invasive cane toads (Bufo marinus) in Australia* (Shine 2009a); and *Cane Toads in Communities- Executive Report* (Bureau of Rural Sciences 2009). These publications provide information on the scope of the cane toad threat and public perceptions of cane toads and their impacts across the known and anticipated range of the species.

1.1 Threat abatement plans

Under section 270 (A) of the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act), the Australian Government:

- develops TAPs
- implements the actions under TAPs that are its direct responsibility
- facilitates the implementation of actions where other groups (e.g. states and territories, industry) share the implementation responsibilities.

The EPBC Act prescribes the process, content and consultation required when making a TAP.

The EPBC Act requires the Australian Government to implement TAPs to the extent to which they apply in areas under Australian Government control and responsibility. In addition, Australian Government agencies must not take any actions that contravene a TAP. Where a TAP applies outside Australian Government areas in states or territories, the Australian Government must seek the cooperation of the affected jurisdictions, with a view to jointly implementing the plan.

The EPBC Act also has requirements relating to Matters of National Environmental Significance (for example listed threatened species and communities, World Heritage listed areas and Ramsar Agreement listed wetlands - see Significant Impact Guidelines (Commonwealth of Australia 2006)). As some of these matters may be affected by a specific threat, appropriate Matters of National Environmental Significance may also be addressed in a TAP.

The Department of the Environment, Water, Heritage and the Arts prepares a five-year project plan for each TAP and assesses progress on the main strategic

actions contained within the TAP on a yearly basis. After five years, each TAP is reviewed to ensure the objectives of the TAP have been achieved.

Mitigating the impact of invasive species is not simply a matter of developing and applying better technical solutions. It also involves the development of better biological and ecological information, as well as understanding and addressing the social and economic factors surrounding the pest. In the case of cane toads, the need to move away from attempts at broad-scale control and eradication to the protection of key biodiversity assets will require the transfer of knowledge on the management of cane toad impacts, as well as support for community effort to limit those impacts.

This new focus, on protection of key assets, results from a recently improved understanding of the impacts of cane toads and from the lack of progress on broad-scale control. Research has demonstrated that these impacts are not generally as negative as may have been expected historically and that adaptation to cane toads can rapidly occur in some native species (Professor Rick Shine [School of Biological Sciences, The University of Sydney] 2010, pers. comm., 12 January)

Communication of scientific evidence regarding which species are at risk will be the key factor in this new approach to protecting biodiversity from cane toad impacts.

1.2 Threat abatement plan for cane toads

1.2.1 The threat


History

Cane toads were introduced to Australia in 1935 as a means of controlling pest beetles in the sugar cane industry. This is a process that was common to many sugar cane or other crop producing areas of the world (including Puerto Rico, Papua New Guinea and Fiji). At some locations, cane toads failed to establish (e.g. Egypt, where they were introduced in 1937). However, in many others locations cane toads survived and established to become pests. Attempts at cane toad management and control have been most extensive in Australia (Global Invasive Species Database 2009).

The success of cane toads in pest insect control in Australia was never determined, as the use of agricultural chemicals for this purpose became widespread soon after their release (Shine 2009b). Cane toads, however, did become very successful at invading the environments of Australia's north. Since 1935, they have dispersed over 2000 km west from their release site at Gordonvale, Queensland and many hundreds of kilometres to the north and south. Their southern dispersal includes areas considered to be marginal cane toad habitat in arid south-west Queensland and the cooler climates and higher altitudes of northern New South Wales (Figure 1).

In the first few decades after cane toads were released in Queensland, they expanded their range at about 10 km per year (Shine 2009c). Since reaching the wet-dry tropics of the Northern Territory, they have progressed, on average, around 55 km each year (Phillips et al. 2007).

135 **Figure 1: Map showing approximate current extent (black line) and anticipated distribution (shaded areas) of cane toads in Australia**
 136 **(Kearney et al. 2008).**



Ecological impacts

Across their range, cane toads have been implicated in a complex web of direct and indirect impacts on native species and potentially the ecological communities in which these species occur. Assessing cane toad impacts through scientific research is very difficult (Shine 2009a). However, available evidence identifies the direct pathway of lethal toxic ingestion of cane toads as the most important impact. Many native Australian predators have evolved in the absence of prey species with the chemical defences present in cane toads. Consequently, predators are vulnerable to being lethally poisoned when cane toads invade and establish in their areas. No species extinction has ever been attributed to the cane toad, however, research has identified vulnerable predator species and other ecological impacts.

Local population extinctions of the endangered marsupial predator, the northern quoll (*Dasyurus hallucatus*), have been observed following the arrival of cane toads in some areas. Lethal toxic poisoning through ingestion of the cane toad has been identified as the cause of these local extinctions. Several species of goannas, snakes and the blue tongued lizard (*Tiliqua scincoides intermedia*) have also been identified as highly vulnerable to lethal poisoning through ingestion of cane toads (Shine 2009a).

Other pathways of cane toad impact on native species have been identified (Shine 2009a) as:

- predation by cane toads (varies, predominantly a minor impact)
- larval competition with frog tadpoles or mosquitoes (minor impact)
- parasite transfer (minor impact)
- competition for terrestrial food (minor impact)
- competition for shelter sites (e.g. usurpation of burrows) (minor impact).

Historically, in the absence of scientific evidence about the complexities of direct and indirect impacts of cane toads, anecdotal information has served to provide guidance on impacts and priorities for managing them.

Within their current and anticipated ranges, cane toads have had a significant impact on a number of Matters of National Environmental Significance. For example, cane toads have impacted:

- listed threatened species such as the northern quoll *Dasyurus hallucatus* – an EPBC listed endangered species
- Wetlands of International Importance listed under the terms of the Ramsar Convention, including iconic wetlands in Kakadu National Park
- World Heritage properties such as the Wet Tropics of Queensland.

1.2.2 Managing the threat

Since 1986, the Australian Government has directed approximately \$11 million to development of a broad-scale means to control cane toads and a further \$9 million to other cane toad research and management activities (details in Table 1).

Table 1: Australian Government funding on cane toad research and management 1986 to 2009.

Area of expenditure	
Research on impacts	\$5,212,518
Research control (long term/biological control)	\$11,111,922
Research control (short/medium term control)	\$1,303,235
Management	\$1,162,117
Community groups	\$1,283,234
Education	\$44,468
TOTAL	\$20,117,494

Over \$11 million of the Australian Government cane toad funding has been provided for the search for a biological control agent in the toads' native habitat in South America, and research directed at modifying a virus in order to disrupt the development of infected cane toad tadpoles. In 2008, an independent review of the CSIRO's cane toad biological control research (Shannon and Bayliss 2008) resulted in funding for this project being discontinued. The review team found that "there are still major technical hurdles to be overcome in the development of a self-disseminating genetically modified cane toad control agent. The long term feasibility of the approach is also questionable on several counts including the availability of an acceptable viral vector, the difficulty of generating an appropriate immune response from virally expressed proteins, and the major hurdle of obtaining approval for release. The lack of a national and international risk assessment and management plan for the release of a virally vectored genetically modified organism regardless of exact product specification is also a major deficit and should be an essential part of any further program in this area".

To date no broad-scale or biological control has been identified and it is unlikely that such a control could be developed and approved for use before the cane toad will have reached its maximum extent (see Figure 1) and impact.

Community action to manually remove cane toads from the landscape has also received Australian Government funding (approx \$1.3 million from 1986 to 2009). Government agencies in Queensland, New South Wales, Western Australia and the Northern Territory have contributed to cane toad control efforts. However, there is no evidence that these endeavours have prevented the continued spread of the pest or significantly limited its impact on Australia's biodiversity. Community action, while satisfying to local communities, does not have the capacity to make any significant changes to the rate of spread of cane toads or to the densities of cane toads beyond specific local areas. However, where community action is focused on cane toad management to protect assets at a local scale it could help maintain priority biodiversity assets.

A decade of effort around Port Macquarie may have resulted in local eradication from that area. However, the cane toad is likely to be towards the southern limits of its "natural" range in this region of northern New South Wales, and climatic factors may

224 have assisted control efforts. An additional factor in the success of this effort is likely
225 to have been the status of the Port Macquarie infestation as an isolated satellite
226 population (Peacock 2007).

227
228 *The biological effects, including lethal toxic ingestion, caused by Cane Toads (Bufo*
229 *marinus)* (Threatened Species Scientific Committee 2005) was listed as a key threatening
230 process under the EPBC Act in 2005 in response, in particular, to concerns about the
231 impact of cane toads on the northern quoll. At the time of this listing, the then
232 Minister for Environment and Heritage decided that the development of a TAP would
233 not be an efficient way to abate the threat posed by cane toads.

234
235 This decision has subsequently been reviewed by the Minister for the Environment,
236 Heritage and the Arts. Consultation with colleagues in state and territory governments
237 about the feasibility, effectiveness and efficiency of developing and implementing a
238 TAP to abate the cane toad threat was undertaken, and national coordination emerged
239 as a dominant theme in support of developing a TAP at this time.

240
241 This TAP provides a national strategy to guide investment and effort by the Australian
242 Government, jurisdictions, research organisations and non-government organisations
243 in abating the impact of cane toads across their known and anticipated range. This
244 TAP identifies key assets (native species and ecological communities) to be protected,
245 discusses protection methods, and identifies the need to develop humane control
246 methods for cane toads.

247
248
249 **Eradication of cane toads is not currently possible. Neither the**
250 **resources nor the technologies required to contain and eradicate cane**
251 **toad numbers on a continental scale are available. The timescales**
252 **required for the development and application of such technologies**
253 **would mean that cane toads will have reached the extent of their**
254 **continental impact regardless of the investment made.**

255
256 **Recognising the new information now available about cane toads and**
257 **their impacts, as well as the failure of past attempts at broad-scale**
258 **control, this TAP takes the approach of identifying and prioritising**
259 **the native species and ecological communities under threat from cane**
260 **toads, and targeting action to protect those assets which have been**
261 **determined to be of the highest priority.**

262
263 **This approach will focus on achieving positive biodiversity outcomes**
264 **for species or ecological communities vulnerable to the presence of**
265 **cane toads. This approach has evolved as the efforts undertaken to**
266 **date have neither provided a broad-scale control method such as a**
267 **biological control, nor an effective answer to the expansion of the**
268 **toads' range through manual removal. Both of these approaches have**
269 **been proven to be an ineffective use of limited natural resource funds.**
270 **This new approach will allow for a more effective and efficient use of**
271 **conservation resources at the national, state, territory and local levels**
272 **than is occurring under current strategies.**

1.2.3 Involvement of stakeholders

The success of this TAP will depend on a high level of **cooperation between all key stakeholders**, including:

- the Australian Government and its agencies
- state and territory conservation and resource management agencies
- local government
- natural resource management agencies and private conservation land management bodies
- research institutes
- industry and entrepreneurs
- Indigenous communities
- other community groups.

The Invasive Animals CRC currently supports a Cane Toad Advisory Group (CTAG). This committee is comprised of Australian Government, state and territory representatives and provides strategic and practical advice on the planning, implementation and delivery of cane toad projects and their outcomes. The CTAG provides a mechanism to focus national and jurisdictional understanding of, and efforts to abate, cane toad impacts and via its links to the Vertebrate Pests Committee will serve as a major coordination point for actions undertaken under this TAP. Major outcomes will be communicated from this group to local government, natural resource management agencies, conservation groups, industry and entrepreneurs, conservation bodies and community groups in each jurisdiction.

Ongoing delivery of awareness and capacity building programs in natural resource management will be required at national, state and regional levels and will make a significant contribution to national implementation of this TAP.

Implementation of some of the objectives of this TAP (e.g. identification and prioritisation of native species and ecological communities) will require specific efforts from the Australian Government and jurisdictions. However, as information is collated, and priorities determined, other stakeholders will have strong locally focused responsibilities for ensuring actions are undertaken to protect biodiversity assets that are impacted by cane toads.

1.3 Definition of priority native species and ecological communities

For the purposes of this TAP, priority native species and ecological communities are those that have been determined through peer-reviewed research to be highly vulnerable at population level to negative impacts from the presence of cane toads.

At the national level, relevant Matters of National Environmental Significance and the National Reserve System will also be considered. For state and territory agencies, this TAP can guide investment based on state or regional conservation priorities. It will be important that managers assess the impacts of cane toads and allocate adequate resources to achieving effective management at all priority sites (national, jurisdictional, regional, local) and that outcomes are measured and assessed on an on-going basis.

Jurisdictional governments, natural resource management groups and community groups will need to determine more localised priority assets and the means by which they will undertake protection and management actions.

2. Objectives and actions

This TAP has three objectives:

- identify priority native species and ecological communities at risk from the impact of cane toads
- reduce the impacts of cane toads on populations of priority native species and ecological communities
- communicate information about cane toads, their impacts and this TAP.

Supporting actions to implement these objectives are listed below.

Objective 1 – Identify priority native species and ecological communities at risk from the impact of cane toads

There are neither the resources nor an appropriate broad-scale control that can be applied to the management of cane toads in a way that would lead to containment and/or eradication of cane toads across their range. However, the Australian Government has a responsibility to manage cane toads on land under its control and where Matters of National Environmental Significance are being impacted by cane toads. Objective 1 addresses the identification of those species and ecological communities at risk from the impact of cane toads.

Ecological communities

There are eight threatened ecological communities listed under the EPBC Act that fall within the current geographic range of the cane toad (Table 2).

Table 2: EPBC Act listed threatened ecological communities within the current cane toad range.

Ecological community	EPBC category
Swamp Tea-tree (<i>Melaleuca irbyana</i>) Forest of South-east Queensland	Critically Endangered
Mabi Forest (Complex Notophyll Vine Forest 5b)	Critically Endangered
White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland	Critically Endangered
Natural Grasslands of the Queensland Central Highlands and the northern Fitzroy Basin	Endangered
Semi-evergreen vine thickets of the Brigalow Belt (North and South) and Nandewar Bioregions	Endangered

Ecological community	EPBC category
The community of native species dependent on natural discharge of groundwater from the Great Artesian Basin	Endangered
Weeping Myall Woodlands	Endangered
Brigalow (<i>Acacia harpophylla</i> dominant and co-dominant)	Endangered

Currently, none of the listing advices for these communities indicate that cane toads are a threat to the community. Further, no other listed ecological communities fall within the predicted future range of cane toads (as shown in Figure 1 Kearney et al. 2008). However, elements of some of the listed communities are impacted, or may in future be impacted, by cane toads as they continue to spread.

Kearney et al. (2008) used the software package Ozclim (CSIRO, Australia) to derive predictions for changes in monthly maximum and minimum air temperature and relative humidity, as well as mean monthly rainfall by 2050. Under this anticipated climate change scenario for Australia, both expansions and contractions in the potential range of the cane toad and in the length of the toads' breeding season have been predicted for 2050. The southern border of cane toad distribution is predicted to move further south by approximately 100 km and be limited by the opposing influences of increasing air temperature and decreasing humidity on the core body temperature of cane toads. In this scenario, further ecological communities listed under the EPBC Act would fall within the range of cane toads.

Native species and ecological communities on off-shore islands may need to be protected from cane toads. Quarantine or emergency management measures to protect these islands may result in the preservation of endemic island species and ecological communities. Further, it may be possible, under particular circumstances, to protect populations of those species identified as highly impacted on the mainland (Table 3) by preserving populations already present on islands or relocating species from the mainland to islands. Islands known to be free of cane toads and which support populations of species highly impacted by cane toads on the mainland, have been identified (Appendices 1, 2 and 3).

Species

Although many individual animals may succumb to lethal toxic ingestion of cane toads, particularly when toads first appear in a new area, the number of species known to be negatively affected at a population level is small. It is this group that forms the highest priority for action under this TAP. Research is currently being undertaken by several groups (e.g. the University of Sydney, Australian National University) to clarify the impact of toads on certain species such as the northern quoll and goanna species. This research may provide insights into priority species for protection over the life of this TAP.

The ecological impact of invasive cane toads (Bufo marinus) in Australia (Shine 2009a) provides an extensive scientific assessment of the impacts of cane toads on native species. A summary of this assessment, listing those native species for which the level of negative population impact by cane toads is high or moderate is

provided (Table 3). This provides an initial assessment of priority for species requiring population level protection. Research is continuing in this area, and the lists may need to be adapted as understanding improves.

Table 3: Current state of knowledge of identified high or moderate negative population level threats to Australian native fauna from the cane toad (modified from Shine 2009a). Lethal toxic ingestion is the most common pathway of impact.

Species	Degree of impact	Authority	Pathway for impact
Proteocephalid Tapeworm*	High	Freeland 1993, 2004	Toads destabilise host / parasite equilibrium
Crocodiles			
Freshwater crocodile (in semi-arid landscapes) <i>Crocodylus johnstoni</i>	High (location dependent)	Letnic et al. 2008	Lethal toxic ingestion
Goannas			
<i>Varanus spp.</i>	High	Freeland 2004; Griffiths and McKay 2007; Doody et al. 2009; Ujvari and Madsen 2009	Lethal toxic ingestion
Skinks			
<i>Tiliqua scincoides intermedia</i>	High	Price-Rees et al. 2010	Lethal toxic ingestion
Snakes			
Northern death adder <i>Acanthophis praelongus</i>	High	Hagman et al. 2009	Lethal toxic ingestion
King brown snake <i>Pseudechis australis</i>	High	G.P. Brown et al. University of Sydney unpublished data	Lethal toxic ingestion
Marsupials			
Northern quoll <i>Dasyurus hallucatus</i>	High	Oakwood 2003 a,b; O'Donnell 2009	Lethal toxic ingestion
Eutherian mammals			
Pale field-rat <i>Rattus tunneyi</i>	Moderate	Watson and Woinarski 2003	Unknown

* **NB**, This tapeworm is a parasite of the spotted python *Antaresia maculosa* and has been described to family level only (Proteocephalidae).

The northern quoll is the only species listed in Table 3 (above) that is also listed under the EPBC Act (as Endangered). Significant actions have taken place to protect the northern quoll through the Northern Territory's *Island Arks* program, as a result of documented decline of the species with the arrival of cane toads (Rankmore et al. 2008). This program has provided 'insurance populations' of the species on two toad-free islands. While the program has been highly successful in establishing populations of northern quolls on the islands, it has not yet attempted to reintroduce any individual animals to their original habitats.

Species, for which there is suspicion, but not scientific certainty, of population level impacts by cane toads, have also been identified (Table 4).

Table 4: Current state of knowledge on uncertain negative population-level threats to Australian native fauna from the cane toad (modified from Shine 2009a).

Species	Type of impact	Degree of impact	Authority
Dragons (Agamidae) Frilled lizard <i>Chlamydosaurus kingii</i>	Lethal toxic ingestion	Reports inconsistent	van Dam et al. 2002; T. Madsen pers. comm.
Birds Rainbow bee-eater <i>Merops ornatus</i>	Usurpation of burrows	unknown	Boland 2004

Future research may require other species to be added to this list, or the list at Table 3.

Recommended actions and priorities

Action	Priority
Action 1.1 Identify native species, ecological communities and off-shore islands currently known to be at high to moderate risk. (Largely completed).	High priority, short term because currently underway
Action 1.2 Identify the ways in which cane toads impact the native species and ecological communities listed in 1.1 (Largely completed).	High priority, short term because currently underway
Action 1.3 Where impact is unknown but may be high, establish and support research to further understand the impact of cane toads on the native species and ecological communities. Where appropriate, research ways to assist with the recovery of priority native species and ecological communities. (Has commenced).	Medium priority, medium term
Action 1.4 Develop a prioritisation tool to guide allocation of resources for protection of native species and communities. Apply it to native species and ecological communities identified: first from Action 1.1, then from Action 1.3.	Low priority, medium term

The criteria to be used in the prioritisation tool (Action 1.4) will include:

- protection of cane toad-free off-shore islands, particularly those that currently support populations of native species identified in Table 3 (Appendices 1, 2 and 3 contain a preliminary list of islands known to fall within this category)
- protection of those species identified in Table 3
- capacity to add species when evidence of impact becomes clear (e.g. species listed under Table 4)
- Matters of National Environmental Significance.

Performance indicators

- Listing of ecological communities, species and off-shore islands at risk from the impacts of cane toads developed by the CTAG and the VPC, agreed by VPC and made available to all stakeholders within 12 months of the making of this TAP.
- Scientific evidence, endorsed by the CTAG and VPC, is gathered for those species for which high impact from cane toads is currently suspected, but not yet confirmed within 18 months of the making of this TAP.
- Research, which improves scientific understanding of impacts on native species and ecological communities; improves understanding of recovery measures; and which informs resourcing agreements between the Australian Government and affected jurisdictions is endorsed by the VPC as it becomes available.
- Prioritisation tool for allocation of resources to ecological communities/species developed and agreed by the CTAG and the VPC within the 24 months of the making of this TAP.

- Prioritisation tool applied at a national level and application encouraged at jurisdictional, regional and local levels within 6 months of the prioritisation tool being agreed by the VPC.

Objective 2 - Reduce the impact on populations of native species and ecological communities

Under Objective 1, actions to determine the priority for the application of resources to the management of cane toads and their impacts will be developed. Listings and mapping of threatened ecological communities and species will be determined. The Australian Government will address those that are on land under its control or are Matters of National Environmental Significance in conjunction with state and territories. These listings and maps will also enable stakeholders to determine state, regional and local priorities and apply appropriate resources to their protection.

The purpose of Objective 2 is to develop effective tools that can be used to reduce the impact of cane toads on native species. The tools will cover all aspects of cane toad management at the planning and response stages, and be broadly applicable.

Use of these guidelines will be the responsibility of all stakeholders, in particular those with land and water management responsibilities in areas identified as being of priority for protection against cane toads. The Australian Government will be monitoring the uptake of management actions in each of the identified priority areas. Where the Australian Government and state/territory governments have mutual obligations (e.g. some Ramsar Wetlands) negotiation of appropriate actions and funding of management actions will be undertaken.

While the purpose of this TAP is not to develop specific cane toad control tools, such as poisons, research is underway that could result in a larger toolkit becoming available over the life of this TAP. These could include:

- development and registration of a humane lethal spray for toads
- use of a larval alarm pheromone to manage cane toad populations within water bodies (Hagman and Shine 2009; Hagman et al. 2009)
- use of a parasitic nematode of cane toads (*Rhabdias pseudosphaerocephala*) identified as present in established populations of Australian cane toads (Dubey and Shine 2008; Shine 2009b)
- development of better traps.

As such tools become available, information about them will be included on the DEWHA cane toad webpage.

Action	Priority and timeframe
<p>Action 2.1 Focus management of cane toad impacts by Australian Government agencies on designated high priority native species and ecological communities, and seek cooperative action on priorities by jurisdictions and other stakeholders.</p> <p>Action 2.1.1 Implement and monitor emergency management of cane toad impacts for known high priority native species and ecological communities (as designated in Table 3) using currently available tools and techniques (e.g. trapping, fencing of small areas, manual removal from designated sites)</p> <p>Action 2.1.2 As new species and communities are added to the list of priority native species and ecological communities via a peer reviewed process, implement or adjust management of cane toad impacts, using available tools and techniques. Tools and techniques will increase with the designation of poisons, development of codes of practice and standard operating procedures and potentially with the development of new techniques.</p>	<p>High priority, short term</p> <p>High priority, short term</p> <p>Medium priority, medium term</p>
<p>Action 2.2 Prepare guidelines, including codes of practice and standard operating procedures, that can be applied to both emergency responses and on-going management for high priority native species and ecological communities for endorsement by the VPC.</p> <p>Action 2.2.1 Australian Government to prepare and implement management plans, (including identifying and addressing gaps in management techniques and tools) for designated high priority species and ecological communities on land managed by Australian Government agencies.</p> <p>Action 2.2.2 Provide the guidelines for emergency and on-going cane toad management to all stakeholders. Liaise with responsible jurisdictions/agencies to encourage the preparation and implementation of such plans in their areas of responsibility. Where mutual obligations exist the Australian Government will work cooperatively to prepare such plans.</p>	<p>Medium priority, medium term</p> <p>Medium priority, medium term</p> <p>Medium priority, medium term</p>

<p>Action 2.2.3 Australian Government to monitor the development and implementation of guidelines and cane toad management plans for designated high priority species and ecological communities.</p>	Medium priority, medium to long term
<p>Action 2.2.4 Australian Government to monitor the literature about the spread and impact of the cane toad and review/amend guidelines and develop new management plans as required.</p>	Medium priority, medium to long term
<p>Action 2.3 Establish guidelines for humane management actions to control cane toads for VPC and Animal Welfare Committee endorsement.</p>	Medium priority, medium term
<p>Action 2.3.1 Distribute guidelines to all Australian Government agencies with land management responsibilities.</p>	Medium priority, medium term
<p>Action 2.3.2 Australian Government to seek cooperative adoption of guidelines by states/territories including incorporation in state based regulations as appropriate.</p>	Medium priority, medium term

487

488 Performance indicators

- 489 • Australian Government agencies advised of this TAP, the designation of high
- 490 priority species and ecological communities, and their management
- 491 responsibilities within six months of the TAP being made.
- 492 • Application of this TAP by Australian Government agencies is monitored by
- 493 these agencies and DEWHA over the life of the TAP.
- 494 • Jurisdictions and stakeholders are advised of this TAP, the national high
- 495 priority species and ecological communities and the jurisdictions/stakeholders'
- 496 management responsibilities within six months of the TAP being made.
- 497 • Management plans agreed with relevant stakeholders (state/territory
- 498 governments) for each of those species impacted by cane toads at a population
- 499 level within eighteen months of this TAP being made.
- 500 • Responses of jurisdiction and other stakeholders are monitored throughout the
- 501 life of the TAP.
- 502 • Additional advice is provided to all stakeholders within six months of new
- 503 species and ecological communities being identified and agreed or removed
- 504 from the list of priority species.
- 505 • Guidelines for emergency and on-going management plans are developed and
- 506 agreed by VPC within two years of the TAP being made.
- 507 • Guidelines are provided to all stakeholders within three months of being
- 508 agreed by VPC.
- 509 • Preparation of plans across all land tenures for high priority species and
- 510 ecosystems are monitored by the Australian Government on an on-going basis
- 511 across the life of the TAP.

- Humane management actions (standard operating procedures and codes of practice for humane treatment of cane toads) are developed and agreed by the VPC within two years of the TAP being made.

Objective 3 – Communicate information about cane toads, their impacts and this TAP

Australians are concerned about the impact of cane toads on the environment. However, community concern is highest when cane toads incursion is recent or imminent and fades over time as the community adjusts to living with cane toads in the environment (Bureau of Rural Sciences 2009). The initial very high level of concern leads to high expectations that environmental agencies will take action to avert the impact of toads.

This TAP acknowledges:

- there are no “magic bullets” that the Australian Government can provide that will eradicate or reduce the cane toad population across Australia
- there are competing conservation requirements for limited conservation funding generating a need to prioritise any allocation of cane toad management funds necessary to those efforts most likely to conserve biodiversity (over \$20 million has been provided by the Australian Government since 1986 to little avail)
- it is clear that, in some cases, other established invasive species, or human activity, are the cause of native species extinctions
- actions to support priority species and threatened ecological communities from the impact of cane toads across Australia must be prioritised, with priority being given to those species that would be most affected at a population level and ecosystems where multiple complex changes may occur
- there remains a need for tools to help all stakeholders at national, state/territory, regional and locals levels to effectively implement and manage cane toad impacts.

While the primary responsibility for managing established pests lies with the states and territories and landholders, all stakeholders can play significant roles in reducing the impacts of cane toads. However, in order to empower stakeholders to take actions that collectively reduce the worst impacts of cane toads it is necessary to communicate:

- the strategic approach detailed in this TAP
- the key priority species and ecosystems that need protection
- guidelines designed to enable action to be undertaken effectively
- standard operating procedures and codes of practice to ensure the humane treatment of cane toads.

State agencies and community groups have produced significant high quality communication materials relating to cane toads. These groups present this material to stakeholders and the general public through regular newsletters and media releases. A number of networks of conservation groups and researchers with an interest in cane toads already exist. These networks can form a link in a communications strategy for

this TAP including communications in regard to developments in toad control methods.

Actions	Priority and timeframe
<p>Action 3.1 Implement a one-stop-shop webpage on the DEWHA website with links to jurisdictional and stakeholder information on cane toads and including information on:</p> <ul style="list-style-type: none"> • the threat cane toads pose to biodiversity • management actions to limit this threat • guidelines for cane toad management • codes of practice and standard operating procedures • management plans for areas designated as high priority as they are developed. 	Medium, priority ongoing
<p>Action 3.2 Ensure monitoring, evaluation and reporting on cane toad management actions is maintained and communicated to stakeholders.</p>	Medium priority, ongoing
<p>Action 3.3 Ensure Australian Government fact sheets and other communications material on cane toads are current and reflect the strategy developed in this TAP.</p>	Medium priority, ongoing

Performance indicators

- Webpage on the DEWHA website holds appropriate information and linkages within 12 months of the Minister making this TAP.
- All co-funded or Australian Government funded cane toad projects include reporting of cane toad management actions, and monitoring of results, and are made available to the public within six months of the completion of the project.
- Cane toad fact sheets and other communications material are revised and made available to the public within 12 months of the Minister making this TAP.
- Threat abatement plan priorities are communicated directly to communities and stakeholders that have expressed concern and interest in cane toad control within three months of any request.

3. Duration, implementation and evaluation of the plan

3.1 Duration of the plan

This TAP reflects the fact that threat abatement will be ongoing, as there is no prospect for national eradication of cane toads.

This TAP must be reviewed by the Minister at intervals of not longer than five years. The Minister's scientific advisory committee, the Threatened Species Scientific Committee, will be provided with annual updates of actions taken under this TAP.

3. 2 Cost of the plan

Funding for TAP actions, along with a range of other responsibilities under the EPBC Act, to be undertaken by the Australian Government, is provided to the Department of the Environment, Water, Heritage and the Arts via the Australian Government budget each year. Currently, this funding sits under Outcome 1 of the Department's budget:

“The conservation and protection of Australia's terrestrial and marine biodiversity and ecosystems through supporting research, developing information, supporting natural resource management, regulating matters of national environmental significance and managing Commonwealth protected areas.”

This budget outcome is allocated to a wide variety of actions including energy efficiency measures, biodiversity conservation, the *Caring for our Country* initiative, the *Australian Biological Resources Study Strategic Plan 2007–2011*, assessment of Commonwealth-managed and all export fisheries, protection of cetaceans as well as development and implementation of individual TAPs.

Over the past two years \$2 million of funding under Caring for our Country has been allocated to both the development of this TAP and a set of projects designed to meet the Caring for our Country business plan targets on cane toads. This TAP sets a new approach to the management of cane toads that focuses on the high priority areas and high priority species as being more effective, efficient and feasible than the broad-scale approaches used in the past.

Where possible, actions under this TAP will be facilitated through existing internal budget allocations where an existing responsibility for biodiversity protection already exists (e.g. the National Reserve System). Where actions required fall outside these existing responsibilities (e.g. development and application of the prioritisation tool), actions will be funded via the Environment Protection and Biodiversity Conservation Program. It is not possible to assign costs to each element of the TAP at this time.

Investment in many of the TAP actions will be determined by the stakeholders, in particular the states and territories. It is not possible to quantify either the uptake of actions or the funding that may be provided by each of the affected jurisdictions. This will be a matter of negotiation (e.g. one tool or resource, funded solely by one jurisdiction, may be shared with other jurisdictions in return for a discounted cost to use or access a different tool or resource).

In addition to funding provided directly by the Australian Government and the jurisdictions, TAP actions are often enacted via existing intergovernmental mechanisms such as the VPC. Funding for these mechanisms is incorporated in normal organisational administrative costs and is not able to be detailed on the basis of costs of an individual TAP.

The total cost of implementation of this TAP, therefore, cannot be quantified at the time of its writing. However, the Australian Government is committed to undertake all the actions listed within the 5 year life of this plan.

This TAP provides a framework for undertaking targeted priority actions. Budgetary and other constraints may affect the achievement of the objectives of this TAP and, as knowledge changes, proposed actions may be modified over the life of the TAP. Australian Government funds may be available to implement key national environmental priorities, such as relevant actions listed in this Plan and actions identified in regional natural resource management plans.

3.3 Implementing the plan

In order to successfully implement this TAP, DEWHA will:

- implement the TAP as it applies to Commonwealth land and will act in accordance with the provisions of the plan
- maintain its strong links with state and territory agencies and with local and regional bodies that are responsible for the management of cane toads
- seek stronger coordination of national action on cane toads under the auspices of the VPC and will draw on expertise from CTAG, state and territory agencies and non-government organisations
- encourage involvement of key stakeholders and experts in cane toad related research and management.

In relation to Australian Government responsibilities, the EPBC Act requires the Director of National Parks to protect, conserve and manage biodiversity and heritage in Commonwealth reserves and conservation zones and to contribute to these factors in areas outside Commonwealth reserves and conservation zones. Collaboration between all stakeholders is required for the successful implementation of this Plan. Local governments assist in delivering state and territory priorities at a local and regional level and consequently may be involved in the management of specific assets as part of jurisdictional actions.

Research priorities for managing the impacts of cane toads should focus on identification of priority biodiversity assets at risk from the impact of cane toads; mechanisms for the protection of those assets found to be of a high priority; and preparation of appropriate tools for stakeholders to use to mitigate the negative impacts of cane toads. All research and monitoring results will be provided to stakeholders via the Department of the Environment, Water, Heritage and the Arts cane toad webpage within the timeframes set under Objective 3.

3.4 Evaluating and reviewing the plan

Section 279 of the EPBC Act provides for the review of this TAP at any time and requires that the TAP be reviewed at intervals of no longer than five years. If evidence is found that the objectives and actions recommended in the TAP need to be updated or modified to prevent species or ecological communities becoming threatened, or that the effectiveness of the TAP can be improved, it can be revised within five years of the release of this TAP. Annual reports on the implementation of the TAP will be provided to the Threatened Species Scientific Committee.

676

677

References

678

679 Boland CRJ (2004). Introduced cane toads are active nest predators and competitors of
680 rainbow bee-eaters: observational and experimental evidence, *Biological Conservation*
681 120:53–62.

682

683 Bureau of Rural Sciences (2009). *Cane Toads in Communities - Executive Report*,
684 Commonwealth of Australia.

685

686 Commonwealth of Australia (2006). *EPBC Act Policy Statement 1.1 Significant Impact*
687 *Guidelines*, available at [http://www.environment.gov.au/epbc/publications/pubs/nes-](http://www.environment.gov.au/epbc/publications/pubs/nes-guidelines.pdf)
688 [guidelines.pdf](http://www.environment.gov.au/epbc/publications/pubs/nes-guidelines.pdf) accessed November 2, 2009.

689

690 Doody JS, Roe J, Mayes P and Ishiyama L (2009). Telemetry tagging methods for some
691 freshwater reptiles, *Marine and Freshwater Research* 60:293–298.

692

693 Dubey S and Shine R (2008). Origin of the parasites of an invading species, the Australian
694 cane toad (*Bufo marinus*): are the lungworms Australian or American?, *Molecular*
695 *Ecology* 17:4418–4424

696

697 Freeland WJ (1993). *Parasites, pathogens and the impact of introduced organisms on the*
698 *balance of nature in Australia*, Pages 171–180 in *Conservation Biology in Australia*
699 *and Oceania*, edited by C. Moritz and J. Kikkawa. Chipping Norton: Surrey Beatty and
700 Sons.

701

702 Freeland WJ (2004). *An assessment of the introduced cane toad's (Bufo marinus Linnaeus)*
703 *impacts on the native Australian fauna, with particular reference to the Eastern*
704 *Kimberley region*, Unpublished Report to the WA Department of Industry and
705 Resources. Palmerston, NT: HLA-Envirosciences Pty Ltd.

706

707 Global Invasive Species Database website (2009). <http://www.issg.org/database/welcome/>
708 accessed November 2, 2009.

709

710 Griffiths AD. and McKay JL (2007). Cane toads reduce the abundance and site occupancy of
711 freshwater goannas *Varanus mertensi*, *Wildlife Research* 34:609–615.

712

713 Hagman M, Hayes R, Capon R and Shine R (2009). Alarm cues experienced by cane toad
714 tadpoles affect post-metamorphic morphology and chemical defences, *Functional*
715 *Ecology* 23:126–132.

716

717 Hagman M, Phillips BL and Shine, R (2009). Fatal attraction: adaptations to prey on native
718 frogs imperil snakes after invasion of toxic toads, *Proceedings of the Royal Society B*:
719 in press.

720

721 Hagman M, and Shine R (2009). Larval alarm pheromones as a potential control for invasive
722 cane toads (*Bufo marinus*) in tropical Australia, *Chemoecology* 19:211–217.

723

724 Kearney M, Phillips BL, Tracy CR, Christian KA, Betts G and Porter W P (2008). Modelling
725 species distributions without using species distributions: the cane toad in Australia
726 under current and future climates, *Ecography* 31: 423–434 .

727

- Letnic M, Webb JK and Shine R (2008). Invasive cane toads (*Bufo marinus*) cause mass mortality of freshwater crocodiles (*Crocodylus johnstoni*) in tropical Australia, *Biological Conservation* 141:1773–1782.
- Oakwood M (2003a). *The effect of cane toads on a marsupial carnivore, the northern quoll, Dasyurus hallucatus*. Unpublished Progress Report February 2003. Darwin: Parks Australia North.
- Oakwood M (2003b). *The effect of cane toads on a marsupial carnivore, the northern quoll, Dasyurus hallucatus*, Unpublished Progress Report August 2003. Darwin: Parks Australia North.
- O'Donnell S (2009). *Alien versus predator: taste aversion learning as a way of enhancing survival in Northern Quolls (Dasyurus hallucatus)*, Honours thesis. Sydney: School of Biological Sciences, University of Sydney.
- Peacock, T (2007). *Community on-ground cane toad control in the Kimberley - A review conducted for the Hon. David Templeman MP Minister for the Environment, Climate Change and Peel*, Invasive Animals Cooperative Research Centre website: http://www.invasiveanimals.com/downloads/Peacock_Community-toad-control-report.pdf accessed 4 Jan 2010.
- Phillips BL, Brown GP, Greenlees M, Webb JK and Shine R (2007). Rapid expansion of the cane toad (*Bufo marinus*) invasion front in tropical Australia, *Austral Ecology* 32, 169–176.
- Phillips BL, Greenlees MJ, Brown GP and Shine R (2009). Predator behaviour and morphology mediates the impact of an invasive species: cane toads and death adders in Australia, *Animal Conservation*: in press.
- Price-Rees, S., Brown GP, Shine R (2010). Are bluetongue lizards (*Tiliqua scincoides intermedia*, Scincidae) threatened by the invasion of toxic cane toads (*Bufo marinus*) through tropical Australia?, *Wildlife Research*, in press.
- Rankmore BR, Griffiths AD., Woinarski JCZ., Ganambarr BL, Taylor R, Brennan K., Firestone K and. Cardoso M (2008). Island translocation of the northern quoll *Dasyurus hallucatus* as a conservation response to the spread of the cane toad *Chaunus (Bufo) marinus* in the Northern Territory, Australia. Northern Territory Department of Natural Resources, Environment, The Arts and Sport website: http://www.nt.gov.au/nreta/wildlife/programs/quoll/pdf/quolltranslocation_final.pdf accessed 16 December 2009.
- Shannon MF, and Bayliss, P (2008). Review of the CSIRO Biological Control of Cane Toad Program to April 2008, Department of the Environment, Water, Heritage and the Arts website: <http://www.environment.gov.au/biodiversity/invasive/publications/cane-toad/pubs/cane-toad-2008.rtf>: accessed December 10 2009.
- Shine R (2009a). *The ecological impact of invasive cane toads (Bufo marinus) in Australia*, School of Biological Sciences, University of Sydney, NSW 2006, Australia.
- Shine R (2009b). Cane toad website: <http://www.canetoadsinoz.com/index.html> accessed November 2 2009.

- Shine R (2009c). University of Sydney cane toad research website
<http://www.bio.usyd.edu.au/Shinelab/cane%20toads/toads.html> accessed November 2
 2009.
- Threatened Species Scientific Committee (TSSC) (2005) *Commonwealth Listing Advice for
 The biological effects, including lethal toxic ingestion, caused by Cane Toads (Bufo
 marinus)* Department of the Environment, Water, Heritage and the Arts website:
<http://www.environment.gov.au/biodiversity/threatened/ktp/cane-toads.html> accessed
 November 2, 2009.
- Ujvari B and Madsen T (2009). Invasion of cane toads associates with a significant increase
 in mortality in a naïve Australian varanid lizard, *Herpetological Conservation*: in press.
- van Dam R, Walden D and Begg G (2002). *A preliminary risk assessment of cane toads in
 Kakadu National Park*, Supervising Scientist Report 164. Darwin: Supervising
 Scientist.
- Watson M and Woinarski J (2003a). *A preliminary assessment of impacts of cane toads on
 terrestrial vertebrate fauna in Kakadu National Park*, Unpublished Report February
 2003. Jabiru: Kakadu Research Advisory Committee.
- Watson M and Woinarski J (2003b). *Vertebrate monitoring and re-sampling in Kakadu
 National Park, 200, Project RS10*, Unpublished Report March 2003. Jabiru: Parks
 Australia.

Appendices

Appendix 1 Cane toad-free islands in Queensland which hold populations of those native species determined to be highly impacted (at the population scale) in the presence of cane toads (i.e. species listed in Table 3).

	<i>Acanthophis praelongus</i> Northern death adder	<i>Pseudechis australis</i> King brown snake	<i>Tiliqua scincoides intermedia</i> Northern blue-tongue lizard	<i>Varanus gouldii</i> Gould's goanna	<i>Varanus indicus</i> Mangrove monitor	<i>Varanus mertensi</i> Merten's water monitor	<i>Varanus panoptes</i> Yellow-spotted monitor	<i>Varanus semiremex</i> Rusty monitor	<i>Varanus timorensis</i> Spotted tree monitor	<i>Varanus tristis</i> Black-headed monitor
AKENS ISLAND				x						
BADU ISLAND	x				x				x	
BOIGU ISLAND					x					
BRIEBIE ISLAND				x	x					
CAP ISLET					x					
CARLISLE ISLAND							x			
CLIFF ISLAND NATIONAL PARK								x		
COMPIGNE ISLAND				x						
COONANGLEBAH (DUNK) ISLAND	x									
COQUET ISLAND								x		
CURTIS ISLAND		x		x						
DARNLEY ISLAND					x					
DAUAN ISLAND					x					
DENHAM ISLAND		x								
DOWAR ISLET					x					x
EBORAC ISLAND	x									
FACING ISLAND				x						
FLINDERS ISLAND				x			x			x
FRASER ISLAND				x						
FRIDAY ISLAND				x	x					
GLOUCESTER ISLAND	x	x								
HINCHINBROOK ISLAND	x						x			
HOOK ISLAND				x						x
HORN ISLAND				x						x
HUMMOCK HILL ISLAND				x						
INGRAM ISLAND								x		
KESWICK ISLAND							x			
LIZARD ISLAND				x						
LLOYD ISLAND								x		
LONG ISLAND	x									
MAER ISLAND					x					x
MAGNETIC ISLAND	x									x
MOA ISLAND				x					x	
MORETON ISLAND				x						
MORNINGTON ISLAND	x	x	x	x					x	x
NORTH KEPPEL ISLAND				x						
NORTH STRADBROKE ISLAND				x						
PRINCE OF WALES ISLAND		x		x	x					x
RESTORATION ISLAND				x						
STANLEY ISLAND					x		x			x
THURSDAY ISLAND				x						
ULUI ISLAND					x				x	x
WAEIR ISLET					x					
WARRABER ISLET									x	
WHITSUNDAY ISLAND				x						

Appendix 2 Cane toad free islands in the Northern Territory which hold populations of those native species determined to be highly impacted (at the population scale) in the presence of cane toads (i.e. species listed in Table 3).

	<i>Dasyurus hallucatus</i> Northern Quoll	<i>Acanthopis praelongus</i> Northern death adder	<i>Pseudechis australis</i> King Brown	<i>Tiliqua scincoides intermedia</i> Northern blue-tongue lizard	<i>Varanus acanthinus</i> Ridge-tailed monitor	<i>Varanus oleobopalm</i> Black-palmed monitor	<i>Varanus gouldii</i> Gould's goanna	<i>Varanus indicus</i> Mangrove monitor	<i>Varanus mertensi</i> Merten's water monitor	<i>Varanus panoptes</i> Yellow-spotted monitor	<i>Varanus scalaris</i> Spotted tree monitor	<i>Varanus timorensis</i> Spotted tree monitor	<i>Varanus tristis</i> Black-headed monitor
ASTELL ISLAND	x												
BATHURST ISLAND		x	x	x			x		x	x		x	x
BICKERTON ISLAND				x			x		x				
BROMBY ISLET					x								
CHANNEL ISLAND	x												
COTTON ISLAND													x
CROKER ISLAND										x	x		
DJEERGAREE ISLAND					x	x							
DORCHERTY ISLAND										x			
DRYSDALE ISLAND										x			
EAST VERNON ISLAND												x	
FIELD ISLAND (KARDANGARL)										x			
GROOTE EYLANDT	x	x	x	x	x	x	x	x	x	x		x	x
GULUWURU ISLAND					x	x							
ILYAUGWAMAJA ISLAND	x												
INGLIS ISLAND	x				x			x					
JIRRGARI ISLAND			x		x		x						
MARCHINBAR ISLAND	x		x		x	x			x	x			
MELVILLE ISLAND			x	x	x		x	x	x			x	x
MOORONGGA ISLAND		x		x									
MUNGWARNDUMANANJA ISLAND	x												
NORTH EAST CROCODILE ISLAND													x
NORTH EAST ISLES	x												
NORTH WEST CROCODILE ISLAND											x		
PERON ISLAND NORTH										x			
POBASSOO ISLAND	x		x		x					x			
PROBABLE ISLAND			x										
RARAGALA ISLAND			x		x	x							
VALENCIA ISLAND													
WIGRAM ISLAND			x			x							
WINCHELSEA ISLAND	x					x	x		x				
YABOOMA ISLAND								x		x	x		

Appendix 3 Cane toad-free islands in Western Australia which hold populations of those native species determined to be highly impacted (at the population scale) in the presence of cane toads (i.e. species listed in Table 3).

	<i>Dasyurus hallucatus</i> Northern Quoll	<i>Acanthopis praelongus</i> Northern death adder	<i>Pseudechis australis</i> King Brown	<i>Varanus acanthiurus</i> Ridge-tailed monitor	<i>Varanus glauerti</i> Kimberley Rock Monitor	<i>Varanus glebovalva</i> Black-palmed monitor	<i>Varanus gouldii</i> Gould's goanna
ADOLPHUS ISLAND	x						
AUGUSTUS ISLAND	x			x	x		
BATHURST ISLAND						x	
BERTHIER ISLAND	x						
BIGGE ISLAND	x	x					
BOONGAREE ISLAND	x					x	x
BYAM MARTIN ISLAND					x	x	
CAFFARELLI ISLAND	x						
CAPSTAN ISLAND	x						
CARLIA ISLAND	x		x				
CHAMPAGNY ISLAND					x		
CORNEILLE ISLAND			x	x			
FENELON ISLAND			x				
GIBBINGS ISLAND					x	x	
HEYWOOD ISLAND					x		
HIDDEN ISLAND	x		x		x	x	
IRVINE ISLAND			x				
KATERS ISLAND						x	
KOOLAN ISLAND						x	
LACHLAN ISLAND				x		x	
LONG ISLAND			x		x		
MIDDLE OSBORNE ISLAND					x		
PASCO ISLAND					x		
PURRUNGUNGKU ISLAND	x						
SAINT ANDREW ISLAND					x		
SIR FREDERICK ISLAND	x				x		
SIR GRAHAM MOORE ISLAND			x	x	x		
SOUTH WEST OSBORNE ISLAND					x	x	
SUNDAY ISLAND			x		x		
UWINS ISLAND	x				x		
WOLLASTON ISLAND	x						

Photo credits

FRONT COVER IMAGES (left to right)

Toad aggregation (Ruchira Somaweera, University of Sydney), Calling male (David Nelson, University of Sydney), Metamorph cane toad (David Nelson, University of Sydney), Slaty-grey snake with cane toad (Zig Madycki, DEWHA)

BACK COVER IMAGES (left to right)

Calling male (David Nelson, University of Sydney), Metamorph cane toad (David Nelson, University of Sydney), Cane toad in leaf litter (Damian McRae, DEWHA), Slaty-grey snake with cane toad (Zig Madycki, DEWHA)

