

Courtesy, Dr Podger 1.11.96
Can be cited if acknowledged. (24)

Consolidated lists (of Podger (1968) and Shearer and Dillon (1995 & 1996) of species of plants from which isolation of *P. cinnamomi* has been attempted directly from surface sterilised tissues of native plants growing naturally in forests and woodlands of the forest region listed in order of the Census of Western Australian Plants.

• Positive record

| Taxon | Formation | | | | | | | |
|-----------------------------------|-------------|----|------------|-----|-------------|---|----|------------|
| | Forest | | | | Woodland | | | |
| | Record | | | | Record | | | |
| | Podger 1968 | b | S & D 1995 | d | Podger 1968 | a | b | c & D 1996 |
| | *a | | c | | | | | d |
| Dennstaedtiaceae | | | | | | | | |
| <i>Pteridium esculentum</i> | 25 | 0 | 5 | 0 | | | | |
| Cycadaceae | | | | | | | | |
| • <i>Macrozamia reidlei</i> | 285 | 2 | 46 | 26 | | | 16 | 25 |
| Podocarpaceae | | | | | | | | |
| • <i>Podocarpus drouynianus</i> | 75 | 1 | | | | | | |
| Cyperaceae | | | | | | | | |
| <i>Cyathochaeta clandestina</i> | | | 2 | 0 | | | | |
| <i>Lepidosperma angustatum</i> | | | 4 | 0 | 60 | 0 | 1 | 0 |
| <i>Lepidosperma tenuis</i> | | | 2 | 0 | | | | |
| Restionaceae | | | | | | | | |
| • <i>Loxocarya cinerea</i> | | | 11 | 18 | | | | |
| <i>Hypolaena exculca</i> | | | | | 38 | 0 | | |
| <i>Hypolaena fastigata</i> | | | | | | | 1 | 0 |
| <i>Restio megalotheca</i> | | | | | 37 | 0 | | |
| Juncaceae | | | | | | | | |
| <i>Juncus planifolius</i> | | | | | 41 | 0 | | |
| Dasygongonaceae | | | | | | | | |
| <i>Calectasia cyanea</i> | | | | | 40 | 0 | | |
| <i>Dasygongon bromeliaefolius</i> | 4 | 0 | | | | | | |
| <i>Lomandra integra</i> | | | 3 | 0 | | | | |
| • <i>Lomandra odora</i> | | | 5 | 36 | | | | |
| • <i>Lomandra sonderi</i> | | | 16 | 54 | | | | |
| Xanthorrhoeaceae | | | | | | | | |
| • <i>Xanthorrhoea gracilis</i> | 7 | 0 | 149 | 38 | 18 | 0 | | |
| • <i>Xanthorrhoea preissii</i> | | | 227 | 27 | 10 | 0 | | |
| Phormaceae | | | | | | | | |
| • <i>Dianella revoluta</i> | | | 2 | 100 | | | | |
| Haemodoraceae | | | | | | | | |
| <i>Blancoa canescens</i> | | | | | 83 | 0 | | |
| <i>Conostylis aculeata</i> | | | 1 | 0 | | | | |
| <i>Conostylis candicans</i> | | | | | 31 | 0 | | |
| <i>Conostylis juncea</i> | | | | | 32 | 0 | | |
| <i>Conostylis pusilla</i> | | | 4 | 0 | | | | |
| <i>Conostylis serrulata</i> | | | 1 | 0 | | | | |
| • <i>Conostylis setigera</i> | 10 | 10 | | | | | | |
| <i>Conostylis setosa</i> | 12 | 0 | 3 | 0 | | | 3 | 0 |
| <i>Phlebocarya ciliata</i> | | | | | | | | |

*a tissue pieces tested b % tissue pieces positive c whole plants tested d % of plants positive. Podger unpublished MSc Thesis. S&D Aust. J. Bot.

| Taxon | Formation | | | | | | | |
|-----------------------------------|-------------|------------|-------------|------------|----------|----|--------|----|
| | Forest | | | | Woodland | | | |
| | Record | | Record | | Record | | Record | |
| | Podger 1968 | S & D 1995 | Podger 1968 | S & D 1995 | a | b | c | d |
| | a | b | c | d | a | b | c | d |
| Iridaceae | | | | | | | | |
| • <i>Patersonia occidentalis</i> | | | 40 | 45 | 57 | 0 | 8 | 12 |
| • <i>Patersonia rufa</i> | | | 15 | 13 | | | | |
| • <i>Patersonia umbrosa</i> | | | 13 | 15 | | | | |
| Casuarinaceae | | | | | | | | |
| • <i>Allocasuarina fraseriana</i> | 40 | 7 | 16 | 12 | | | 5 | 20 |
| Proteaceae | | | | | | | | |
| • <i>Adenanthera barbigerus</i> | | | 117 | 31 | | | | |
| <i>Adenanthera cygnorum</i> | | | | | | | 3 | 0 |
| • <i>Adenanthera obovata</i> | 82 | 12 | | | | | 3 | 66 |
| • <i>Banksia attenuata</i> | | | | | 184 | 5 | 38 | 58 |
| • <i>Banksia grandis</i> | 213 | 2 | 97 | 37 | | | 7 | 43 |
| • <i>Banksia ilicifolia</i> | | | | | 75 | 23 | 25 | 76 |
| • <i>Banksia littoralis</i> | 117 | 21 | 15 | 60 | | | | |
| • <i>Banksia menziesii</i> | | | | | 26 | 23 | 5 | 80 |
| • <i>Banksia quercifolia</i> | | | | | 16 | 37 | | |
| <i>Banksia seminuda</i> | 25 | 0 | | | | | | |
| • <i>Dryandra armata</i> | | | 1 | 100 | | | | |
| <i>Dryandra bipinnatifida</i> | | | 1 | 0 | | | | |
| • <i>Dryandra carduacea</i> | | | 18 | 22 | | | | |
| • <i>Dryandra nivea</i> | 4 | 0 | 31 | 16 | | | | |
| • <i>Dryandra sessilis</i> | 12 | 0 | 64 | 19 | | | | |
| <i>Grevillea synaphaea</i> | | | 1 | 0 | | | | |
| <i>Grevillea trifida</i> | 28 | 0 | | | | | | |
| <i>Grevillea wilsonii</i> | | | 4 | 0 | | | | |
| <i>Hakea lissocarpa</i> | | | 2 | 0 | | | | |
| • <i>Hakea ruscifolia</i> | | | 4 | 25 | | | | |
| <i>Hakea amplexicaulis</i> | | | 2 | 0 | | | | |
| <i>Hakea undulata</i> | | | 6 | 0 | | | | |
| • <i>Isopogon attenuatus</i> | 15 | 40 | | | | | | |
| <i>Isopogon dubius</i> | | | 1 | 0 | | | | |
| • <i>Isopogon formosus</i> | 36 | 19 | | | | | | |
| • <i>Isopogon sphaerocephalus</i> | 16 | 0 | 9 | 16 | | | 1 | 0 |
| • <i>Persoonia longifolia</i> | 133 | 7 | 118 | 17 | | | | |
| <i>Petrophile biloba</i> | 31 | 0 | | | | | | |
| • <i>Petrophile linearis</i> | | | 4 | 0 | | | 6 | 66 |
| <i>Petrophile obovatus</i> | | | 2 | 0 | | | | |
| • <i>Petrophile striata</i> | | | 1 | 100 | | | 1 | 14 |
| • <i>Stirlingia latifolia</i> | | | | | | | | |
| <i>Synaphea petiolaris</i> | | | 2 | 0 | | | | |
| • <i>Xylomelum occidentale</i> | 23 | 43 | | | | | 2 | 0 |
| Pittosporaceae | | | | | | | | |
| <i>Billardiera variifolia</i> | | | | | | | 3 | 0 |
| Mimosaceae | | | | | | | | |
| • <i>Acacia campylophylla</i> | | | 1 | 100 | | | | |
| <i>Acacia drummondii</i> | | | 2 | 0 | | | | |
| <i>Acacia hastulata</i> | 20 | 0 | | | | | | |
| • <i>Acacia huegelii</i> | | | | | 16 | 25 | | |

| Taxon | Formation | | | | | | | |
|-----------------------------------|-------------|------------|-------------|------------|-------------|------------|-------------|------------|
| | Forest | | | | Woodland | | | |
| | Record | | Record | | Record | | Record | |
| | Podger 1968 | S & D 1995 |
| | a | b | c | d | a | b | c | d |
| Mimosaceae (contd) | | | | | | | | |
| <i>Acacia lateriticola</i> | | | 5 | 0 | | | | |
| <i>Acacia preissiana</i> | | | 1 | 0 | | | | |
| • <i>Acacia pulchella</i> | | | 11 | 0 | | | 20 | 5 |
| <i>Acacia saligna</i> | | | | | | | 1 | 0 |
| • <i>Acacia stenoptera</i> | | | | | | | 1 | 100 |
| <i>Acacia urophylla</i> | | 1 | 0 | | | | | |
| Papilionaceae | | | | | | | | |
| • <i>Aotus ericoides</i> | | | | | 22 | 23 | | |
| • <i>Aotus passerinoides</i> | | | | | 73 | 8 | | |
| • <i>Bossiaea aquifolium</i> | 16 | 0 | 10 | 20 | | | | |
| • <i>Bossiaea eriocarpa</i> | 11 | 18 | | | | | 3 | 33 |
| • <i>Bossiaea ornata</i> | | | 4 | 75 | | | | |
| • <i>Burtonia conferta</i> | | | | | 13 | 8 | 1 | 0 |
| <i>Daviesia costata</i> | 27 | 0 | | | | | | |
| • <i>Daviesia decurrens</i> | | | | | 17 | 35 | | |
| <i>Daviesia pectinata</i> | 62 | 0 | | | | | | |
| • <i>Daviesia polyphylla</i> | | | 4 | 25 | | | | |
| • <i>Daviesia rhomboides</i> | | | 2 | 50 | | | | |
| • <i>Dillwynia uncinata</i> | 15 | 47 | | | | | | |
| • <i>Gompholobium knightianum</i> | | | 5 | 20 | | | | |
| <i>Hovea chorizemifolia</i> | | | 4 | 0 | | | | |
| • <i>Hovea elliptica</i> | 15 | 47 | | | | | | |
| <i>Hovea pungens</i> | | | | | | | | |
| <i>Jacksonia horrida</i> | | | | | | | 3 | 0 |
| • <i>Kennedia coccinea</i> | | | 1 | 100 | | | | |
| • <i>Pultenaea reticulata</i> | 26 | 9 | | | | | | |
| <i>Sphaerolobium medium</i> | | | 6 | 0 | | | | |
| <i>Viminaria juncea</i> | | | | | | | 2 | 0 |
| Rutaceae | | | | | | | | |
| <i>Boronia gracilipes</i> | 24 | 0 | | | | | | |
| <i>Boronia spathulata</i> | | | 1 | 0 | | | | |
| <i>Crowea angustifolia</i> | 20 | 0 | | | | | | |
| Tremandraceae | | | | | | | | |
| • <i>Tetrapetra viminea</i> | 60 | 25 | | | | | | |
| Euphorbiaceae | | | | | | | | |
| • <i>Amperea ericoides</i> | 24 | 12 | | | | | | |
| <i>Phyllanthus calycinus</i> | | | 9 | 0 | | | | |
| Rhamnaceae | | | | | | | | |
| • <i>Trymalium ledifolium</i> | 9 | 0 | 16 | 6 | | | | |
| Sterculiaceae | | | | | | | | |
| • <i>Lasiopetalum floribundum</i> | 21 | 38 | 2 | 0 | | | | |
| • <i>Lasiopetalum glabratum</i> | | | 2 | 100 | | | | |
| • <i>Thomasia grandiflora</i> | 168 | 29 | | | | | 35 | 43 |
| • <i>Thomasia pauciflora</i> | | | | | | | | |
| Dilleniaceae | | | | | | | | |
| • <i>Hibbertia acerosa</i> | 11 | 9 | | | | | | |
| • <i>Hibbertia amplexicaulis</i> | | | 8 | 50 | | | | |

| Taxon | Formation | | | | | | | |
|------------------------------------|-------------|------------|-------------|------------|----------|----|--------|-----|
| | Forest | | | | Woodland | | | |
| | Record | | Record | | Record | | Record | |
| | Podger 1968 | S & D 1995 | Podger 1968 | S & D 1995 | | | | |
| | a | b | c | d | a | b | c | d |
| Dilleniaceae (contd) | | | | | | | | |
| • <i>Hibbertia cunninghamii</i> | 35 | 77 | | | | | | |
| • <i>Hibbertia hypericoides</i> | | | 5 | 100 | | | 13 | 69 |
| • <i>Hibbertia linearis</i> | | | 4 | 100 | | | | |
| • <i>Hibbertia montana</i> | | | 6 | 100 | | | | |
| • <i>Hibbertia quadricola</i> | | | 1 | 100 | | | | |
| • <i>Hibbertia silvestris</i> | | | 1 | 100 | | | | |
| • <i>Hibbertia subvaginata</i> | | | | | 46 | 13 | 3 | 33 |
| • <i>Hibbertia vaginata</i> | | | | | 15 | 86 | | |
| Thymelaeaceae | | | | | | | | |
| <i>Pimelea ciliata</i> | | | 2 | 0 | | | | |
| <i>Pimelea suaveolens</i> | | | 2 | 100 | | | | |
| Myrtaceae | | | | | | | | |
| • <i>Agonis hypericifolia</i> | 47 | 2 | | | | | | |
| <i>Agonis parviceps</i> | 50 | 0 | | | | | | |
| <i>Astartea fascicularis</i> | 14 | 0 | | | | | | |
| <i>Baeckia camphorosmae</i> | | | 1 | 0 | | | | |
| • <i>Beaufortia sparsa</i> | 17 | 23 | | | | | | |
| • <i>Calytrix flavescens</i> | | | | | 34 | 23 | 1 | 0 |
| <i>Calytrix fraseri</i> | | | | | | | 3 | 0 |
| <i>Eucalyptus calophylla</i> | | | 17 | 0 | | | | |
| • <i>Eucalyptus marginata</i> | 262 | 5 | | | | | 11 | 18 |
| <i>Eucalyptus todtsiana</i> | | | | | 50 | 0 | | |
| • <i>Hypocalymma angustifolium</i> | 15 | 66 | 6 | 0 | | | 1 | 0 |
| • <i>Hypocalymma cordifolium</i> | 35 | 17 | | | | | 4 | 25 |
| • <i>Hypocalymma robustum</i> | 18 | 17 | 2 | 100 | | | 1 | 100 |
| • <i>Kunzea ericifolia</i> | | | | | | | 5 | 40 |
| • <i>Leptospermum ellipticum</i> | | | | | 27 | 48 | | |
| • <i>Melaleuca thymoides</i> | | | | | | | 21 | 71 |
| • <i>Pericalymma ellipticum</i> | | | | | | | 4 | 25 |
| <i>Scholtzia involucrata</i> | | | | | | | 2 | 0 |
| • <i>Verticordia densiflora</i> | | | | | 39 | 31 | | |
| • <i>Verticordia huegelii</i> | | | | | 33 | 6 | | |
| • <i>Verticordia plumosa</i> | | | | | 30 | 70 | | |
| Apiaceae | | | | | | | | |
| • <i>Platysace compressa</i> | 16 | 37 | 1 | 100 | | | | |
| <i>Platysace juncea</i> | | | | | 10 | 0 | | |
| <i>Xanthosia atkinsoniana</i> | | | 2 | 0 | | | | |
| Epacridaceae | | | | | | | | |
| <i>Andersonia caerulea</i> | | | | | 37 | 0 | | |
| <i>Andersonia sprengelioides</i> | | | | | 13 | 0 | | |
| • <i>Astroloma microcalyx</i> | | | 3 | 33 | | | | |
| <i>Astroloma pallidum</i> | | | 4 | 0 | | | | |
| • <i>Conostephium pendulum</i> | | | | | | | 3 | 66 |
| • <i>Leucopogon australis</i> | 55 | 38 | | | 29 | 31 | | |

| Taxon | Formation | | | | | | | |
|-------------------------------------|-------------|------------|-------------|------------|----------|----|--------|----|
| | Forest | | | | Woodland | | | |
| | Record | | Record | | Record | | Record | |
| | Podger 1968 | S & D 1995 | Podger 1968 | S & D 1995 | | | | |
| | a | b | c | d | a | b | c | d |
| Epacridaceae (contd) | | | | | | | | |
| • <i>Leucopogon capitellatus</i> | | | 5 | 100 | | | | |
| • <i>Leucopogon conostephioides</i> | | | | | | | 10 | 90 |
| <i>Leucopogon cordatus</i> | | | | | 30 | 0 | | |
| • <i>Leucopogon glabellus</i> | 63 | 6 | | | | | | |
| • <i>Leucopogon lasiostachyus</i> | | | | | 15 | 13 | | |
| • <i>Leucopogon nutans</i> | | | 11 | 45 | | | | |
| <i>Leucopogon oxycedrus</i> | | | 1 | 0 | | | | |
| <i>Leucopogon pendulus</i> | 13 | 0 | | | | | | |
| <i>Leucopogon polymorphus</i> | | | | | 11 | 0 | | |
| • <i>Leucopogon propinquus</i> | | | 5 | 40 | | | | |
| • <i>Leucopogon pulchellus</i> | 5 | 40 | | | | | | |
| • <i>Leucopogon verticillatus</i> | 20 | 40 | 6 | 66 | | | | |
| <i>Lysinema ciliatum</i> | | | | | 11 | 0 | | |
| • <i>Monotoca tamariscina</i> | 58 | 9 | | | | | | |
| <i>Sphenotoma capitatum</i> | 37 | 0 | | | | | | |
| • <i>Sphenotoma squarrosum</i> | 25 | 36 | | | | | | |
| • <i>Styphelia tenuiflora</i> | | | 12 | 66 | | | | |
| Lamiaceae | | | | | | | | |
| <i>Hemigenia incana</i> | | | | | 18 | 0 | | |
| <i>Hemiandra linearis</i> | | | | | 27 | 0 | | |
| Goodeniaceae | | | | | | | | |
| <i>Dampieria alata</i> | | | | | 50 | 0 | | |
| • <i>Dampieria linearis</i> | | | 1 | 0 | 56 | 23 | | |
| • <i>Leschenaultia biloba</i> | | | 3 | 33 | 30 | 0 | | |
| <i>Scaevola striata</i> | | | 1 | 0 | | | | |