



Department of **Biodiversity,  
Conservation and Attractions**

## Understanding sources of feral cats in Dryandra Woodland through DNA analysis

Version: 1.2

Approved by:

Last Updated: 28 February 2023

Custodian: Kym Ottewell

Review date: 24/02/2023

Version number	Date approved DD/MM/YYYY	Approved by	Brief Description
1.1	17/02/2023	Kym Ottewell	Draft report to PHCC
1.2	28/02/2023	Kym Ottewell	Final report to PHCC



**Biodiversity and  
Conservation Science**

# Understanding sources of feral cats in Dryandra Woodland through DNA analysis

Robyn Shaw and Kym Ottewell

Final report to Peel Harvey Catchment Council  
February 2023



Department of **Biodiversity,  
Conservation and Attractions**

Department of Biodiversity, Conservation and Attractions  
Locked Bag 104  
Bentley Delivery Centre WA 6983  
Phone: (08) 9219 9000  
Fax: (08) 9334 0498

[www.dbca.wa.gov.au](http://www.dbca.wa.gov.au)

© Department of Biodiversity, Conservation and Attractions on behalf of the State of Western Australia 2023  
February 2023

This work is copyright. You may download, display, print and reproduce this material in unaltered form (retaining this notice) for your personal, non-commercial use or use within your organisation. Apart from any use as permitted under the *Copyright Act 1968*, all other rights are reserved. Requests and enquiries concerning reproduction and rights should be addressed to the Department of Biodiversity, Conservation and Attractions.

ISSN [ISSN] (print)  
ISSN [ISSN] (online)  
ISBN [ISBN] (print)  
ISBN [ISBN] (online)

This report/document/publication was prepared by Robyn Shaw and Kym Ottewell

Questions regarding the use of this material should be directed to:  
Kym Ottewell, Senior Research Scientist  
Biodiversity and Conservation Science  
Department of Biodiversity, Conservation and Attractions  
Locked Bag 104  
Bentley Delivery Centre WA 6983  
Phone: (08) 9219 9086  
Email: [kym.ottewell@dbca.wa.gov.au](mailto:kym.ottewell@dbca.wa.gov.au)

The recommended reference for this publication is:  
Shaw, R. and Ottewell, K., 2023, *Understanding sources of feral cats in Dryandra Woodland through DNA analysis. Final report to the Peel Harvey Catchment Council*. Department of Biodiversity, Conservation and Attractions, Perth.

This document is available in alternative formats on request.



# Contents

Acknowledgments .....	vii
Summary .....	viii
1 Background.....	1
1.1 Dryandra Woylie/Numbat Predator Control Project.....	1
1.2 Project aims .....	2
2 Materials and Methods.....	3
2.1 Sampling location and material .....	3
2.2 DNA extraction and SNP genotyping .....	4
2.3 SNP filtering .....	4
2.4 Relatedness and parentage analysis .....	5
2.5 Population genetic structure and genetic diversity .....	5
2.6 Fine-scale genetic structure .....	6
3 Results.....	7
3.1 SNP genotyping and filtering.....	7
3.2 Relatedness and parentage analysis .....	7
3.3 Population genetic structure and genetic diversity .....	9
3.4 Fine-scale genetic structure .....	13
4 Discussion.....	15
Appendices .....	17
Glossary .....	35
References.....	37

## Appendices

Appendix 1 Sample information .....	17
Collection details for <i>Felis catus</i> samples for which DArTSeq SNP genotypes were able to be generated.....	17
Appendix 2 SNP Filtering .....	23
SNP filtering and data cleaning protocol, including filtering steps and the thresholds chosen.....	23
Appendix 3 Pairwise relatedness .....	25
Pairwise relatedness for ‘highly related pairs’ (where relatedness $\geq 0.2$ ). .....	25
Appendix 4 Genetic diversity estimates 2016 - 2021 .....	33
Estimates of genetic diversity parameters over sampling years. ....	33

## Figures

Figure 1 Map of the study area, including spatial groupings and the number of individuals (n) within each group.....	3
Figure 2 Pairwise relatedness patterns across the study landscape, with circles representing the number of 'highly related pairs' of individuals within each group, and lines representing the number of 'highly related pairs' among groups. ....	8
Figure 3 Map of the study region with sample locations and the first two axes of the Principal Coordinate Analysis (for the full data set and the spatially trimmed data set), with colours representing the spatial groupings of individuals.....	10
Figure 4 Cross entropy plot from the TESS3 analysis, showing K values one through to six.....	10
Figure 5 Genetic summary statistics and standard error ( $A_r$ , $H_e$ and $H_o$ ), or parametric confidence intervals ( $N_e$ ) across each year for which sampling sizes were greater than or equal to 10 individuals. ....	12
Figure 6 Mean genetic diversity (internal heterozygosity, $P_{Ht}$ ) for each spatial grouping, interpolated across the study area, with grey points and labels representing the sample sizes used in each group average. ....	13
Figure 7 Correlograms for the total data set, and for females and males, showing fine-scale spatial genetic structure ( $r$ ) for increasing distance classes. Error bars represent 95% bootstrap confidence intervals and significant fine-scale spatial genetic structure is detected when confidence intervals do not overlap zero.....	14

## Tables

Table 1 Filters and sample sizes for each data set, where all data sets underwent base line filters and additional filters are listed for specific analyses.....	4
Table 2 Individuals identified as full-siblings (i.e., same mother and father), potentially from the same litter, with the spatial group the individual was sampled from in parentheses. ....	9
Table 3 Parentage assignment, with the spatial group the individual was sampled from in parentheses. ....	9
Table 4 Genetic summary statistics for the total population across all years. ....	11

## Acknowledgments

We thank Peter Lacey, Marissah Kruger, Brian Macmahon and Deanna Rasmussen (DBCA) for coordinating access to and preparation of the samples and metadata. Tissue samples were provided by DBCA, local landholders and community, local feral animal contractors, supported by Peel-Harvey Catchment Council. Funding for the genetics analysis and reporting was provided through PHCC's Dryandra Woonta project supported by the Australian Government.

# Summary

This report presents findings from genetic analysis of feral cat populations located in Dryandra Woodland and surrounds, in the wheatbelt of Western Australia.

Key results include:

- Of 176 tissue samples sent for genomic sequencing (DArTseq), high quality genotypes were obtained for 135 individuals.
- Overall, there was low genetic structure across the study area suggesting feral cats have a high capacity for dispersal in this landscape.
- High relatedness was detected within cat populations to the north and east of the study area suggesting a stable population structure in these areas, representing putative source populations.
- Conversely, individuals within the southern forest blocks were largely unrelated and there was evidence of movement of related individuals between other areas and this population, suggestive of sink dynamics.
- Positive relatedness amongst individual cats was detected at up to 20 km suggesting this is the scale of the genetic neighborhood of cats at Dryandra.
- Of years with sufficient numbers of samples for comparison (2016, 2017, 2018), genetic diversity showed a declining trend in 2018, compared to 2016 and 2017. Effective population size also declined in 2017 and 2018 compared to 2016.
- Trends in genetic diversity and effective population size suggest that feral cat control has resulted in population contraction in the study area.

# 1 Background

Feral cats (*Felis catus*) are recognised as one of the most harmful invasive species globally (Doherty et al., 2016). Termed ‘feral’ when living independently of humans in the wild, feral cats have a substantial negative impact on native wildlife through predation, competition, and disease transmission (Doherty et al., 2017). They have been implicated in the severe population decline and even extinction of many species (Doherty et al., 2016). Feral cats are also a major driver of mammal declines in Australia, particularly those species that fall within the size range of their preferred prey, known as the ‘critical weight range’ (35 g to 5.5 kg) (Burbidge & McKenzie, 1989). Estimates suggest that there are between 2.1 to 6.3 million feral cats in total across 99.8% of Australia's land area, including both natural and human modified environments (Legge et al., 2017). Thus, effective conservation management strategies are needed to mitigate the impact of feral cats on native wildlife, particularly to maintain viable populations of ‘critical weight range’ threatened Australian mammals.

Genetic data can provide valuable information about the history and ecology of invasive species (Le Roux & Wiecezorek, 2009). For example, previous studies in Australia reveal that feral cats likely originated from Europe following human colonisation (Koch et al., 2015; Spencer et al., 2016). Feral cats show high levels of genetic diversity and gene flow across Australia, with limited population genetic structure, suggesting they have a high capacity for dispersal (Spencer et al., 2016). To date, studies using genetics to understand feral cat populations have mainly used mitochondrial and microsatellite genetic markers. However, Single Nucleotide Polymorphisms (SNPs) obtained via genomic sequencing can provide increased resolution to detect subtle patterns of population genetic structure and relatedness (Fischer et al., 2017; Telfer et al., 2015), and therefore may yield additional insights into the dynamics of feral cat populations.

Genetic analyses can help guide the design of management and eradication programs through identifying patterns of genetic structure, gene flow, diversity and relatedness (Koch et al., 2020). This information can highlight areas of the landscape with high connectivity, indicating possible migration routes, as well as potential source populations when new areas are colonised. For example, a study on Christmas Island found that feral cats were able to disperse and breed successfully over the entire island (Koch et al., 2020). Efforts targeting the population as a whole reduced the effective population size over a five-year period. Another study on Dirk Hartog Island, Western Australia, found that the feral cat population on the island was likely reproductively isolated from the mainland, making it feasible to eradicate the feral cat population long-term (Koch et al., 2014).

## 1.1 Dryandra Woylie/Numbat Predator Control Project

The Dryandra Woodland National Park (comprising of Lol Gray, Highbury and Montague State Forests) is located 160 kilometres south-east of Perth in the Department of Biodiversity, Conservation and Attractions's (DBCA) Wheatbelt Region. It comprises 17 discrete blocks scattered across a north-south distance of about 50 kilometres, separated and contained by large areas of agricultural land, where edge effects and fragmentation exacerbate impacts of introduced species. Dryandra Woodland represents the largest and most diverse remnants of native vegetation in

the central Western Australian wheatbelt. It is home to over 800 native plant species and 24 mammal, 41 reptile, 8 frog and 98 bird species (DEC, 2011).

Woylies and numbats are listed as critically endangered and endangered, respectively, under Western Australia's Biodiversity Conservation Act 2016. They are also listed as endangered under the Commonwealth's Environmental Protection and Biodiversity Act 1999. Dryandra Woodland is one of only two sites containing wild populations of these species and the localised extinction of either species would result in an irreparable loss of genetic diversity of the species. Prior to feral cat control (detailed below) there was strong evidence that feral cats represented the primary predator of both woylies and numbats in the Woodland, particularly given fox numbers had been reduced by management (Marlow et al., 2015).

The Dryandra Woylie/Numbat Predator Control Project commenced in April 2015 to conserve populations of numbat and woylie in the Dryandra Woodlands through on-ground feral predator management. The Project is an integrated feral cat-fox control program at Dryandra Woodland and adjoining private land. This project aims to recover the wild populations of the woylie and numbat that occur there.

The Project builds upon existing operational management and associated research projects within DBCA estate, including the Western Shield program, protecting highly susceptible fauna from fox and cat predation. This project acknowledges that baiting on conservation estate alone may not be effective in mitigating feral cat numbers and subsequent predation, and therefore combines multiple efforts and tools to control introduced predators. The active participation of neighbours to control feral cats and foxes on their property is needed to increase the effectiveness of the Department's Western Shield program and is critical for the long-term viability of these and other fauna species in Dryandra Woodland. The project targeted cats and foxes through an integrated approach of baiting on reserve, trapping and shooting programs (DBCA, 2017).

The Peel-Harvey Catchment Council (PHCC) partnered with the DBCA Wheatbelt Region and Project Numbat in April 2017 through the Farmers for Fauna project to assist landholders adjacent to Dryandra Woodland to achieve effective off-reserve feral cat and fox control to help prevent invasion and re-invasion into the woodland. The involvement and support of the community, particularly adjacent landowners and managers are important to the conservation of Dryandra Woodland's biodiversity. Their involvement in off-reserve activities increases the effectiveness of integrated feral cat and fox management within Dryandra Woodland. As part of these activities, DNA samples are collected from humanely euthanized animals to contribute to research on their population dynamics.

## 1.2 Project aims

This project involves undertaking a SNP-based genetic analysis of samples collected following cat control activities, with the aim to:


- Investigate potential source populations of feral cats in Dryandra.
- Explore patterns of relatedness between Dryandra and potential source populations.
- Explore genetic evidence for effectiveness of cat control.
- Provide information that can help adapt and focus control efforts to manage the threat of feral cats in this landscape.

## 2 Materials and Methods

### 2.1 Sampling location and material

Feral cat control was undertaken by DBCA Wheatbelt Regional staff within Dryandra Woodlands. In addition, feral cat control off-reserve activities were undertaken by Dryandra Woodland's neighbouring landholders and managers, with support from PHCC. Ear notches were obtained from humanely euthanised individual cats and stored in 70-100% ethanol or 20% DMSO prior to laboratory analyses. Tissue samples were also collected from feral cat carcasses killed by car strikes or suspected baiting deaths found within Dryandra Woodlands and surrounding neighbouring properties (see Appendix 1 for sample information). A total of 176 tissue samples were collected from animals in Dryandra woodland and surrounding townships and agricultural properties (Figure 1) between December 2015 and November 2021.

Samples were grouped spatially based on natural breaks in the landscape and major townsites for visualisation purposes and to enable the comparison of relatedness and parentage estimates among Dryandra and surrounds and potential source populations (Figure 1). Individuals for which coordinate information was missing were removed, as analyses relied on accurate spatial data.



*Figure 1 Map of the study area, including spatial groupings and the number of individuals (n) within each group.*

## 2.2 DNA extraction and SNP genotyping

Genomic DNA was extracted from tissue samples using a standard ‘salting out’ protocol (Sunnucks & Hales, 1996) with the addition of 3  $\mu$ L 10 mg/mL RNase to the TNES buffer to remove RNA contamination. A total of 176 tissue samples were sent to Diversity Arrays Technology Pty Ltd (DART), for library preparation, sequencing, quality control and SNP genotyping (DARTseq). DARTseq<sup>TM</sup> uses enzyme digestion to prepare samples for reduced representation sequencing on an Illumina HiSeq2500, followed by read assembly, quality control and SNP calling through DART’s proprietary software (Cruz et al., 2013; Kilian et al., 2012; Melville et al., 2017).

## 2.3 SNP filtering

SNP datasets contain a large volume of genetic markers, for which individual testing and evaluation is not feasible. Therefore, filtering these data based on sensible quality control thresholds is essential to reduce noise, artefacts and errors that can be introduced in the laboratory or through genotyping error (O’Leary et al., 2018). We followed filtering protocols outlined in Shaw et al., (2022) to rigorously filter the raw SNP dataset provided by DART, with the goal to obtain a set of genetic markers that accurately represented independent loci. Thresholds were determined by visualising the raw data in R version 4.1.2 (R Core Team, 2022). Full details about the SNP filtering and data cleaning protocol can be found in Table 1 and Appendix 2.

*Table 1 Filters and sample sizes for each dataset, where all datasets underwent base line filters and additional filters are listed for specific analyses.*

Dataset	Analyses	Filters	Sample sizes
Total (baseline)	All (total, baseline dataset)	Call rate (individual $\geq 0.55$ , locus $\geq 0.9$ ) Read count ( $\geq 20$ and $\leq 150$ ) Repeatability average ( $\geq 0.95$ ) Minor allele frequency ( $\geq 0.025$ ) Secondaries (1 SNP per sequence) Sex-linked SNPs (removed) Linkage disequilibrium ( $\geq 0.5$ ) Missing coordinate data (individuals removed)	Individuals = 135 SNPs = 3,241
A	Relatedness	No additional filters	Individuals = 135 SNPs = 3,241
B	Parentage	Locus call rate (increased to $\geq 0.95$ ) Minor allele frequency (increased to $\geq 0.3$ ) Hardy Weinberg Equilibrium (loci that do not conform to HWE removed)	Individuals = 135 SNPs = 387
C	Population genetic structure	Highly related individuals (removed) Spatial trimming (1 individual/500 m radius)	Individuals = 32 SNPs = 3,241

Dataset	Analyses	Filters	Sample sizes
D	Genetic diversity statistics Internal heterozygosity Fine-scale genetic structure	Highly related individuals (removed) Hardy Weinberg Equilibrium (loci that do not conform to HWE removed)	Individuals = 72 SNPs = 3,105
E	Temporal genetic diversity statistics Effective population size	Highly related individuals (removed) Hardy Weinberg Equilibrium (loci that do not conform to HWE removed) Subset to years with large enough sample sizes (ten or more individuals = 2016, 2017 and 2018)	Individuals = 59 SNPs = 3,105

## 2.4 Relatedness and parentage analysis

We calculated pairwise relatedness (Wang, 2002) in the R package related (Pew et al., 2015), using dataset A (Table 1). Pairs of individuals with a relatedness estimate of  $\geq 0.2$  were identified as ‘highly related’, as these represent plausible values for half-siblings through to parent-offspring relationships. We then summarised these findings spatially, by visualising the number of ‘highly related pairs’ within and between spatial groups, to explore potential source populations into Dryandra.

To go beyond broad estimates of relatedness, the R package SEQUOIA (Huisman, 2017) was used to assign parentage and identify full sibling clusters within dataset B (Table 1). SEQUOIA is a maximum likelihood method that combines genetic data with demographic information (priors). Here, the sex of individuals (if known) and the minimum and maximum potential year of birth were used as priors in the analysis. The maximum year of birth was taken as the year of collection. We assumed feral cats live an average of five years, so the minimum birth year was taken to be five years before the collection year. We reduced the dataset down to a more informative set of SNPs (387), by filtering for Hardy Weinberg Equilibrium, increasing the call rate to 95% and increasing the minor allele frequency to 0.3, as recommended by Huisman (2017). Again, parentage and full-sibling assignment was summarised in relation to spatial groupings to determine whether parents of Dryandra individuals could be identified in surrounding source locations.

After summarising relatedness and parentage results, we removed one individual from each of the ‘highly related pairs’, as close relatives can bias population genetic analyses (Wang, 2018). This resulted in 63 individuals being removed from the dataset (Table 1).

## 2.5 Population genetic structure and genetic diversity

Population genetic structure was investigated to identify any landscape elements that may act as barriers or limit cat movement and to evaluate the potential for assigning migrant individuals in Dryandra to original source populations. To achieve this, the dataset was first trimmed spatially to include just one sample per 1 km<sup>2</sup> (dataset C, Table 1), creating an even distribution across the landscape to reduce bias associated with heavily sampled locations. Next, a Principal Coordinate

Analysis (PCoA) was performed in the R package *dartr* (Gruber et al., 2018), to determine if there were natural genetic clusters (representing ‘populations’) in the data. We then used the TESS3 algorithm in the R package *tess3r* (Caye & Francois, 2016) to estimate individual ancestry coefficients, while also taking geographic information into account. We tested for one through to six potential populations (K) with 50 repetitions for each value and a maximum of 200 optimisation iterations. We used the default settings for the remaining parameters and masked 10% of the data to use for the cross-validation. If population genetic structure is present, individuals will fall within K population clusters, as indicated by a plateau, or change in slope in the cross-entropy criterion (based on the best performing run with the lowest root mean squared error).

Genetic diversity summary statistics were then calculated within any population/s detected in the above *tess3r* analysis, with the HWE filtered dataset D (Table 1, Appendix 2). Observed heterozygosity ( $H_o$ ), expected heterozygosity ( $H_e$ ) and inbreeding coefficient ( $F$ ) were calculated with the R package *adegenet* (Jombart, 2008), while allelic richness (AR) was calculated in *PopGenReport* (Adamack & Gruber, 2014). Summary statistics were also calculated temporally for years with ten or more individuals (considered here as an adequate sample size for such analyses), which included 2016, 2017 and 2018 (dataset E, Table 1), to investigate whether there were substantial changes in genetic diversity with ongoing cat control. Effective population size ( $N_e$ ) was also estimated using the Linkage Disequilibrium method for each of these years (dataset E, Table 1), using *NeEstimator V2.1* (Do et al., 2014), implemented in the *dartr* package (Gruber et al., 2018).  $N_e$  estimates were reported alongside parametric confidence intervals, estimated with chi-square approximation (Waples, 2006).

Additionally, genetic diversity was estimated at the individual level with the R function *GENHET* (Coulon, 2010), which calculates the proportion of heterozygous loci for each individual (PHt) (Aparicio et al., 2006). We calculated this for all individuals in dataset D and took the average value within spatial groups, interpolating PHt across the landscape, using an inverse distance weighting scheme, to show areas of high or low genetic diversity.

## 2.6 Fine-scale genetic structure

Fine-scale patterns of genetic structure (at the individual level) were explored to understand any limitations to cat movement using spatial autocorrelation analysis of multilocus SNP genotypes for dataset D (Table 1), following methods outlined in Banks & Peakall, (2012), Double et al., (2005) and Smouse & Peakall (1999). The autocorrelation coefficient,  $r$ , describes the genetic similarity between all individuals within specified geographic distance classes and was calculated using the R package *dartr* (Gruber et al., 2018). The  $r$  coefficient was calculated for distance classes of increasing size, for all individuals and separately for females and males, to test for differences in dispersal between the sexes. The distance at which  $r$  is no longer significantly positive (where confidence intervals overlap zero) approximates the extent of positive genetic structure (Double et al., 2005). Statistical significance was tested by generating 95% bootstrap confidence intervals using 1000 bootstrap samples.

## 3 Results


### 3.1 SNP genotyping and filtering

Of the 176 samples sent to DArT, 20 samples failed to sequence. Library construction was challenging and required multiple attempts for many samples. Additionally, the composition of several samples was dominated by microbial contamination, likely where tissue samples were obtained from animals that had experienced field decomposition and degradation prior to sampling. The raw DArT dataset contained 12,098 SNPs for 156 individuals, although a further 15 individuals were removed during filtering due to a large amount of missing data and a further six individuals were removed due to missing coordinate information. After filtering, the total (baseline) dataset included 3,241 independent SNP loci for 135 individuals (Table 1). This dataset was filtered further for specific analyses, as described in Table 1 (with further detail about each specific filter provided in Appendix 2).

### 3.2 Relatedness and parentage analysis

There were 316 pairs of 'highly related individuals' ( $r \geq 0.2$ ) in the dataset (Figure 2, Appendix 3), with most of these occurring in the northeast of the study area. Of the two Groups with the highest sample sizes, 21% of pairwise individual comparisons in Group 9 were 'highly related', compared to only 0.4% in Group 2. For all remaining Groups, the percentage varied between 0 to 33.3% (although some of these Groups had sample sizes as low as three individuals). Thus, it is important to note that these results do not account for sampling effort.

To infer movement of cats between patches, we identified between 1 – 41 'highly related pairs' shared among spatial Groups. Related individuals were found between Group 2 and five different potential source populations. The number of other groups sharing related pairs was lower, ranging from 0 – 3.



*Figure 2 Pairwise relatedness patterns across the study landscape, with circles representing the number of 'highly related pairs' of individuals within each group, and line weight representing the number of 'highly related pairs' among groups.*

Parentage analysis revealed a similar trend, with more parent-offspring relationships and full-sibling clusters assigned in the northeast (Group 9) and east (Group 3) of the study area compared to within Group 2 (Tables 1 – 2). Of the 11 full-sibling clusters (representing potential litters, Table 1), eight were from Group 9. The remaining sibling clusters were made up of individuals from Group 3 and 10. Two sibling clusters suggest dispersal of individuals within litters between Group 9 and 10, and between Group 9 and Group 2. Parents (mothers and/or fathers) were assigned for 12 individuals in Group 3 and 9 (Table 2). All parents were from the same location (Group) as their offspring, with the exception of one father-offspring pair from Group 10 and 9, respectively, suggesting Group 10 was the potential source of juvenile recruitment into Group 9.

*Table 2 Individuals identified as full-siblings (i.e., same mother and father), potentially from the same litter, with the spatial group the individual was sampled from in parentheses.*

Cluster	Offspring 1	Offspring 2	Offspring 3	Offspring 4	Offspring 5
1	T14328 (9)	T14331 (9)	T14422 (9)	NA	NA
2	T14374 (9)	T14373 (9)	NA	NA	NA
3	T14268 (9)	T14376 (9)	T14375 (9)	T14409 (9)	T14336 (9)
4	T14278 (9)	T14387 (9)	T14424 (9)	NA	NA
5	T14400 (3)	T14402 (3)	T14406 (3)	T14399 (3)	NA
6	T14300 (10)	T14301 (10)	T14329 (9)	T14341 (9)	T14389 (9)
7	T14374 (9)	T14380 (9)	T14373 (9)	NA	NA
8	T14376 (9)	T14409 (9)	NA	NA	NA
9	T14279 (9)	T14424 (9)	NA	NA	NA
10	T14278 (9)	T14387 (9)	NA	NA	NA
11	T14314 (2)	T14336 (9)	NA	NA	NA

*Table 3 Parentage assignment, with the spatial group the individual was sampled from in parentheses.*

id	Mother	Father
T14387 (9)	NA	T14302 (10)
T14328 (9)	NA	T14326 (9)
T14424 (9)	NA	T14326 (9)
T14326 (9)	NA	T14331 (9)
T14380 (9)	T14325 (9)	NA
T14382 (9)	T14325 (9)	T14383 (9)
T14280 (9)	T14390 (9)	T14330 (9)
T14398 (3)	T14395 (3)	NA
T14400 (3)	T14396 (3)	NA
T14402 (3)	T14396 (3)	NA
T14406 (3)	T14396 (3)	NA
T14399 (3)	T14396 (3)	NA

### 3.3 Population genetic structure and genetic diversity

The first two axes of the PCoA cumulatively explained 6.2% or 9.5% of the variation in the dataset, depending on whether the full or the spatially trimmed dataset was used (Figure 3). In both cases, it did not reveal any genetic clustering of samples indicative of population genetic structure resulting from barriers to animal movement. Likewise, the cross-entropy score for the TESS3 analysis of both datasets decreased linearly for increasing values of K (numbers of populations), suggesting that the samples in this study form one genetic population (Figure 4).




Figure 3 Map of the study region with sample locations and the first two axes of the Principal Coordinate Analysis (for the full dataset and the spatially trimmed dataset), with colours representing the spatial groupings of individuals.




Figure 4 Cross entropy plot from the TESS3 analysis, showing K values one through to six.

The mean allelic richness for the population was  $1.997 \pm 0.006$ , with lower observed ( $0.199 \pm 0.002$ ) than expected heterozygosity ( $0.236 \pm 0.003$ ), potentially indicating the presence of some inbreeding ( $F = 0.144 \pm 0.003$ , Table 3). These values were fairly consistent across 2016 to 2018, although with a slight decrease in 2018 (Figure 5). Effective population size was higher in 2016 ( $N_e = 244$ , CIs = 227 – 263.7) than in later years where confidence intervals overlapped (2017:  $N_e = 153.2$ , CIs = 148.6 – 158.1; 2018:  $N_e = 133.6$ , CIs = 117.6 – 154.5; Figure 5). Sample sizes were lower in years 2019 – 2021 but indicated that  $N_e$  also remained low in 2021 (Appendix 4). Spatial interpolation of mean internal heterozygosity (Figure 6) showed that genetic diversity was highest in the northeast, and generally decreased towards the west of the study area.

*Table 4 Genetic summary statistics for the total population across all years.*

<b>Metric</b>	<b>Mean</b>	<b>Standard error</b>
<b>Observed heterozygosity (Ho)</b>	0.199	$2.16 \times 10^3$
<b>Expected heterozygosity (He)</b>	0.236	$2.49 \times 10^3$
<b>Inbreeding coefficient (F)</b>	0.144	$3.28 \times 10^3$
<b>Allelic richness (AR)</b>	1.997	$0.59 \times 10^3$




Figure 5 Genetic summary statistics and standard error ( $A_r$ ,  $H_e$  and  $H_o$ ), or parametric confidence intervals ( $N_e$ ) across each year for which sampling sizes were greater than or equal to 10 individuals.




Figure 6 Mean genetic diversity (internal heterozygosity,  $PHt$ ) for each spatial grouping, interpolated across the study area, with grey points and labels representing the sample sizes used in each group average.

### 3.4 Fine-scale genetic structure

Multi-locus genetic spatial autocorrelation detected significant fine-scale spatial genetic structure up to 20 km, suggesting that this is the scale of the genetic neighbourhood of individuals in the study area (Figure 7).

There was no significant difference in spatial genetic structure between the sexes, meaning that we did not detect any signature of sex-biased dispersal. However, this could potentially be due to the low female sample sizes resulting in large confidence intervals in this analysis.




Figure 7 Correlograms for the total dataset, and for females and males, showing fine-scale spatial genetic structure ( $r$ ) for increasing distance classes. Error bars represent 95% bootstrap confidence intervals and significant fine-scale spatial genetic structure is detected when confidence intervals do not overlap zero.

## 4 Discussion

Genetic analysis can provide insight into population dynamics of invasive species, in particular by inferring population genetic structure and the spatial scale of dispersal, as well as source/sink dynamics (Burgess et al., 2022; Koch et al., 2020; Watson et al., 2021). Insight into these aspects of invasive species biology can assist managers to design and effectively apply control methods. Here, we used high resolution SNP genotyping to assess patterns of relatedness and dispersal in the feral cat population captured in the Dryandra woodland and surrounding areas, in the wheatbelt region of Western Australia.

Overall, we found there was no detectable population genetic structure in the feral cat population in the studied landscape, suggesting that cats are moving freely and there are no quantifiable barriers to dispersal. This is in line with previous genetic studies on feral cats in Australia, which have also failed to find genetic structure at the scale of islands (Koch et al., 2020), marshes and surrounding uplands (Cowen et al., 2019), to much of the Australian continent (Spencer et al., 2016; Koch et al., 2015). Taken together, these genetic results indicate feral cats have a very high capacity for dispersal which does not appear to be influenced by the spatial configuration of agricultural versus woodland areas, which is likely a limiting factor for dispersal in other native species.

Whilst we found no strong structure at the landscape-scale, analysis of patterns of genetic relatedness amongst individual cats indicated the presence of low, but significant, fine-scale spatial genetic structure up to 20 km (i.e., cats are positively related over this distance). This result can be interpreted as defining the genetic neighbourhood of *Felis catus* in the study region. The genetic neighbourhood area ( $N_a$ ) spans the area where animals could mate randomly and is related to the effective size of a population (Wright, 1969). Male-biased dispersal has been reported for feral cats in the Fortescue Marsh (Cowen et al., 2019), however, sample sizes in our study were insufficient to detect a significant effect of dispersal differences between male and female cats, with spatial autocorrelation results being similar among the sexes.

As was observed in Cowen et al., (2019), the high dispersal capacity of feral cats (i.e., leading to low genetic structure) prohibits the use of population assignment methods to explicitly detect source-sink population dynamics in the current study. Nevertheless, qualitative assessment of patterns of genetic relatedness, particularly the distribution of siblings and parent-offspring pairs amongst spatial groups, and genetic diversity allows some inference on putative source populations in the region. Here, we found that feral cats located towards the north and east of the study area (especially Group 9, located in the outlying blocks of the Montague State forest) comprised high numbers of related individuals suggestive of a stable population structure and a putative source of cats moving into the surrounding landscape. Multiple sibling clusters (families) were detected in Groups 9 and 10 and small family groups dominated Group 3 located on agricultural properties southwest of Cuballing. There was evidence of dispersal of one half of a sibling pair dispersing between Group 9 and Group 2, to the south of the area. Group 2 located in the southern forest blocks had few internally-related individuals, but many related individuals between groups. This pattern is suggestive of a disrupted population structure and potentially represents a population sink. Even though we cannot be confident in the

direction of movement of animals, the lower overall relatedness and lower genetic diversity of this southern population of cats supports this observation. Higher genetic diversity in the north and eastern populations suggest a larger population size in this region and, given the proximity to human settlements, that these areas may be sources. Further sampling of cat populations closer to town infrastructure (e.g. rubbish tips) may also help illuminate the role of 'urbanised' populations of cats and whether they are contributing as sources.

Coordinated feral cat control in the Dryandra woodland has been ongoing since April 2017. Although sample sizes varied between years, we found that genetic diversity was similar across years between 2016 and 2018, although with a declining trend observed in 2018. Effective population size ( $N_e$ ) also showed a declining trend and was significantly lower in both 2017 and 2018, compared to 2016. While sample sizes precluded analysis for some later years, we found that  $N_e$  remained low in 2021. We can infer from these results that the control program has reduced the overall feral cat population in the area, and the decrease in effective population size is indicative of this bottleneck (Hedrick, 2011). Additional information on changes in cat abundance, e.g. from camera traps or track counts following an appropriate statistical design, would be beneficial in corroborating these results and confirming an overall reduction in cat numbers due to control methods, rather than a disruption in population dynamics due to other factors that may be unrelated to cat control.

As indicated in Cowen et al., (2019), the use of genetic monitoring to measure the impact and effectiveness of invasive species control programs has potential but ensuring adequate sample sizes over a long enough time-frame, and a large enough scale is critical to the success of such studies (Cowen et al., 2019). For this reason, it is recommended that feral cat samples continue to be collected and stored appropriately for future genetic analysis. Genetic monitoring can complement other methodologies estimating abundance and/or occupancy in demonstrating the success of control programs.

Given the challenge of working with high dispersal species such as the feral cat, we nevertheless were able to find qualitative evidence of potential source populations, and an estimate of the distance individuals are dispersing when establishing new territories across the study area (20 km). Feral cat and fox control seems to have been most effective in the southern forest blocks and future efforts may need to specifically target cat populations in the north and eastern blocks to reduce overall pressure in the area (while maintaining southern control efforts). Additional sampling around townships may assist in determining these as a source of cats moving into woodland areas. Nonetheless, information on spatial genetic patterns in the feral cat population in Dryandra Woodland and surrounding areas demonstrates the importance of coordinated efforts amongst conservation agencies, local Natural Resource Management groups and private landholders in implementing landscape-scale cat management in this system.

# Appendices

## Appendix 1 Sample information

Collection details for *Felis catus* samples for which DArTSeq SNP genotypes were able to be generated (i.e., excluding information for 20 samples that failed to sequence).

Tissue ID	Location	Date	Sex	Latitude	Longitude	ID Tag
T14258	West Cuballing	1/1/2021	NA	-32.809	117.111	010121U1
T14259	West Cuballing	1/1/2021	NA	-32.809	117.111	010121U2
T14260	West Cuballing	1/1/2021	NA	-32.809	117.111	010121U3
T14261	North of Montague	1/3/2016	F	-32.724	117.111	010316F1
T14262	South of Lol Gray	1/5/2018	NA	-32.819	117.001	010518U1
T14263	West Yornanning	1/6/2017	M	-32.753	117.028	010617M1
T14264	South of Yornaning	1/10/2020	F	-32.749	117.169	011020F1
T14265	South of Yornaning	1/10/2020	F	-32.749	117.169	011020F2
T14266	West Cuballing	2/10/2017	M	-32.809	117.111	021017M1
T14267	Narrogin	3/2/2020	M	-32.929	117.155	030220M1
T14268	West Yornanning	4/6/2017	M	-32.753	117.028	040617M1
T14269	West Cuballing	4/11/2018	NA	-32.809	117.111	041118U1
T14270	West Cuballing	4/11/2018	NA	-32.809	117.111	041118U2
T14273	West Yornanning	5/6/2017	M	-32.753	117.028	050617M1
T14274	West of Contine	5/9/2017	NA	-32.853	116.886	050917U1
T14275	Narrogin	5/10/2016	M	-32.929	117.155	051016M1
T14276	West Cuballing	6/4/2018	NA	-32.809	117.111	060418U1
T14278	West Yornanning	6/6/2017	F	-32.753	117.028	060617F1
T14279	West Yornanning	6/6/2017	M	-32.753	117.028	060617M1
T14280	West Yornanning	6/6/2017	M	-32.753	117.028	060617M2
T14281	West Yornanning	6/6/2017	M	-32.753	117.028	060617M3
T14283	Forestry Road	6/7/2017	M	-32.719	117.041	060717M1
T14284	Narrogin	7/7/2016	M	-32.929	117.155	070716M1

<b>Tissue ID</b>	<b>Location</b>	<b>Date</b>	<b>Sex</b>	<b>Latitude</b>	<b>Longitude</b>	<b>ID Tag</b>
<b>T14285</b>	West Cuballing	7/9/2021	NA	-32.809	117.111	070921U1
<b>T14286</b>	NA	7/11/2017	M	NA	NA	071117M1
<b>T14287</b>	NA	7/11/2017	M	NA	NA	071117M2
<b>T14288</b>	NA	7/11/2017	M	NA	NA	071117M3
<b>T14289</b>	NA	7/11/2017	M	NA	NA	071117M4
<b>T14290</b>	NA	7/11/2017	M	NA	NA	071117M5
<b>T14291</b>	NA	7/11/2017	M	NA	NA	071117M6
<b>T14292</b>	West Cuballing	8/5/2020	NA	-32.809	117.111	080520U1
<b>T14294</b>	South of Lol Gray	8/8/2016	F	-32.819	117.001	080816F1
<b>T14295</b>	East of forest blocks	8/8/2016	M	-32.818	116.804	080816M1
<b>T14296</b>	West Yornanning	8/8/2019	F	-32.753	117.028	080819F1
<b>T14297</b>	West Yornanning	8/8/2019	NA	-32.753	117.028	080819U1
<b>T14298</b>	Narrogin	8/10/2018	M	-32.929	117.155	081018M1
<b>T14300</b>	North of Lol Gray	9/6/2017	F	-32.712	116.958	090617F1
<b>T14301</b>	North of Lol Gray	9/6/2017	F	-32.712	116.958	090617F2
<b>T14302</b>	North of Lol Gray	9/6/2017	M	-32.712	116.958	090617M2
<b>T14303</b>	South of Lol Gray	9/8/2016	M	-32.820	117.001	090816M1
<b>T14304</b>	NA	9/8/2018	M	NA	NA	090818M1
<b>T14305</b>	NA	9/8/2018	M	NA	NA	090818M2
<b>T14306</b>	Dryandra	9/12/2017	M	-32.790	116.966	091217M1
<b>T14307</b>	Dryandra	10/12/2017	M	-32.804	116.971	101217M1
<b>T14308</b>	North of Contine	11/4/2016	M	-32.881	117.013	110416M1
<b>T14310</b>	East of Yornaning	11/6/2020	M	-32.743	117.121	110620M1
<b>T14311</b>	Forestry Road	11/6/2021	NA	-32.721	117.041	110621U1
<b>T14312</b>	Forestry Road	11/6/2021	NA	-32.721	117.041	110621U2
<b>T14313</b>	South of Lol Gray	11/7/2016	F	-32.819	117.001	110716F1
<b>T14314</b>	North of Contine	11/10/2016	M	-32.837	116.938	111016M1
<b>T14315</b>	Cuballing townsite	11/10/2018	M	-32.821	117.172	111018M1
<b>T14317</b>	West Cuballing	14/5/2020	NA	-32.809	117.111	140520U1
<b>T14318</b>	Cuballing	14/8/2017	F	-32.810	117.186	140817F1

*Understanding sources of feral cats in Dryandra Woodland*

<b>Tissue ID</b>	<b>Location</b>	<b>Date</b>	<b>Sex</b>	<b>Latitude</b>	<b>Longitude</b>	<b>ID Tag</b>
<b>T14319</b>	Cuballing	14/8/2018	M	-32.818	117.183	140818M1
<b>T14320</b>	Cuballing	14/10/2018	NA	-32.808	117.200	141018U1
<b>T14321</b>	NA	14/11/2017	F	NA	NA	141117F1
<b>T14322</b>	West Cuballing	14/11/2017	M	-32.809	117.111	141117M1
<b>T14323</b>	Cuballing	14/12/2015	F	-32.816	117.178	141215F1
<b>T14324</b>	Contine	15/3/2019	F	-32.837	116.935	150319F1
<b>T14325</b>	South of Yornaning Road	15/4/2017	F	-32.753	117.036	150417F1
<b>T14326</b>	North of Montague	15/4/2017	M	-32.715	117.097	150417M1
<b>T14328</b>	Forestry Road	15/6/2017	F	-32.705	117.063	150617F2
<b>T14329</b>	North of Montague	15/6/2017	F	-32.697	117.076	150617F3
<b>T14330</b>	North of Montague	15/6/2017	M	-32.697	117.076	150617M2
<b>T14331</b>	North of Montague	15/6/2017	M	-32.697	117.076	150617M3
<b>T14332</b>	North of Contine	15/10/2016	M	-32.840	116.957	151016M1
<b>T14335</b>	Cuballing townsite	16/10/2018	NA	-32.821	117.172	161018U1
<b>T14336</b>	North of Yornaning Road	17/2/2017	M	-32.742	117.015	170217M1
<b>T14338</b>	Narrogin	17/5/2021	M	-32.929	117.155	170521M1
<b>T14339</b>	Narrogin	17/6/2017	M	-32.929	117.155	170617M1
<b>T14340</b>	West Yornanning	18/6/2017	F	-32.753	117.028	180617F1
<b>T14341</b>	West Yornanning	18/6/2017	F	-32.753	117.028	180617F2
<b>T14342</b>	West Yornanning	18/6/2017	F	-32.753	117.028	180617F3
<b>T14343</b>	West Yornanning	18/6/2017	F	-32.753	117.028	180617F4
<b>T14344</b>	North of Contine	18/10/2016	M	-32.853	116.957	181016M1
<b>T14345</b>	South of Lol Gray	18/11/2016	M	-32.848	117.023	181116M1
<b>T14346</b>	North of Yornaning Road	19/2/2017	M	-32.742	117.015	190217M1
<b>T14347</b>	Contine	19/3/2019	M	-32.899	116.996	190319M1
<b>T14348</b>	North of Contine	19/5/2016	NA	-32.845	116.926	190516U1
<b>T14349</b>	West Yornanning	19/7/2017	F	-32.753	117.028	190717F1
<b>T14350</b>	North of Yornaning Road	19/7/2017	F	-32.720	117.023	190717F2
<b>T14351</b>	West Yornanning	19/7/2017	F	-32.753	117.028	190717F3
<b>T14353</b>	North of Yornaning Road	19/7/2017	M	-32.720	117.023	190717M2

<b>Tissue ID</b>	<b>Location</b>	<b>Date</b>	<b>Sex</b>	<b>Latitude</b>	<b>Longitude</b>	<b>ID Tag</b>
<b>T14354</b>	North of Montague	19/7/2017	M	-32.697	117.076	190717M3
<b>T14355</b>	East of forest blocks	19/8/2016	M	-32.819	116.804	190816M1
<b>T14356</b>	North of Contine	19/10/2016	M	-32.848	116.990	191016M1
<b>T14357</b>	Cuballing townsite	19/10/2018	NA	-32.821	117.172	191018U1
<b>T14358</b>	NA	19/10/2020	F	NA	NA	191020F1
<b>T14359</b>	NA	19/10/2020	M	NA	NA	191020M1
<b>T14360</b>	Narrogin	19/12/2015	M	-32.835	116.963	191215M1
<b>T14363</b>	West Yornanning	20/5/2017	F	-32.753	117.028	200517F1
<b>T14364</b>	Narrogin	20/7/2016	M	-32.929	117.154	200716M1
<b>T14365</b>	Cuballing	20/8/2018	NA	-32.816	117.178	200818U1
<b>T14366</b>	West Cuballing	20/8/2021	M	-32.809	117.111	200821M1
<b>T14368</b>	NA	20/10/2017	M	NA	NA	201017M1
<b>T14369</b>	North of Contine	20/11/2016	F	-32.830	117.007	201116F1
<b>T14371</b>	West Cuballing	21/6/2018	NA	-32.809	117.111	210618U1
<b>T14372</b>	East of Cuballing	21/12/2016	F	-32.809	117.111	211216F1
<b>T14373</b>	North of Yornaning Road	22/2/2017	M	-32.727	117.009	220217M1
<b>T14374</b>	South of Yornaning Road	22/4/2017	F	-32.752	117.027	220417F1
<b>T14375</b>	South of Yornaning Road	22/4/2017	F	-32.753	117.028	220417F2
<b>T14376</b>	South of Yornaning Road	22/4/2017	M	-32.753	117.028	220417M1
<b>T14377</b>	South of Contine	22/9/2019	NA	-32.916	117.006	220919U1
<b>T14378</b>	Yornaning	22/11/2017	M	-32.742	117.165	221117M1
<b>T14379</b>	North of Yornaning Road	23/2/2017	M	-32.742	117.015	230217M1
<b>T14380</b>	South of Yornaning Road	23/4/2016	M	-32.752	117.032	230416M1
<b>T14381</b>	York Williams Road	23/4/2019	NA	-32.733	116.889	230419F1
<b>T14382</b>	West Yornanning	23/5/2017	F	-32.753	117.028	230517F1
<b>T14383</b>	West Yornanning	23/5/2017	M	-32.753	117.028	230517M1
<b>T14384</b>	West Yornanning	23/5/2017	M	-32.753	117.028	230517M2
<b>T14385</b>	West Cuballing	23/5/2021	F	-32.809	117.111	230521F1
<b>T14386</b>	West Yornanning	23/6/2021	NA	-32.753	117.028	230621U1
<b>T14387</b>	North of Montague	23/8/2017	F	-32.697	117.076	230817F1

*Understanding sources of feral cats in Dryandra Woodland*

<b>Tissue ID</b>	<b>Location</b>	<b>Date</b>	<b>Sex</b>	<b>Latitude</b>	<b>Longitude</b>	<b>ID Tag</b>
<b>T14388</b>	West Yornanning	23/8/2017	F	-32.753	117.028	230817F2
<b>T14389</b>	North of Montague	23/8/2017	F	-32.697	117.076	230817F3
<b>T14390</b>	North of Montague	23/8/2017	F	-32.697	117.076	230817F4
<b>T14391</b>	West Yornanning	23/8/2017	M	-32.753	117.028	230817M1
<b>T14392</b>	West Yornanning	23/8/2017	M	-32.753	117.028	230817M2
<b>T14393</b>	West of Contine	23/8/2017	NA	-32.859	116.860	230817U3
<b>T14394</b>	Narrogin	23/11/2021	M	-32.929	117.155	231121M1
<b>T14395</b>	Southwest of Cuballing	25/4/2017	F	-32.845	117.138	250417F1
<b>T14396</b>	North of Yornaning Road	25/4/2017	F	-32.846	117.137	250417F2
<b>T14397</b>	Southwest of Cuballing	25/4/2017	F	-32.845	117.137	250417F3
<b>T14398</b>	Southwest of Cuballing	25/4/2017	F	-32.846	117.138	250417F4
<b>T14399</b>	Southwest of Cuballing	25/4/2017	F	-32.846	117.137	250417F5
<b>T14400</b>	Southwest of Cuballing	25/4/2017	F	-32.846	117.137	250417F6
<b>T14401</b>	Southwest of Cuballing	25/4/2017	F	-32.847	117.138	250417F7
<b>T14402</b>	Southwest of Cuballing	25/4/2017	F	-32.846	117.138	250417F8
<b>T14403</b>	West Yornanning	25/4/2017	F	-32.754	117.028	250417F9
<b>T14404</b>	Southwest of Cuballing	25/4/2017	M	-32.846	117.138	250417M1
<b>T14405</b>	North of Yornaning Road	25/4/2017	M	-32.846	117.138	250417M2
<b>T14406</b>	Southwest of Cuballing	25/4/2017	M	-32.846	117.138	250417M3
<b>T14407</b>	Southwest of Cuballing	25/4/2017	M	-32.845	117.138	250417M4
<b>T14409</b>	South of Yornaning Road	25/4/2017	M	-32.753	117.028	250417M6
<b>T14410</b>	Contine	25/5/2017	F	-32.875	116.923	250517F1
<b>T14411</b>	West Yornanning	25/5/2017	M	-32.753	117.028	250517M3
<b>T14414</b>	Contine	26/5/2017	M	-32.859	116.939	260517M1
<b>T14415</b>	West Cuballing	27/3/2018	M	-32.809	117.111	270318M1
<b>T14416</b>	Contine	27/5/2017	M	-32.866	116.942	270517M1
<b>T14417</b>	Dryandra	28/2/2016	M	-32.845	117.137	280216M1
<b>T14419</b>	Southwest of Cuballing	28/4/2017	M	-32.846	117.139	280417M1
<b>T14422</b>	Forestry Road	28/6/2017	M	-32.719	117.041	280617M1
<b>T14423</b>	West Yornanning	28/9/2020	NA	-32.753	117.028	280920U1

<b>Tissue ID</b>	<b>Location</b>	<b>Date</b>	<b>Sex</b>	<b>Latitude</b>	<b>Longitude</b>	<b>ID Tag</b>
<b>T14424</b>	North of Yornaning Road	29/1/2017	F	-32.737	117.057	290117F1
<b>T14425</b>	East of forest blocks	29/6/2016	M	-32.827	116.777	290616M1
<b>T14426</b>	Southwest of Cuballing	29/6/2017	F	-32.856	117.127	290617F1
<b>T14427</b>	West Cuballing	29/10/2018	F	-32.809	117.111	291018F1
<b>T14428</b>	Narrogen	30/6/2016	F	-32.929	117.154	300616F1
<b>T14429</b>	West Yornanning	30/7/2017	M	-32.753	117.028	300717M1
<b>T14431</b>	South of Contine	30/11/2016	F	-32.916	117.006	301116F1
<b>T14432</b>	South of Contine	30/11/2016	M	-32.916	117.005	301116M1
<b>T14433</b>	South of Turners Road	31/7/2018	NA	-32.825	117.032	310718U1
<b>T14434</b>	Narrogen	31/7/2018	NA	-32.903	117.182	310718U2
<b>T14435</b>	NA	15/6/2017	M	NA	NA	150617M1
<b>T14436</b>	NA	9/6/2017	M	NA	NA	090617M1
<b>T14437</b>	NA	NA	M	NA	NA	NA

## Appendix 2 SNP Filtering

**SNP filtering and data cleaning protocol, including filtering steps and the thresholds chosen.**

The total dataset underwent filters 1 to 9. Additional filters were then performed where relevant, depending on the assumptions of the specific analysis.

Filter	Description	Thresholds (no. loci remaining)
<b>1. Call rate (individual)</b>	The threshold for excluding low quality individuals was chosen by balancing the trade-off between number of samples, importance of lower quality samples (i.e., in unique versus well sampled locations) and number of loci.	$\geq 0.55$ (16 individuals removed, 12,098 loci remaining)
<b>2. Call rate (locus)</b>	To determine the call rate threshold across loci, the relationship between call rate and population genetic summary statistics (heterozygosity estimates and F-statistics) was explored, under the assumption that data quality and biologically meaningful population genetic metrics should not be correlated. The threshold that minimised this correlation, while retaining a large enough dataset for adequate statistical power was chosen.	$\geq 0.9$ (6,123)
<b>3. Read count</b>	Upper and lower thresholds were chosen for average read count, as lower read counts under call heterozygotes (sample bias), while very high read counts may represent paralogous regions. To determine the average read count upper and lower thresholds, the relationship between read count and population genetic summary statistics was explored (as above). Thresholds that minimised this correlation, while retaining a large enough dataset for adequate statistical power was chosen.	$\geq 20$ and $\leq 150$ (4,329)
<b>4. Repeatability average</b>	The repeatability filter threshold was chosen to exclude poor quality loci, while not filtering out SNPs with high levels of heterozygosity (as heterozygote genotype calls may vary between replicates due to slight variations in reference/SNP reads).	$\geq 0.95$ (4,329)
<b>5. Minor allele frequency</b>	A MAF threshold that balanced the trade-off between including SNPs that represented sequencing error, versus removing true low frequency/private alleles was chosen. A minimum threshold that equated to each allele appearing at least twice in the dataset was used.	$\geq 0.025$ (3,874)
<b>6. Secondaries</b>	Non-independent SNPs were removed by randomly retaining one locus per sequence (and excluding other loci found in the same fragment).	1 SNP/ sequence (3,691)

Filter	Description	Thresholds (no. loci remaining)
<b>7. Sex-linked</b>	Loci that BLASTed to the X or Y chromosome in the <i>Felis catus</i> genome were removed.	Removed: "NC_058386.1_chromosome_X" "NW_025408526.1_chromosome_X_chrX_random_Un_scaffold_68" (3,593)
<b>8. Linkage disequilibrium</b>	The LD filter was performed last, as many non-independent SNPs were removed by the earlier filters. This means that performing the LD filter last maximises the number of quality loci that are retained. <i>SNPRelate</i> version 1.20.1 (Zheng et al., 2012) was used to perform pairwise genotypic correlations within a sliding window of 500,000 base pairs, removing SNPs with a correlation of $\geq 0.5$ .	$\geq 0.5$ (3,241)
<b>9. Missing data</b>	Individuals with missing coordinate information were removed, as all analyses required spatial information.	Missing coordinate information (6 individuals removed, 3,241 loci remaining)
<b>10. Relatedness</b>	Wang's pairwise relatedness (Wang, 2002) was calculated, using a relatedness threshold that likely represented half-siblings and above. One individual from each highly related pair was removed, to avoid biasing genetic analyses.	$\geq 0.2$ (63 individuals removed, 3,241 loci remaining)
<b>11. Spatial trimming</b>	To avoid biasing genetic clustering analyses by including uneven sampling of locations, samples were thinned to only include one randomly chosen individual within a 500 m radius. Note that spatial thinning was used for PCoA and genetic clustering analyses only (and these results were compared to the full dataset).	1 individual/500 m radius (40 individuals removed, 3,241 loci remaining)
<b>12. Hardy Weinberg Equilibrium</b>	For analyses that assume HWE, the R package <i>dartr</i> (Gruber et al., 2018) was used to detect loci that significantly departed from HWE assumptions, which were then removed from the dataset.	Significantly out of HWE (3,105)

## Appendix 3 Pairwise relatedness

Pairwise relatedness for 'highly related pairs' (where relatedness  $\geq 0.2$ ).

Individual 1	Individual 2	Wang's relatedness
T14326	T14424	0.7143
T14325	T14380	0.6976
T14340	T14342	0.6468
T14328	T14422	0.6466
T14377	T14378	0.6461
T14341	T14389	0.6457
T14340	T14279	0.6312
T14300	T14389	0.6271
T14300	T14341	0.6225
T14330	T14280	0.6149
T14326	T14436	0.609
T14328	T14331	0.6049
T14301	T14389	0.602
T14329	T14436	0.5867
T14300	T14301	0.5822
T14302	T14387	0.5815
T14263	T14411	0.5775
T14331	T14422	0.5766
T14374	T14375	0.5753
T14395	T14398	0.5734
T14301	T14436	0.5722
T14280	T14390	0.5674
T14389	T14436	0.5665
T14424	T14436	0.5621
T14326	T14331	0.5551
T14301	T14424	0.5526
T14411	T14383	0.5523
T14402	T14396	0.5505
T14325	T14382	0.5478
T14326	T14301	0.547
T14341	T14436	0.5437
T14326	T14422	0.5357
T14396	T14399	0.5319
T14301	T14341	0.5289
T14382	T14383	0.5285
T14396	T14406	0.5282

<b>Individual 1</b>	<b>Individual 2</b>	<b>Wang's relatedness</b>
<b>T14424</b>	T14389	0.5274
<b>T14376</b>	T14409	0.5248
<b>T14341</b>	T14424	0.5223
<b>T14326</b>	T14328	0.5214
<b>T14329</b>	T14389	0.514
<b>T14326</b>	T14389	0.5102
<b>T14374</b>	T14373	0.5099
<b>T14300</b>	T14424	0.5075
<b>T14432</b>	T14431	0.5052
<b>T14329</b>	T14341	0.5049
<b>T14343</b>	T14300	0.5008
<b>T14279</b>	T14436	0.5003
<b>T14279</b>	T14342	0.4986
<b>T14301</b>	T14329	0.4972
<b>T14340</b>	T14330	0.4962
<b>T14400</b>	T14396	0.4926
<b>T14397</b>	T14398	0.4921
<b>T14363</b>	T14383	0.491
<b>T14329</b>	T14424	0.4888
<b>T14300</b>	T14278	0.4873
<b>T14374</b>	T14383	0.4801
<b>T14273</b>	T14336	0.4794
<b>T14313</b>	T14425	0.4728
<b>T14376</b>	T14336	0.4681
<b>T14300</b>	T14329	0.4672
<b>T14402</b>	T14399	0.4661
<b>T14326</b>	T14341	0.4632
<b>T14409</b>	T14336	0.4578
<b>T14268</b>	T14336	0.454
<b>T14326</b>	T14329	0.4517
<b>T14343</b>	T14424	0.4501
<b>T14375</b>	T14377	0.4496
<b>T14429</b>	T14336	0.4496
<b>T14374</b>	T14382	0.4437
<b>T14346</b>	T14379	0.4403
<b>T14343</b>	T14301	0.4395
<b>T14343</b>	T14387	0.438
<b>T14349</b>	T14273	0.4372
<b>T14289</b>	T14378	0.4369
<b>T14424</b>	T14422	0.4355
<b>T14343</b>	T14341	0.4346

Individual 1	Individual 2	Wang's relatedness
T14325	T14263	0.43
T14331	T14424	0.4226
T14419	T14396	0.4198
T14343	T14329	0.4163
T14400	T14406	0.4152
T14374	T14409	0.4116
T14268	T14376	0.4115
T14343	T14342	0.4089
T14332	T14417	0.4059
T14374	T14376	0.4057
T14278	T14387	0.4055
T14330	T14281	0.4052
T14409	T14429	0.4023
T14343	T14389	0.3965
T14263	T14391	0.396
T14326	T14279	0.3953
T14329	T14387	0.3949
T14374	T14411	0.3944
T14376	T14429	0.3938
T14401	T14398	0.3937
T14387	T14424	0.3907
T14302	T14342	0.3902
T14279	T14330	0.3899
T14328	T14436	0.3894
T14397	T14407	0.3888
T14302	T14340	0.3883
T14300	T14326	0.385
T14330	T14390	0.3848
T14279	T14424	0.3831
T14301	T14279	0.3809
T14401	T14395	0.3806
T14269	T14427	0.3791
T14406	T14399	0.3784
T14258	T14259	0.3782
T14278	T14341	0.373
T14302	T14281	0.3659
T14301	T14278	0.3648
T14328	T14279	0.3637
T14373	T14383	0.3637
T14328	T14424	0.3619
T14325	T14374	0.3608

<b>Individual 1</b>	<b>Individual 2</b>	<b>Wang's relatedness</b>
<b>T14278</b>	T14389	0.36
<b>T14343</b>	T14278	0.3595
<b>T14268</b>	T14374	0.3584
<b>T14325</b>	T14391	0.3578
<b>T14268</b>	T14409	0.3569
<b>T14310</b>	T14330	0.3553
<b>T14374</b>	T14429	0.3546
<b>T14328</b>	T14329	0.3545
<b>T14258</b>	T14260	0.3539
<b>T14380</b>	T14382	0.3531
<b>T14382</b>	T14373	0.3501
<b>T14310</b>	T14302	0.3488
<b>T14269</b>	T14270	0.3484
<b>T14259</b>	T14260	0.3482
<b>T14340</b>	T14281	0.3475
<b>T14374</b>	T14336	0.3469
<b>T14375</b>	T14409	0.3454
<b>T14411</b>	T14382	0.3441
<b>T14342</b>	T14424	0.3438
<b>T14402</b>	T14406	0.3432
<b>T14340</b>	T14280	0.3426
<b>T14342</b>	T14387	0.3423
<b>T14368</b>	T14388	0.339
<b>T14329</b>	T14304	0.3384
<b>T14270</b>	T14427	0.337
<b>T14278</b>	T14424	0.3361
<b>T14279</b>	T14422	0.3357
<b>T14268</b>	T14429	0.3344
<b>T14400</b>	T14402	0.3317
<b>T14375</b>	T14336	0.3296
<b>T14343</b>	T14281	0.3295
<b>T14328</b>	T14390	0.329
<b>T14269</b>	T14415	0.3255
<b>T14422</b>	T14436	0.3246
<b>T14409</b>	T14383	0.323
<b>T14263</b>	T14380	0.3214
<b>T14279</b>	T14389	0.3211
<b>T14302</b>	T14330	0.3171
<b>T14310</b>	T14281	0.3165
<b>T14329</b>	T14281	0.3161
<b>T14343</b>	T14326	0.3158

Individual 1	Individual 2	Wang's relatedness
T14301	T14422	0.3152
T14340	T14422	0.3147
T14329	T14279	0.3143
T14330	T14342	0.3139
T14330	T14422	0.313
T14281	T14424	0.3123
T14343	T14436	0.3109
T14422	T14390	0.3108
T14342	T14422	0.3087
T14289	T14377	0.3085
T14389	T14422	0.3071
T14340	T14390	0.3068
T14301	T14342	0.3064
T14331	T14436	0.3062
T14342	T14436	0.3061
T14376	T14375	0.3059
T14263	T14382	0.3054
T14384	T14403	0.3054
T14305	T14403	0.305
T14281	T14387	0.3041
T14363	T14382	0.304
T14311	T14312	0.3036
T14273	T14409	0.3036
T14297	T14383	0.3035
T14395	T14407	0.3029
T14263	T14374	0.3021
T14411	T14391	0.3008
T14395	T14397	0.3007
T14310	T14340	0.3
T14329	T14422	0.2983
T14328	T14281	0.298
T14301	T14328	0.2979
T14297	T14374	0.2979
T14328	T14340	0.2941
T14328	T14389	0.2938
T14342	T14281	0.2934
T14314	T14336	0.2926
T14328	T14341	0.2923
T14325	T14373	0.2919
T14300	T14304	0.2913
T14349	T14403	0.2905

<b>Individual 1</b>	<b>Individual 2</b>	<b>Wang's relatedness</b>
<b>T14326</b>	T14387	0.2887
<b>T14302</b>	T14279	0.2881
<b>T14404</b>	T14405	0.2877
<b>T14300</b>	T14342	0.286
<b>T14328</b>	T14342	0.2849
<b>T14325</b>	T14411	0.2836
<b>T14300</b>	T14387	0.2824
<b>T14341</b>	T14387	0.282
<b>T14331</b>	T14389	0.2814
<b>T14279</b>	T14281	0.2795
<b>T14302</b>	T14390	0.2788
<b>T14329</b>	T14331	0.2788
<b>T14343</b>	T14302	0.2769
<b>T14300</b>	T14281	0.2766
<b>T14278</b>	T14329	0.2766
<b>T14387</b>	T14389	0.2753
<b>T14374</b>	T14363	0.2751
<b>T14342</b>	T14389	0.2743
<b>T14301</b>	T14331	0.2741
<b>T14301</b>	T14281	0.2729
<b>T14331</b>	T14281	0.2719
<b>T14353</b>	T14321	0.2712
<b>T14341</b>	T14281	0.2704
<b>T14328</b>	T14330	0.27
<b>T14429</b>	T14383	0.2698
<b>T14387</b>	T14436	0.2691
<b>T14375</b>	T14429	0.2687
<b>T14326</b>	T14342	0.268
<b>T14411</b>	T14373	0.2673
<b>T14310</b>	T14390	0.2665
<b>T14273</b>	T14374	0.2663
<b>T14268</b>	T14375	0.2661
<b>T14279</b>	T14280	0.266
<b>T14326</b>	T14278	0.2659
<b>T14343</b>	T14304	0.2655
<b>T14331</b>	T14342	0.2651
<b>T14341</b>	T14331	0.2642
<b>T14279</b>	T14331	0.2628
<b>T14300</b>	T14422	0.2627
<b>T14280</b>	T14281	0.2617
<b>T14279</b>	T14341	0.2614


Individual 1	Individual 2	Wang's relatedness
T14300	T14436	0.2608
T14301	T14387	0.2604
T14297	T14409	0.26
T14266	T14320	0.2591
T14279	T14387	0.2591
T14329	T14342	0.2589
T14302	T14424	0.2577
T14380	T14391	0.2562
T14341	T14342	0.2554
T14400	T14399	0.2554
T14341	T14422	0.2553
T14374	T14380	0.2537
T14281	T14422	0.2528
T14278	T14342	0.2519
T14302	T14422	0.2498
T14302	T14278	0.2467
T14375	T14373	0.2467
T14300	T14331	0.2465
T14310	T14422	0.2446
T14273	T14429	0.2445
T14273	T14376	0.2444
T14262	T14371	0.2437
T14310	T14342	0.2421
T14331	T14390	0.242
T14398	T14407	0.2403
T14340	T14424	0.24
T14314	T14417	0.2392
T14281	T14390	0.2392
T14406	T14407	0.2386
T14310	T14424	0.2384
T14297	T14376	0.2363
T14321	T14350	0.2355
T14297	T14411	0.2349
T14304	T14387	0.2346
T14278	T14436	0.2334
T14349	T14384	0.2334
T14374	T14391	0.2333
T14264	T14265	0.2322
T14281	T14389	0.2307
T14280	T14342	0.23
T14303	T14373	0.229

<b>Individual 1</b>	<b>Individual 2</b>	<b>Wang's relatedness</b>
<b>T14326</b>	T14281	0.228
<b>T14332</b>	T14336	0.2273
<b>T14297</b>	T14382	0.2263
<b>T14343</b>	T14279	0.2261
<b>T14321</b>	T14373	0.2256
<b>T14330</b>	T14424	0.2249
<b>T14340</b>	T14387	0.2246
<b>T14409</b>	T14382	0.2242
<b>T14300</b>	T14328	0.2227
<b>T14384</b>	T14363	0.2184
<b>T14304</b>	T14389	0.2174
<b>T14280</b>	T14422	0.2166
<b>T14340</b>	T14331	0.2162
<b>T14273</b>	T14375	0.2154
<b>T14304</b>	T14424	0.2142
<b>T14328</b>	T14280	0.2133
<b>T14305</b>	T14363	0.2119
<b>T14296</b>	T14336	0.2117
<b>T14409</b>	T14411	0.2117
<b>T14343</b>	T14422	0.21
<b>T14323</b>	T14398	0.21
<b>T14302</b>	T14329	0.2094
<b>T14375</b>	T14383	0.2091
<b>T14363</b>	T14373	0.2087
<b>T14304</b>	T14281	0.2084
<b>T14302</b>	T14328	0.208
<b>T14396</b>	T14407	0.2078
<b>T14310</b>	T14328	0.2065
<b>T14296</b>	T14409	0.2061
<b>T14281</b>	T14436	0.205
<b>T14303</b>	T14383	0.2046
<b>T14263</b>	T14373	0.2044
<b>T14417</b>	T14336	0.2039
<b>T14403</b>	T14391	0.2015

## Appendix 4 Genetic diversity estimates 2016 - 2021

### Estimates of genetic diversity parameters over sampling years.

Allelic richness ( $A_r$ ), expected heterozygosity ( $H_e$ ), observed heterozygosity ( $H_o$ ) and effective population size ( $N_e$ ). Note that years with sample sizes  $n < 10$  have low confidence. Effective population size could not be calculated for 2019 and 2020.





## Glossary

Single Nucleotide Polymorphism	a single nucleotide polymorphism (abbreviated SNP, pronounced 'snip') is a genomic variant at a single nucleotide base position in the DNA where one nucleotide (adenine, thymine, cytosine, or guanine) is replaced with another (e.g. adenine is replaced with cytosine). SNPs are the most common type of genetic variation present between individuals and are ubiquitous through the genome.
alleles	an allele is one of two or more versions of a DNA sequence (a single base or a segment of bases) at a given genomic location.
Heterozygosity (He)	the presence of two different alleles at a particular gene location, as opposed to homozygosity (same alleles at a particular gene location).
inbreeding coefficient ( $F_{IS}$ )	the probability that two alleles at any locus in an individual are identical by descent from the common ancestor of the two parents. A measure of population-level inbreeding.
effective population size ( $N_e$ )	a theoretical measure of the number of breeding individuals in a randomly mating population that would give rise to the observed genetic diversity. Effective population size is often contrasted with the census population size ( $N_c$ ), with $N_e$ typically being less than $N_c$ .



## References

- Adamack, A. T., & Gruber, B. (2014). POP GEN REPORT: Simplifying basic population genetic analyses in R. *Methods in Ecology and Evolution*, 5(4), 384–387. <https://doi.org/10.1111/2041-210X.12158>
- Aparicio, J. M., Ortego, J., & Cordero, P. J. (2006). What should we weigh to estimate heterozygosity, alleles or loci? *Molecular Ecology*, 15(14), 4659–4665. <https://doi.org/10.1111/j.1365-294X.2006.03111.x>
- Banks, S. C., & Peakall, R. (2012). Genetic spatial autocorrelation can readily detect sex-biased dispersal. *Molecular Ecology*, 21(9), 2092–2105. <https://doi.org/10.1111/j.1365-294X.2012.05485.x>
- Burbidge, A. A., & McKenzie, N. L. (1989). Patterns in the modern decline of western Australia's vertebrate fauna: Causes and conservation implications. *Biological Conservation*, 50(1–4), 143–198. [https://doi.org/10.1016/0006-3207\(89\)90009-8](https://doi.org/10.1016/0006-3207(89)90009-8)
- Burgess, B. T., Irvine, R. L., & Russello, M. A. (2022). Population genomics of Sitka black-tailed deer supports invasive species management and ecological restoration on islands. *Communications Biology*, 5(1), 223. <https://doi.org/10.1038/s42003-022-03159-5>
- Caye, K., & Francois, O. (2016). *tess3r: Inference of Spatial Population Genetic Structure* (1.1.0). R package.
- Coulon, A. (2010). GENHET: An easy-to-use R function to estimate individual heterozygosity. *Molecular Ecology Resources*, 10(1), 167–169. <https://doi.org/10.1111/j.1755-0998.2009.02731.x>
- Cowen, S., Clausen, L., Algar, D., & Comer, S. (2019). Using genetics to evaluate the success of a feral cat (*Felis catus*) control program in north-western Australia. *Animals*, 9(12), Article 12. <https://doi.org/10.3390/ani9121050>
- Cruz, V. M. V., Kilian, A., & Dierig, D. A. (2013). Development of DArT marker platforms and genetic diversity assessment of the U.S. collection of the new oilseed crop *Lesquerella* and related species. *PLoS ONE*, 8(5), e64062. <https://doi.org/10.1371/journal.pone.0064062>
- DBCA (2017). *Department of Biodiversity and Conservation and Attractions, 2017, Dryandra Sanctuary Management Plan 2017-2027*. Department of Biodiversity and Conservation and Attractions, Perth.
- DEC (2011). *Dryandra Woodland Management Plan No. 70*. Department of Environment and Conservation and Conservation Commission of Western Australia, Perth, WA.
- Do, C., Waples, R. S., Peel, D., Macbeth, G. M., Tillett, B. J., & Ovenden, J. R. (2014). NeEstimator v2: Re-implementation of software for the estimation of contemporary effective population size ( $N_e$ ) from genetic data. *Molecular Ecology Resources*, 14(1), 209–214. <https://doi.org/10.1111/1755-0998.12157>
- Doherty, T. S., Dickman, C. R., Johnson, C. N., Legge, S. M., Ritchie, E. G., & Woinarski, J. C. Z. (2017). Impacts and management of feral cats *Felis catus*

- in Australia. *Mammal Review*, 47(2), 83–97.  
<https://doi.org/10.1111/mam.12080>
- Doherty, T. S., Glen, A. S., Nimmo, D. G., Ritchie, E. G., & Dickman, C. R. (2016). Invasive predators and global biodiversity loss. *Proceedings of the National Academy of Sciences*, 113(40), 11261–11265.  
<https://doi.org/10.1073/pnas.1602480113>
- Double, M. C., Peakall, R., Beck, N. R., & Cockburn, A. (2005). Dispersal, philopatry, and infidelity: Dissecting local genetic structure in superb fairy-wrens (*Malurus cyaneus*). *Evolution*, 59(3), 625–635.
- Fischer, M. C., Rellstab, C., Leuzinger, M., Roumet, M., Gugerli, F., Shimizu, K. K., Holderegger, R., & Widmer, A. (2017). Estimating genomic diversity and population differentiation – an empirical comparison of microsatellite and SNP variation in *Arabidopsis halleri*. *BMC Genomics*, 18(1), 69.  
<https://doi.org/10.1186/s12864-016-3459-7>
- Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2018). DART: An R package to facilitate analysis of SNP data generated from reduced representation genome sequencing. *Molecular Ecology Resources*, 18(3), 691–699.  
<https://doi.org/10.1111/1755-0998.12745>
- Hedrick, P. W. (2011). *Genetics of Populations*, 2nd ed. Jones and Barlett.
- Huisman, J. (2017). Pedigree reconstruction from SNP data: Parentage assignment, sibship clustering and beyond. *Molecular Ecology Resources*, 17(5), 1009–1024. <https://doi.org/10.1111/1755-0998.12665>
- Jombart, T. (2008). adegenet: A R package for the multivariate analysis of genetic markers. *Bioinformatics*, 24(11), 1403–1405.  
<https://doi.org/10.1093/bioinformatics/btn129>
- Kilian, A., Wenzl, P., Huttner, E., Carling, J., Xia, L., Blois, H., Caig, V., Heller-Uszynska, K., Jaccoud, D., Hopper, C., Aschenbrenner-Kilian, M., Evers, M., Peng, K., Cayla, C., Hok, P., & Uszynski, G. (2012). Diversity Arrays Technology: A generic genome profiling technology on open platforms. In F. Pompanon & A. Bonin (Eds.), *Data Production and Analysis in Population Genomics: Methods and Protocols* (pp. 67–89). Humana Press.  
[https://doi.org/10.1007/978-1-61779-870-2\\_5](https://doi.org/10.1007/978-1-61779-870-2_5)
- Koch, K., Algar, D., & Schwenk, K. (2014). Population structure and management of invasive cats on an Australian Island. *The Journal of Wildlife Management*, 78(6), 968–975. <https://doi.org/10.1002/jwmg.739>
- Koch, K., Algar, D., Searle, J. B., Pfenninger, M., & Schwenk, K. (2015). A voyage to Terra Australis: Human-mediated dispersal of cats. *BMC Evolutionary Biology*, 15(1), 262. <https://doi.org/10.1186/s12862-015-0542-7>
- Koch, K., Pink, C., Hamilton, N., & Algar, D. (2020). A population genetic study of feral cats on Christmas Island. *Australian Journal of Zoology*, 68(3), 120.  
<https://doi.org/10.1071/ZO20081>
- Le Roux, J., & Wieczorek, A. m. (2009). Molecular systematics and population genetics of biological invasions: Towards a better understanding of invasive species management. *Annals of Applied Biology*, 154(1), 1–17.  
<https://doi.org/10.1111/j.1744-7348.2008.00280.x>

- Legge, S., Murphy, B. P., McGregor, H., Woinarski, J. C. Z., Augusteyn, J., Ballard, G., Baseler, M., Buckmaster, T., Dickman, C. R., Doherty, T., Edwards, G., Eyre, T., Fancourt, B. A., Ferguson, D., Forsyth, D. M., Geary, W. L., Gentle, M., Gillespie, G., Greenwood, L., ... Zewe, F. (2017). Enumerating a continental-scale threat: How many feral cats are in Australia? *Biological Conservation*, 206, 293–303. <https://doi.org/10.1016/j.biocon.2016.11.032>
- Marlow, N. J., Thomas, N. D., Williams, A. A. E., Macmahon, B., Lawson, J., Hitchen, Y., Angus, J., & Berry, O. (2015). Cats (*Felis catus*) are more abundant and are the dominant predator of woylies (*Bettongia penicillata*) after sustained fox (*Vulpes vulpes*) control. *Australian Journal of Zoology*, 63(1), 18. <https://doi.org/10.1071/ZO14024>
- Melville, J., Haines, M. L., Boysen, K., Hodkinson, L., Kilian, A., Smith Date, K. L., Potvin, D. A., & Parris, K. M. (2017). Identifying hybridization and admixture using SNPs: Application of the DArTseq platform in phylogeographic research on vertebrates. *Royal Society Open Science*, 4, 161061. <https://doi.org/10.1098/rsos.161061>
- O'Leary, S. J., Puritz, J. B., Willis, S. C., Hollenbeck, C. M., & Portnoy, D. S. (2018). These aren't the loci you're looking for: Principles of effective SNP filtering for molecular ecologists. *Molecular Ecology*, 27(16), 3193–3206. <https://doi.org/10.1111/mec.14792>
- Pew, J., Wang, J., Muir, P., & Frasier, T. (2015). *Related: An R package for analyzing pairwise relatedness data based on codominant molecular markers*. R package.
- R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Shaw, R. E., Spencer, P. B., Gibson, L. A., Dunlop, J. A., Kinloch, J. E., Mokany, K., Byrne, M., Moritz, C., Davie, H., Travouillon, K. J., & Ottewell, K. M. (2022). Linking life history to landscape for threatened species conservation in a multiuse region. *Conservation Biology*, 00, 00–00. <https://doi.org/10.1111/cobi.13989>
- Smouse, P. E., & Peakall, R. (1999). Spatial autocorrelation analysis of individual multiallele and multilocus genetic structure. *Heredity*, 82(5), 561–573. <https://doi.org/10.1038/sj.hdy.6885180>
- Spencer, P. B. S., Yurchenko, A. A., David, V. A., Scott, R., Koepfli, K.-P., Driscoll, C., O'Brien, S. J., & Menotti-Raymond, M. (2016). The Population Origins and Expansion of Feral Cats in Australia. *Journal of Heredity*, 107(2), 104–114. <https://doi.org/10.1093/jhered/esv095>
- Sunnucks, P., & Hales, D. F. (1996). Numerous transposed sequences of mitochondrial cytochrome oxidase I-II in aphids of the genus Sitobion (Hemiptera: Aphididae). *Molecular Biology and Evolution*, 13(3), 510–524. <https://doi.org/10.1093/oxfordjournals.molbev.a025612>
- Telfer, E. J., Stovold, G. T., Li, Y., Silva-Junior, O. B., Grattapaglia, D. G., & Dungey, H. S. (2015). Parentage reconstruction in *Eucalyptus nitens* using SNPs and microsatellite markers: A comparative analysis of marker data power and robustness. *PLOS ONE*, 10(7), e0130601. <https://doi.org/10.1371/journal.pone.0130601>

- Wang, J. (2002). An estimator for pairwise relatedness using molecular markers. *Genetics*, 160(3), 1203–1215. <https://doi.org/10.1093/genetics/160.3.1203>
- Wang, J. (2018). Effects of sampling close relatives on some elementary population genetics analyses. *Molecular Ecology Resources*, 18(1), 41–54. <https://doi.org/10.1111/1755-0998.12708>
- Waples, R. S. (2006). A bias correction for estimates of effective population size based on linkage disequilibrium at unlinked gene loci. *Conservation Genetics*, 7(2), 167–184. <https://doi.org/10.1007/s10592-005-9100-y>
- Watson, K. M.-A., Mikac, K. M., & Schwab, S. G. (2021). Population genetics of the invasive red fox, *Vulpes vulpes*, in south-eastern Australia. *Genes*, 12(5). <https://doi.org/10.3390/genes12050786>
- Wright, S. (1969). *Evolution and the genetics of populations: Vol. 2. The theory of gene frequencies*.
- Zheng, X., Levine, D., Shen, J., Gogarten, S. M., Laurie, C., & Weir, B. S. (2012). A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics*, 28(24), 3326–3328. <https://doi.org/10.1093/bioinformatics/bts606>