



Restoring an appropriate fire regime to fragmented vegetation: a vital attributes approach

Carl Gosper, Colin Yates and Suzanne Prober
CSIRO Sustainable Ecosystems and
WA Department of Environment and Conservation
2 December 2009



Department of
Environment and
Conservation



Our environment, our future



South-West Western Australia: a global biodiversity hotspot

- High richness and endemism in vascular plants
- Highly threatened
- Mediterranean climate
- Fire-prone landscape
- Poor understanding of historic fire regimes
- Fire is a key process driving vegetation composition, health and recruitment
- Many plants possess traits that enable them to persist following fire



The WA wheatbelt

Highly fragmented

As little as 2-3% cover in some districts

Altered disturbance regimes

- Fire frequency, season, extent &/or intensity
- Different sources of ignition
- Loss of connectivity (for passage of fire)
- Fire exclusion and suppression
- Different parts of the landscape appear to be experiencing different fire intervals



CSIRO. Restoring an appropriate fire regime to fragmented vegetation



Restoring fire?

Conservation risks of infrequent fire:

- Vegetation community change
- Senescence
- Lack of recruitment opportunities

Uncertainty as to what constitutes an appropriate fire regime for biodiversity conservation

Study aim:

Can appropriate fire intervals for plant communities be reconstructed from sampling the vital attributes of fire-sensitive examples of key Plant Functional Types?



Department of
Environment and
Conservation

Our environment, our future



Species selection

Functional types vulnerable to fire interval

- Focus on processes and mechanisms that fire impacts species
- Means of persistence through fire (sprouter vs. obligate seeder)
- Means of seed survival (persistent soil-stored vs. canopy-stored)

Serotinous species

- Seed bank survival reliant on adult survival
- Seed bank exhausted in each disturbance
- Recruit predominately after disturbance



Serotinous species selection

Obligate seeders

Reliant of recruitment from seed for persistence after fire

Vulnerable to fire intervals:

- less than juvenile period
- greater than adult longevity (senescence)

Sprouters

Rely on adult survival, often having low rates of recruitment

Vulnerable to:

- Adult mortality (competitive interactions, disturbance, disease)

Paired contrasts

Family	Obligate seeders	Sprouters
Proteaceae	<i>Banksia violacea</i>	<i>B. pteridifolia</i>
Proteaceae	<i>Hakea pandanycarpa</i>	<i>H. incrassata</i>
Proteaceae	<i>Petrophile glauca</i>	<i>P. seminuda</i>
Myrtaceae	<i>Beaufortia micrantha</i>	<i>Leptospermum spinescens</i>



Department of
Environment and
Conservation

Our environment, our future



Methods

- Stratification by time since fire
- 9 vegetation ages
- Range: 2 to > 46 yrs post-fire
- Species sampled at 4 sites per fire age (more for > 46 yrs)
- 10 individuals per site

Field measurements:

- Mortality
- Fruit number



Department of
Environment and
Conservation

Our environment, our future



Study community

Tallerack mallee-heath

- Tallerack *Eucalyptus pleurocarpa*
- Scattered mallees over dense shrub layer
- Myrtaceae and Proteaceae dominant
- This and similar communities (kwongan) are widespread across Mediterranean-climate SW Western Australia



Acceptable fire intervals

We use two methods to estimate minimum and maximum fire intervals

Minimum fire intervals

- Time to maturation and commencement of accumulation of seed bank
- Primary (obligate seeders) or secondary (sprouters) juvenile period
- Our estimates derived from (i) the proportion of individuals with fruit and (ii) mean fruit crop size



Department of
Environment and
Conservation

Our environment, our future




CSIRO

Minimum fire intervals – method 1

Sprouter - *Banksia pteridifolia*

Proportion of individuals with fruit

- Time since fire when $\geq 50\%$ of individuals carry fruit
- Derived from regression of the proportion of sampled individuals carrying fruit per site with time since fire




Secondary
juvenile period
~ 8 yrs



Minimum fire intervals – method 1

Obligate seeder - *Petrophile glauca*



Primary juvenile period ~ 15 yrs




Minimum fire intervals – method 2

Sprouter - *Banksia pteridifolia*


Mean fruit number

- Time since fire when mean number of fruits per plant reaches 25% of maximum
- Derived from regression of count data with time since fire (sites combined)



Minimum fire intervals – method 2

Obligate seeder - *Petrophile glauca*



Primary juvenile period ~ 20 yrs



Acceptable fire intervals

Maximum fire interval

- Period when time since fire exceeds plant and/or seed bank longevity
- Appropriate metrics have not been well defined
- Our estimates derived from (i) the proportion of dead individuals and (ii) mean fruit crop size (measuring decline of the seed bank)




Maximum fire intervals – method 1

Sprouter - *Banksia pteridifolia*


Proportion of dead individuals

- Time since fire when $\geq 25\%$ of individuals are dead
- Derived from regression of the proportion of dead individuals per site with time since fire



Maximum fire intervals – method 1

Obligate seeder - *Petrophile glauca*



Max. interval > 46 yrs




Maximum fire intervals – method 2

Mean fruit number


Sprouter - *Banksia pteridifolia*

- Time since fire when mean number of fruits per plant declines to 25% of maximum
- Derived from regression of count data with time since fire (sites combined)



Maximum fire intervals – method 2

Obligate seeder - *Petrophile glauca*



Fruit crop declined in long unburnt vegetation, but did not reach 25% threshold



Minimum fire intervals – all species

Time since fire to:	≥ 50% of popn.	25% of max. crop
Obligate seeder (primary juv. period)		
<i>Banksia violacea</i>	Did not reach	19 yrs
<i>Hakea pandanycarpa</i>	36 yrs	36 yrs
<i>Petrophile glauca</i>	15 yrs	20 yrs
<i>Beaufortia micrantha</i>	8 yrs	14 yrs
Sprouter (secondary juv. period)		
<i>Banksia pteridifolia</i>	8 yrs	3 yrs
<i>H. incrassata</i>	No relationship	No relationship
<i>P. seminuda</i>	No relationship	No relationship
<i>Leptospermum spinescens</i>	7 yrs	5 yrs



Department of
Environment and
Conservation

Our environment, our future



Maximum fire intervals – all species

Time since fire to:	25% mortality	25% of max. crop
Obligate seeder		
<i>Banksia violacea</i>	> 46 yrs [#]	No decline
<i>Hakea pandanocarpa</i>	> 46 yrs	> 46 yrs
<i>Petrophile glauca</i>	> 46 yrs	> 46 yrs [#]
<i>Beaufortia micrantha</i>	> 46 yrs [#]	> 46 yrs [#]
Sprouter		
<i>Banksia pteridifolia</i>	> 46 yrs [#]	No decline
<i>H. incrassata</i>	No relationship	No relationship
<i>P. seminuda</i>	30-40 yrs [#]	No relationship
<i>Leptospermum spinescens</i>	20-25 yrs	19-39 yrs

[#]Significant relationship with time since fire (increase in mortality, or reduction in fruit crop, in vegetation of a certain age), but did not reach relevant threshold

Appropriate fire intervals

- Based on the species sampled, an appropriate minimum fire return interval ~36 yrs
- Among obligate seeders, several show senescence under very long fire-return intervals (>> 50 yrs)
- Among sprouters, higher mortality and lower fruit crop size of some species at intermediate ages suggests that repeated intermediate intervals would lead to population declines
- From a plant vital attribute perspective, an appropriate fire return interval range may be variable, but within the range 36 - >>50 yrs



Department of
Environment and
Conservation

Our environment, our future



Restoration of fire interval

How do these estimates compare to recent fire regimes?

- Recent average fire intervals vary with landscape context (Blair Parsons unpubl. data):
 - Small remnants: ~340 yrs
 - Large remnants: ~70 yrs
 - Continuous vegetation: ~40 yrs
- Active fire introduction may be appropriate in small remnants
- A increase in fire interval may be desirable in continuous vegetation
- Challenges in fire regime restoration in remnants:
 - Weed invasion
 - Operational constraints



Department of
Environment and
Conservation

Our environment, our future



Vital attributes approach

Vital attributes:

- useful for deriving estimates of minimal fire intervals
- less useful in deriving estimates of maximum fire intervals, due to the lack of long-term fire history data
- do not provide any information on other aspects of fire regime that would provide guidance in restoring fire regimes in remnants (season, patchiness, extent, intensity)



Department of
Environment and
Conservation

Our environment, our future



CSIRO

Carl Gosper, Suzanne Prober and Colin Yates

**Department of Environment and Conservation and
CSIRO Sustainable Ecosystems**

Phone: +61 8 9333 6442

Email: carl.gosper@csiro.au

Web: <http://www.csiro.au/science/WAFireFragment.html>

© CSIRO Australia, 2009

All photos: C. Gosper. Fire age polygons kindly provided by DEC Fire Management Services.

www.csiro.au

Thank you



Department of
Environment and
Conservation



Our environment, our future

Contact Us

Phone: 1300 363 400 or +61 3 9545 2176

Email: enquiries@csiro.au Web: www.csiro.au

