

LIBRARY

Department of Biodiversity,
Conservation and Attractions

This PDF has been created for digital preservation. It may be used for research but is not suitable for other purposes. It may be superseded by a more current version or just be out-of-date and have no relevance to current situations.




FIG 4. AFTER LOGGING

Note that:

1. Some potential crop trees surplus to stocking needs may remain after the commercial operation because they are below marketable size.
2. Occasional crop trees below marketable size may be retained in what is otherwise a gap. However, every opportunity should be taken to remove them commercially if possible.
3. Small groups of poles have little chance of surviving the felling of surrounding large trees.
4. Cull trees remain after commercial operations.
5. Some "non-crop trees" may be retained if their removal would cause excessive damage to crop trees.

LEGEND




FIG 5. FOLLOW-UP SILVICULTURAL TREATMENT

Following logging, surplus stems are removed to produce a stand containing patches of:

- Partial cutting (or shelterwood) to promote the development of seedlings into ground coppice (Objective 1).
- Complete removal of the overstorey to allow ground coppice to develop into poles (Objective 2).
- Thinning to concentrate the stand growth onto fewer stems (Objective 3).

Note that:

1. Some gaps which result are too small to be effective regeneration areas and larger than necessary for a thinning. This cannot always be avoided.
2. This gap has been extended by the removal (or poisoning) of cull trees. It has not been extended far beyond the gap initiated by the commercial operation.
3. Where insufficient crop trees exist for a complete thinning, individual crop trees are released from competition.

Coppicing may be carried out in regeneration areas but coppice is discouraged in thinning areas where it will produce undesirable competition for the crop trees.



Treemarking and Silviculture in the Jarrah Forest

A TRAINING BRIEF FOR OPERATIONAL STAFF
PREPARED BY
J. BRADSHAW

SILVICULTURE BRANCH

Department of Conservation & Land Management

1985



Department of Conservation and Land Management, W.A.

SILVICULTURAL OBJECTIVES

The jarrah forest is a complex mosaic of stand structure, age and quality. This is due to natural variation and a number of logging and silvicultural treatments carried out in the last 100 years. Although discrete structures may be recognised (Fig 2), changes occur continuously throughout the forest, often blending from one to another without distinct boundaries.

Given this complexity, what is the most appropriate silvicultural system to use? How should it be treemarked?

The objective of treemarking is to create the sort of stand structure most suited to the particular stage of development of that stand - whether it be for regrowth development or acceleration of tree growth. Because the forest is so variable, a fixed prescription will not work for every stand, but there are basic principles which apply to all stands.

If a tree is to be removed from a stand it should be done to achieve one of three silvicultural objectives:

1. **To reduce competition to allow seedlings to develop into ground coppice (large lignotubers)** (i.e. a partial removal of the overstorey to retain a forest cover while the regeneration develops.)
2. **To allow ground coppice to develop unimpeded into saplings, poles and trees.** (i.e. a complete removal of the overstorey.)
3. **To promote growth on the remaining trees** (i.e. a thinning. Regeneration is not wanted at this stage.)

Only one of these objectives should be sought in any one patch of forest at one time. The desirable size of this patch is 4 x mature tree height (or more) in diameter. If the patches are smaller than this, the felling of surrounding larger trees in the future will cause excessive damage to the regrowth in these patches. The existing stand structure will not always allow the 4 x tree height rule to be applied; nevertheless we should aim to push the stand in this direction whenever the opportunity arises.

When the emphasis is placed on selecting the trees to be retained as part of the next generation of forest a logical pattern begins to emerge. The trees available for cutting are those which are surplus to requirements as crop trees. When the commercial logging and the follow-up silvicultural treatment have been completed, the result should be a forest in a condition suitable for the optimum development of each particular stage of growth in each part of the stand.

THE QUALITIES OF A CROP TREE

A crop tree should be in the codominant level, have a potentially useful bole (at least 3m free of any defect that would preclude its use as a sawlog) and have a good crown. Crown size is important since future growth will depend upon it. There is no size limit to crop trees and there is no obligation to remove a tree simply because it has reached a marketable size.




FIG 1. CROWN SIZE AND INCREMENT ARE CLOSELY CORRELATED. (LONERAGAN)




FIG 2. STAND STRUCTURE & POTENTIAL CROP TREES

The stands depicted here show a variety of structural types:

- A - Even-aged (from clear felling)
- B - Grouped (from group selection cutting and follow up treatment of culls)
- C - Uniform (from single tree selection cutting or group selection cutting with no follow-up)
- D - Virgin (virgin forest may have a variety of structures but a preponderance of large sizes is common)

Regardless of the structural type, an initial identification of potential crop trees will show a pattern which will indicate whether a thinning or a regeneration cut is the most appropriate.

LEGEND





FIG 3. SELECTION OF CROP TREES

- In this patch there are few potential crop trees. Because there is inadequate ground coppice, a partial cut retaining some additional "non-crop trees" is carried out. The retained trees provide seed and a shelterwood while lignotubers develop (Objective 1).
- These patches contain few potential crop trees but have adequate stocking of ground coppice. All marketable stems (including potential crop trees) are removed to release the ground coppice, i.e. no stems are marked for retention (Objective 2).
- This patch contains sufficient potential crop trees to warrant thinning (Objective 3). Thinning need not be restricted to pole sizes and is a valid option for all vigorous trees. Potential crop trees, surplus to those required for a specified density, are not marked for retention and may be removed.