

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP ^{-Lower} Foothills SITE NUMBER SERP01


DATE TRIP 23/10/96 RECORDERS AM KKM DL

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Chaffield Rd
Mud Map Draw a sketch of the location of the site below



SERP01 - in flagged stand of Acacia horridula

Road Location

Geographic Location Latitude 32 22 39.2 S Longitude 115 59 55.3 E Altitude 70m
GPS ☒ Accuracy Jawahdale PDGP 2.7
Map Sheet

Photograph Name Number #35 - roll #3?

Topographic Position (Circle the correct response)
Upland ☐ Lower slope ☒
Upper slope ☐ Valley ☐
Midslope ☐ Washline ☐

2. SITE DATA Circle the correct response

Slope flat ☐ gentle ☒ steep ☐ Aspect N NE E SE S SW W ☒ NW

Surface Soil clayey, loamy grit Depth to Rock 0-1m Colour Grey to Dk Brown

Exposed Rock type granite % surface cover 2 size range 0.5 - 10cm

Litter 60 20 % surface cover Depth (cm) 2-5cm

Bare Ground | % surface cover)

Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

7
lower slope!
Ridge Hill
shelf
colluvium
shale
laterite
(from geology map)






BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Muir)- use pencil only

3. VEGETATION STRUCTURE AND COVER

NH Quartzite!

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class 2-10% 10-30% 30-70% over 70%

TREES				MALLEES	
	over 15m	5-15m	under 5m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		10-30%			
DOMINANT SPECIES		<i>E. lane-poolii</i>			
SHRUBS			SHRUBS		
	over 2m	2m - 1.5m	1.5 - 1m	1-0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)		2-10%		10-30%	30-70
DOMINANT SPECIES		<i>K. australis</i>		<i>X. acanthostachya</i> <i>H. trifurcata</i> <i>Dry. ovata</i>	<i>Melaleuca scabra</i> <i>Caloch. tomentos</i> <i>Hibb. hypericoides</i>
GRASSES		HERBS	SEDGES	under 0.5m	
	over 0.5m				
LIFE FORM					
COVER CLASS (%)	Not really a 2-10% layer	2-10%			
DOMINANT SPECIES	<i>N. alopec.</i>	Mixed			

4. VEGETATION CONDITION

Comments

Open *E. lane-poolii* woodland with
Emergent *Kingia australis* & *X. acanthostachya* a
low heath upstage. DRE *Acacia horridula* - lower
slope of scarp face. 5/5! Excellent condition
Rich sclerophyllous (proteaceous & myrtaceous) heath.

10

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIFIC PRESENCE Label each plant with plants number, site code, date and plant's name or working name if required

SITE No	SERPISI	Record on Sheet	• Column 1	plant name
---------	---------	-----------------	------------	------------

Date 23.10.96 • Column 2 plant number

13-12-96(R) • Column 3 flowering time- TICK if species flowering

• Column 4 Identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Euc. lane-poolei</i>	✓	✓		<i>Hakea lessocarpha</i>	✓	—	1/2	<i>Hydrocotyle callicarpa</i>	✓	✓	1/2
<i>Petrophile striata</i>	✓	✓	1/2	<i>Xanth. acanthostachya</i>	✓	—	1/2	<i>Drosera menziesii</i>	✓	✓	✓
<i>Melaleuca aff. scabra</i>	✓	✓	1/2	<i>Kingia australis</i>	✓	—	1/2	<i>Romulea rosea</i>	✓	✓	✓
<i>Acacia obovata</i>	✓	✓	1/2	<i>Hypocalymma angustifolium</i>	✓	—	1/2	<i>Dampiera alata</i>	✓	✓	1/2
<i>Hibbertia hypericoides</i>	✓	✓	1/2	<i>Isopogon asper</i>	✓	—	1/2	<i>Stylidium hispidum</i>	✓	✓	✓
<i>Gompholobium polymorphum</i>	✓	✓	1/2	<i>Gompholobium marginatum seedling</i>	✓	—	✓	<i>Styl. brunonianum</i>	✓	✓	✓
<i>Bossiqea eriocarpa</i>	✓	✓	1/2	<i>Dryandra lindleyi</i>	✓	—	1/2	<i>Goodenia caerulea</i>	✓	✓	R
<i>Gompholobium knightianum</i>	✓	✓	1/2	<i>Asteroloma pallidum</i>	✓	—	1/2	<i>Styl. pycnostachya (leaves)</i>	✓	✓	0
<i>Leucopogon pulchellus</i>	✓	✓	1/2	GRASSES				<i>Drosera odorifera</i>	✓	✓	✓
<i>Hibbertia sp. acerosa</i>	✓	✓	0	<i>Stipa pycnostachya</i>	✓	✓	✓	<i>Lomandra sericea</i>	✓	✓	1/2
<i>Darwinia thymoides</i>	✓	✓	1/2	<i>Briza minor</i>	✓	✓	✓	<i>Cassynia - porneiformis</i>	✓	✓	✓
<i>Xanthorrhoea gracilis</i>	✓	✓	0	<i>Amphipogon stricta</i>	✓	✓	✓	<i>Conostylis setosa</i>	✓	✓	✓
<i>Acacia horridula</i>	✓	✓	1/2	<i>Neurachne absepeuroidea</i>	✓	✓	1/2	<i>Paterosonia umbrosa v. xanthina</i>	✓	✓	1/2
<i>Acacia nervosa</i>	✓	✓	1/2					<i>Haemodorum laxum</i>	✓	✓	1/2
<i>Eriostemon spicatus</i>	✓	✓	1/2					<i>Cassynia glabella</i>	✓	✓	✓
<i>Hibbertia confertata</i>	✓	✓	1/2					<i>Drosera patulifolia</i>	✓	✓	✓
<i>Grevillea pilulifera</i>	✓	✓	1/2					<i>Hypochaeris glabra</i>	✓	✓	1/2
<i>Acacia feretifolia</i>	✓	✓	0	HERBS				<i>Tricoryne elatior</i>	✓	✓	✓
<i>Labeckia punctata</i>	✓	✓	0	<i>Conostylis setosa (?) sericea</i>	✓	✓	R	<i>Sonchus oleraceus</i>	✓	✓	1/2
<i>Lechenaulia biloba</i>	✓	✓	1/2					<i>Arthropodium capillipes</i>	✓	✓	✓
<i>Gompholobium sp.</i>	✓	✓	1/2	<i>Comesperma virgatum</i>	✓	✓	0	<i>Stylidium bulbiferum</i>	✓	✓	✓
<i>Acacia latericicola</i>	✓	✓	1/2	<i>Paterosonia occidentalis</i>	✓	✓	✓	<i>Thysanotus tenuifolius</i>	✓	✓	✓
<i>Dryandra bipinnatifida</i>	✓	✓	1/2	<i>Isotoma hypochaerifolius</i>	✓	✓	R	SEDGES			
<i>Pimelea suaveolens</i>	✓	✓	✓	<i>Oreocaularia apiculata</i>	✓	✓	1/2	<i>Tetraria capillaris</i>	✓	✓	1/2
<i>Baeckea camphorosma</i>	✓	✓	1/2	<i>Lomandra hemaphrodita</i>	✓	✓	1/2	<i>Schoenus grammophyllus</i>	✓	✓	1/2
				<i>Wahlenbergia preissii</i>	✓	✓	1/2	<i>Lepidosperma angustatum (leaf)</i>	✓	✓	✓
<i>Hakea trifurcata</i>	✓	✓	1/2	<i>Leuenhookia pusilla</i>	✓	✓	✓	<i>Schoenus sp. serapi</i>	✓	✓	✓
<i>Hakea stenocarpa</i>	✓	✓	1/2	<i>Thelymitra binita</i>	✓	✓	0	<i>Lorocarya fasciculata</i>	✓	✓	1/2
<i>Dryandra armata</i>	✓	✓	1/2	<i>Laxmannia squarrosa</i>	✓	✓	✓	<i>Tetraria octandra</i>	✓	✓	1/2
<i>Ocoteanthus torulosis</i>	✓	✓	1/2	<i>Xanthosia huegelii</i>	✓	✓	1/2	<i>Mesomelaena tetragona</i>	✓	✓	1/2
<i>Olearia paucidentata</i>	✓	✓	✓	<i>Trachymene pilosa</i>	✓	✓	✓	<i>Caladenia flava</i>	✓	✓	✓
<i>Hakea undulata</i>	✓	✓	1/2					<i>Stylidium carnosum</i>	✓	✓	✓

• Styl pycnostachy - enter ✓ OK

Ca_HCO3	CEC	exCa	exNa	exK	exMg	Alt	Lat
63	63	63	63	63	63	63	63

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP01

Record on Sheet

Date 23-10-96

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>B. lane-poolii</i>	✓		✓	<i>Dryandra nitida</i>	✓			<i>Harmiodorum</i> sp. (strapl?)	✓		✓
				<i>Dryandra armata</i>	✓		✓	<i>Cassytha racemosa</i>	✓		✓
				<i>Dryandra bipinnatifida</i>	✓		✓				
SHRUBS				<i>Xanthosia hueglinii</i>				<i>Hakea undulata</i>	✓		
<i>Hakea lissocarpa</i>	✓			<i>Astroloma pallidum</i>				<i>Leucopogon</i> sp.	✓		✓
<i>Xan. acanthostachya</i>	✓			<i>Thomasia</i> sp.				<i>Hydrocotyle</i>	✓		✓
<i>Kingia MALLEES australis</i>	✓			<i>Hakea stenocarpa</i>	✓			<i>Thysanotus</i> (Bival' rosette)			✓
<i>Calothamnus torulosus</i>	✓		✓					<i>Trachymene pilosa</i>			
<i>Metaleuca ? scrub</i>	✓		✓	GRASSES				<i>Sonchus oleraceus</i>	✓		
<i>Hypocalymma angustifolium</i>	✓		✓	<i>Neuradme alopecuroides</i>	✓			<i>Stipa</i> sp.			
SHRUBS				<i>Lomandra laevis</i>				<i>Briza minor</i>			
<i>Petrophile striata</i>			✓	<i>Hypericium glabra</i>				<i>Hypochaeris glabra</i>			
<i>Eriostemon spiratum</i>	✓		✓	<i>Trichosyne ? elatior</i> (PF)				SEDGES			
* <i>Hibbertia communata</i>	✓		✓	<i>Lomandra - single grey leaf ?</i> → as coll before	✓			<i>Loxocarya pasiculata</i>	✓		✓
<i>Leopogon asper</i> (dead)			✓	<i>Lomandra setacea</i> caespitosa				* <i>Tetraria axillaris</i>	✓		✓
<i>Acacia hauridula</i>	✓		✓	<i>Anthopogon ? Anthopodium</i>				Very fine sedge ?			
<i>Darwinia thymoides</i>	✓		✓	<i>Wahlenbergium ? Wahlenbergia</i>			✓	<i>Genus - black base</i>	✓		✓
<i>A. nuxia</i> <i>Acacia</i> sp. broad phyllode	✓		✓	HERBS				<i>Tetraria capillaris</i>	✓		✓
<i>A. nuxia</i> <i>Acacia</i> sp. - crinkled broad leaf	✓		✓	<i>Prosera strobilifera</i>			✓	<i>Lepidosperma terete</i> rough	✓		✓
<i>Hakea trifurcata</i>	✓		✓	<i>Stylidium pilulifera</i>	✓		✓	<i>Mesomelaena tragana</i>			✓
<i>Dampiera alba</i>	✓		✓	<i>Stylidium bruceianum</i>	✓		✓	<i>Lepidosperma angustifolium</i>	✓		✓
<i>Bossea eriocarpa</i>	✓		✓	<i>Paterosia ? occidentalis</i>	✓		✓				
<i>Acacia drummondii</i>			✓	<i>Stylidium bulbiferum</i>				<i>Caladenia flava</i>			
<i>Hakea camphorosa</i>	✓		✓	<i>Conostylis cerosa</i>	✓		✓	<i>Stylidium carnosum</i>	✓		✓
<i>Lechmanulka biloba</i> (nf) ??	✓		✓	<i>Prosera palda</i>			✓	Small annual xanthosia			✓
Small foliose seedling - ??				<i>Prosera menziesii</i>			✓	<i>Stylidium bulbiferum</i>			✓
<i>Compholobium knightianum</i>	✓		✓	<i>Paterosia ? (conostylis setosa)</i>	✓		✓	<i>Crotophobium trinerum</i> (- Polymorph -)			✓
* <i>Hibbertia hypericoides</i>	✓		✓	Small climbing <i>Prosera</i>			✓	<i>Olearia paucidentata</i>			✓
<i>Isopogon</i> seedling - ??	✓		✓	<i>Remulla rosea</i>			✓	<i>Hibb. communata</i> (seedling)			✓
<i>Hypericaria laspidula</i>				<i>Spatzmania squarosa</i>				<i>Levenhookia axillaris</i>			
<i>Cervilla hylephora</i> seedling	✓		✓	<i>Larrea</i>				<i>Pinelia suaveolens</i>			

✓ *Compholobium pilulifera marginata* (seedling) ✓• *Hydrocotyle* ✓

Operculina

Label each plant with plants number, site code, date and plant's name or working name if required

Record on Sheet

SERPØ1

Rescore ↗

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

[illegible]

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine N.P. SITE NUMBER SERP02


DATE TRIP 23.10.96 RECORDERS AM DL

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location 400D Survey track on westernmost ridge-line KEY access only

Geographic Location Latitude 32 22 27.1 S Longitude 116 00 08.8 E Altitude 180m
GPS Accuracy 7 SAT
Map Sheet PDP 2-8

Photograph Name _____ Number # 36+37 Row 3

Topographic Position (Circle the correct response)
☒ Upland Ridge ☐ Lower slope
☐ Upper slope ☒ Valley
☐ Midslope ☐ Washline

2. SITE DATA Circle the correct response

Slope flat ☒ gentle ☐ steep 2.5 Aspect 172° 5 - 11° -90
180 135
☐ N ☐ NE ☐ E ☒ SE ☐ S ☐ SW ☐ W ☐ NW

Surface Soil gritty loam. Depth to Rock 0-10cm Colour grey to dk brown

Exposed Rock type granite % surface cover 65-70% size range 20-50cm Broken-granite

Litter 10 % surface cover _____ Depth (cm) 2cm




Bare Ground 1% % surface cover _____

with rocks 70% 70%

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class					
	2-10%	10-30%	30-70%	over 70%	
TREES					
	over 15m	5-15m	under 5m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		1% - 2% ADS	5% (5m)		
DOMINANT SPECIES		E. calophylla for plot			
SHRUBS					
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)				30-70%	2-10%
DOMINANT SPECIES				Calyptra aculeata Grev. bipinnatifida	Hibbs? commutata/montana
GRASSES		HERBS	SEDGES	over 0.5m	
LIFE FORM					
COVER CLASS (%)	10-30%	30-70%			
DOMINANT SPECIES	Biza maximz Avena sp.	Styl. bulbiferus Chorizanthes aulis Hypochaeris glabra			

4. VEGETATION CONDITION

Comments x Weedy from adj. farmland - 3 1/2 4/5 (v. weedy!)
Recent drought - death of Allocas humilis
on ridge. v. weedy open & subject to intense
heat/exposure. ADS - Allocas. huegeliana
Labiachya punctata

3-3 1/2

3

SERP 02
PG 1

* Check Orthosandrus for shooting leaf bases

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP02

Record on Sheet

Date 23.10.96

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

check that grev. pilulifera is
other plant was not
tymnaria angust

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
E. calophylla	✓			Drosera pallidifolia?			✓	Melotia? - Asclepiaceae	✓		
				3 cauline leaves				unid. Scrophs? ①	✓		
SHRUBS				in whorl				Anagallis arvensis	✓		
Tymnaria latifolia	✓		✓				✓	Drosera menziesii			
Catylax aculeata?/acutifolia	✓		✓	Wahlenbergia sp.			✓	Ursinia anthemoides	✓		
Pyramnia quercifolia?	✓		✓					Daucus glochidiatus	✓		
MALLEES								unid. Scrophs ②	✓		
Hibbertia commutata/mortona				Medicago/Trifolium	✓		✓	fraespiphyllum (leaf orchid)			
Hemigenia incana	✓		✓	GRASSES				Drosera glandulifera			
Grevillea bipinnatifida	✓			Briza maxima	✓		✓	Leuechodia sp.			
Macrorhiza veidtii	✓			Avena sp. ? fabriz	✓		✓	trachymene pilosa			
SHRUBS				Romulea rosea	✓		✓	Isoplepis sp.			
Dodonaea sp. aratropa	✓		✓	Wahlenbergia Gaulis (check wood)			✓	Calamita clenchii?			
Thysanotus patersonii	✓		✓	Hemra Hemra			✓	SEDGES			
Didopeltis sp	✓		✓	Aera Calophylla			✓	Loxocaryx flexuosus?	✓		
Xanthosia candida	✓		✓	Unid small grass (Volpia)			✓	Lepodosperma sword leat.	✓		
Hemianthera pinnata				patersonia? Shooting leaf base (?)	✓		✓				
Arctostaphylos purpurea	✓		✓	Briza Minor	✓		✓	Pleurachne (AOT)?			
Hakea lasiocarpa lissocarpa	✓			HERBS				Lobelia (AOT)			
Baronia sp?	✓		✓	Cheilanthes austrotenuifolia	✓		✓				
Allocasuarina humilis?	✓		✓	Dioscorea hastifolia	✓		✓				
Hakea undulata	✓			Rhodanthe sp			✓				
Xanthosia preissii	✓			Stylidium bulbiferum	✓		✓				
Burchardia congesta	✓		✓	Caesia micrantha	✓		✓				
				Drosera stolonifera			✓				
				Hypochaeris glabra			✓				
				Unid. Poliose herb with 3 connate carpels			✓	Baronia!			
				Arthroopodium capillipes	✓		✓	Didopeltis	✓		
				Drosera - small							
				Arctostaphylos p. lesa							
				Antigonon antiquus. Apicose?							
				Tridentata leaf. H. site	✓		✓				

✓ off Hovea pungens??
check id.

Arctostaphylos
Pleurachne?

Thelymitra
crinita



BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP02

Record on Sheet

Date 13.12.96

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
								Yellow fw. <i>Lineum</i>			
								<i>Actinotus</i> sp			
								White everesting- <i>Myrtaceae</i>			
								(<i>Senecio</i>)			
								White <i>Scroph</i>			
								<i>Ranuncula vasezi</i>			
MALLEES								<i>Sonchus oleraceus</i>			
								<i>Medicago</i>			
				GRASSES				<i>Flouza pungens</i>			
								<i>Leucophaea purpurea</i>			
								<i>Boys sphacelophila</i>			
SHRUBS								<i>Asi</i>			
								SEDGES			
								Large <i>Hibbertia</i> !			
								<i>linata??</i>			
								<i>Gomph. marginatum</i>			
								Opp-located pink flw herbs			
				HERBS				check <i>Alloxas. hueglicana??</i>			
								replant			
								<i>Alloxas humilis</i> is			
								also checks			
								replant			
								Prob. <i>A. hueglicana</i>			

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SELP82

Record on Sheet

Date 23-10-96

13-12-96 (R)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Allocasuarina huegeliana</i>	✓		✓	NE <i>Prasophyllum</i> sp. leaf only				<i>Thysanotus manglicus</i>	✓		✓
<i>Euc. calophylla</i> (seedling)	✓		✓	<i>Drosera glanduligera</i> ?	✓		✓	<i>Hypochaeris glabra</i>	✓	✓	✓
↓ seed sampling!				<i>Gonolobum marginatum</i>	✓		✓	<i>Caesia micrantha</i>	✓	✓	✓
								<i>Trifolium campestre</i>	✓	✓	✓
								<i>Thelymitra viridis</i>	✓		✓
								<i>Drosera menziesii</i>	✓		✓
<i>Xanthorrhoea preissii</i>	✓		✓					<i>Drosera stolonifera</i>	✓		✓
<i>Hakea undulata</i>	✓		✓					<i>Cheilanthes australis</i>	✓		✓
<i>Macropia riedlii</i>	✓		✓	GRASSES				<i>Rhodanthe corymbosa</i>	✓	✓	✓
<i>Grevillea bipinnatifida</i>	✓		✓	<i>Briza minor</i>	✓		✓	<i>Pavementaria latifolia</i>	✓	✓	✓
NE <i>Hibbertia commutata</i> ?				<i>Mira canyophyllen</i>	✓	✓	✓	<i>Banksia mixago</i>	✓	✓	✓
<i>Allocasuarina humilis</i>	✓	✓	✓	<i>Vulpia muricoides</i>	✓	✓	✓	<i>Ursinia anthemoides</i>	✓	✓	✓
<i>Acacia pulchella</i>	✓	✓	✓	<i>Briza maxima</i>	✓	✓	✓	<i>Sonchus oleraceus</i>	✓		R
<i>Boronia racemosa</i>	✓	✓	✓	<i>Avena barbata</i>	✓	✓	✓	<i>Boronia spirocephala</i>	✓		R
<i>Hemigenia incana</i>	✓	✓	✓					<i>Xanthosia candida</i>	✓	✓	✓
<i>Diplopeltis huegelii</i> var <i>huegelii</i>	✓	✓	✓					<i>Stylidium bulbiferum</i>	✓	✓	✓
<i>Calceolaria acutifolia</i>	✓	✓	✓					<i>Actinotus leucocephalus</i>	✓	✓	✓
<i>Trimalium ledifolium</i> var <i>ros</i>	✓	✓	✓					<i>Centaurium euthyaca</i>	✓	✓	R
ADS: <i>Labechia lanceolata</i>	✓	✓	✓	HERBS				<i>Daucus glochidialis</i>	✓	✓	✓
<i>Dryandra quercifolia</i>	✓	✓	✓	ADS: <i>Trachymene coerulescens</i>	✓	✓	✓	<i>Trachymene pilosa</i>	✓	✓	✓
<i>Dodonaea ceratocarpa</i>	✓	✓	✓	<i>Dioscorea hastifolia</i>	✓	✓	✓	<i>Homalosciadium homalocarpum</i>	✓	✓	✓
<i>Hibbertia mylnei</i>	✓	✓	✓	<i>Linum trigynum</i>	✓	✓	✓	<i>Galium divaricatum</i>	✓	✓	✓
<i>Plowea pungens</i>	✓	✓	✓	NE <i>Xanthosia huegelii</i>	✓			<i>Pro. Arthropodium capillipes</i>	✓		✓
<i>Acacia pulchella</i>	✓		✓	<i>Fraxinaria ovata</i> ??	✓			SEDGES			
<i>Hakea lissocarpa</i>	✓		✓	<i>Wahlenbergia preissii</i>	✓	✓	✓	<i>Desmodium aspera</i>	✓	✓	✓
				<i>Levenhookia pusilla</i>	✓	✓	✓	<i>Isolepis marginatum</i>	✓	✓	✓
<i>Dryandra proemorsa</i>				ADS: <i>Pleurosorus rutifolius</i>	✓	✓	✓				
(<i>proemorsa</i> is				<i>Anagallis arvensis</i>	✓	✓	✓				
southern sandplains !!)				<i>Pomarea rosea</i>	✓	✓	✓				
				<i>Drosera macrantha</i>	✓	✓	✓				
				<i>Burchardia congesta</i>	✓	✓	✓				
				<i>Rhodanthe manglicus</i>	✓	✓	✓				

id=✓

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP SITE NUMBER SERP03

DATE TRIP 23.10.96 RECORDERS AM DL


DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Walkin on bearing!

Mud Map Draw a sketch of the location of the site below



Road Location * Walk in from survey track ~200m

Geographic Location Latitude 32 22 28.4 S Longitude 116 00 07.2 E Altitude (210) (650)
GPS ☒ Accuracy DDPP 3.3
Map Sheet Jawahdale

Photograph Name Number # 37 Roll # 3

Topographic Position (Circle the correct response)
Upland ☐ Lower slope ☐
Upper slope ☒ Valley ☐
Midslope ☐ Washline ☐

2. SITE DATA Circle the correct response

Slope flat ☐ gentle ☒ steep ☐ Aspect N NE E SE S SW W NW

Surface Soil clayey, gritty, loam Depth to Rock 30-7100 cm Colour Dk Brown-Grey

Exposed Rock type granite % surface cover 0.01 % size range 0-2-10 cm

Litter 780 % surface cover Depth (cm) 20-30



Bare Ground 0 % surface cover

Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class		2-10%	10-30%	30-70%	over 70%
TREES					
	over 15m	5-15m	under 5m	MALLEES over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		10-30% (30-40%)	10-30% 30-40%		
DOMINANT SPECIES		<i>E. calophylla</i>	<i>E. calophylla</i> <i>E. marginata</i>		
SHRUBS					
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	SHRUBS under 0.5m
LIFE FORM					
COVER CLASS (%)			10-30%	30-70%	30-70%
DOMINANT SPECIES			<i>A. pulchella</i> <i>H. hissarayda</i> <i>Ty. ledifolium</i>	<i>Xen. preissii</i> <i>Bossia ornata</i>	<i>Hyp. angust.</i> <i>Hibb. hyperi</i> <i>Dry. nivalis</i>
GRASSES					
	GRASSES	HERBS	SEDGES	over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)		thin sward			10-30%
DOMINANT SPECIES					<i>L. tenuis</i> (rough) <i>T. capillaris</i>

4. VEGETATION CONDITION

Comments low (3-8m) Savanna Mauri on edge of west-facing 1st ridge - overlooking foothills. Thick X. praeii ulstorey. Roo grazing. Some fine history & some Briza from adjacent farmland. score 4/5.
Dense ulstorey & good soil.

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP 03

Record on Sheet

Date 28.10.96

13.12.96(R)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Eucalyptus calophylla</i> ✓✓								<i>Tetradlea nuda</i> ✓✓			
<i>Eucalyptus marginata</i> ✓✓								<i>Comesperma ciliatum</i> ✓✓			✓R
								<i>Stylidium hispidum</i> ✓✓			✓R
<i>Hakea issocarpa</i> ✓✓			✓R					<i>Dampiera alata</i> ✓✓			✓R
<i>Gnaveillea pilulifera</i> ✓✓			✓R					<i>Burchardia congesta</i> ✓✓			✓
<i>Dryandra lindleyana</i> ✓✓			✓R					<i>Stylidium bulbiferum</i> ✓✓			✓R
<i>Boraginaceae</i> sp. <i>Navona</i> (id.) ✓✓			✓R								
<i>Leucopogon capitellatus</i> ✓✓			✓R	<i>Neurachne aboperculoides</i> ✓✓			✓R	<i>Paterosia babionoides</i> ✓✓			✓R
<i>Hypochaeris angustifolia</i> ✓✓			✓R	GRASSES				<i>Lomandra hermaphrodita</i> ✓✓			✓
<i>Xanthorrhoea gracilis</i> ✓✓			✓R	<i>Briza maxima</i> ✓✓			✓R	<i>Haemodorum</i> sp. <i>strepens</i> ✓✓			✓
<i>Acacia pulchella</i> ✓✓			✓R	<i>Ara canyophylla</i> ✓✓			✓R				
<i>Astroloma pallidum</i> ✓✓			✓R	<i>Tetradlea laevis</i> ✓✓			✓R	<i>Haemodorum</i> sp. <i>narrow leaf</i> ✓✓			✓
<i>Chorizanthe dicksonii</i> ✓✓			✓R	<i>Drya pycnostachya</i> ✓✓			✓R				
<i>Leucopogon pulchellus</i> ✓✓			✓R	<i>Danthonia setacea</i> ✓✓			✓R	<i>Lagenifera huegelii</i> ✓✓			✓
<i>Olearia paucidentata</i> ✓✓			✓R	<i>Micraena stipitoides</i> ✓✓			✓R				
<i>Hibbertia hypericoides</i> ✓✓			✓R	<i>Briza minor</i> ✓✓			✓				
<i>Leucopogon propinquus</i> ✓✓			✓R	<i>Dirichlachne crinita</i> ✓✓			✓				
<i>Baekelandia campylosma</i> ✓✓			✓R	<i>Vulpia</i> sp. ✓✓			✓				
<i>Tynantrum ledifolium</i> var. <i>rosm.</i> ✓✓			✓R	HERBS							
<i>Boraginaceae</i> sp. <i>ovata</i> (id.) ✓✓			✓R	<i>Opervularia apiciflora</i> ✓✓			✓R				
<i>Hibbertia communata</i> ✓✓			✓R	<i>Stackhousia monogyna</i> ✓✓			✓R				
				<i>Scaevola calliptera</i> ✓✓			✓				
<i>Lechenanthea biloba</i> seedling? ✓✓			✓R	<i>Leueihostia puella</i> ✓✓			✓R	SEDGES.			
<i>Hovea charizemifolia</i> ✓✓			✓	<i>C. micrantha</i> ? single leaf form ✓✓			✓R	<i>Tetradlea extensa</i> ✓✓			✓R
<i>Xanthorrhoea pleissii</i> ✓✓			✓R	<i>Cassia micrantha</i> ✓✓			✓R	<i>Tetradlea capillaris</i> ✓✓			✓R
<i>Macroglossa nedllei</i> ✓✓			✓R	<i>Drosera erythrorhiza</i> ✓✓			✓	<i>Loxorhiza fasciculata</i> ✓✓			✓R
				<i>Anthropodium capillipes</i> ✓✓			✓R				
				<i>Dianella revoluta</i> ✓✓			✓R				
				<i>Cassutha paniformis</i> ✓✓			✓R				
				<i>Lomandra nigricans</i> ✓✓			✓R				
				<i>Stylidium brunonianum</i> ✓✓			✓				
				<i>Cassutha glabella</i> ✓✓			✓R				

<i>Arenophryne rotunda</i> , <i>Pseudophryne guentheri</i> & <i>Uperoleia russelli</i>	PSEUDOPH
<i>Neobatrachus</i> sp_B, <i>N. sp_C</i> & <i>N. wilsmorei</i>	NEOSP_AA
GECKOS	
<i>Diplodactylus alboguttatus</i> A & <i>D. alboguttatus</i> B	DIPALBOG
<i>Diplodactylus pulcher</i> A & <i>D. pulcher</i> B	DIPPULCH
<i>Diplodactylus rankini</i> , <i>D. spinigerus</i> & <i>D. strophurus</i>	DIPLOD_1
<i>Diplodactylus squarrosus</i> & <i>Rhynchoedura ornata</i>	DIP_RHYN
<i>Gehyra punctata</i> & <i>G. variegata</i>	GEHYRA
PYGOPIDS	
<i>Aprasia fusca</i> , <i>A. haroldi</i> & <i>A. smithi</i>	APRASIA
<i>Delma australis</i> A, <i>D. australis</i> B, <i>D. butleri</i> , <i>D. nasuta</i> , <i>Pygopus nigriceps</i> & <i>Pygopus lepidopodus</i>	DEL_PYGO
DRAGONS	
<i>Ctenophorus badius</i> & <i>C. maculatus</i>	CTENOP_1
<i>Ctenophorus femoralis</i> & <i>C. scutulatus</i>	CTENOP_2
<i>Ctenophorus inermis</i> & <i>C. reticulatus</i>	CTENOP_3
<i>Ctenophorus rubens</i> A & <i>C. rubens</i> B	CTENOP_4
<i>Tympanocryptus parviceps</i> & <i>T. adelaidensis</i>	TYMPANOC
SKINKS	
<i>Cryptoblepharus carnabyi</i> & <i>C. plagiocephalus</i>	CRYPTOBL
<i>Ctenotus alleni</i> , <i>C. fallens</i> , <i>C. lesueuri</i> , <i>C. mimetes</i> , <i>C. saxatilis</i> & <i>C. severus</i>	CTENOT_1
<i>Ctenotus hanloni</i> A, <i>C. hanloni</i> B & <i>C. uber</i>	CTENOT-2
<i>Cyclodomorphus celatus</i> , <i>C. melanops</i> & <i>Eremiascincus fasciolatus</i>	CYC_ERE
<i>Egernia depressa</i> & <i>E. stokesii</i>	EGERNIA
<i>Lerista connivens</i> , <i>L. kendricki</i> A, <i>L. kendricki</i> B & <i>L. varia</i>	LERIST_1
<i>Lerista decora</i> & <i>L. planiventralis</i>	LERIST_2
<i>Lerista elegans</i> , <i>L. haroldi</i> , <i>L. muelleri</i> A, <i>L. muelleri</i> B, <i>L. muelleri</i> C & <i>L. muelleri</i> D	LERIST_3
<i>Lerista humphriesi</i> , <i>L. praepedita</i> A & <i>L. praepedita</i> B	LERIST_4
<i>Lerista kennedyensis</i> & <i>L. uniduo</i>	LERIST_5
<i>Lerista lineopunctulata</i> A & <i>L. macropisthopus</i>	LERIST_6
<i>Menetia amaura</i> , <i>M. greyi</i> A, <i>M. greyi</i> B & <i>M. greyi</i> C	MENET_1
<i>Menetia surda</i> A, <i>M. surda</i> B & <i>Notoscincus ornatus</i>	MENET_2
<i>Morethia lineocellata</i> & <i>M. obscura</i>	MORETH_2
<i>Tiliqua multifasciata</i> , <i>T. occipitalis</i> & <i>T. rugosus</i>	TILIQUEA
GOANNAS	
<i>Varanus brevicauda</i> , <i>V. caudolineatus</i> & <i>V. eremius</i>	VAR_SMAL
SNAKES	
<i>Demansia calodera</i> & <i>D. psammophis</i>	SNAKE_S1
<i>Denisonia fasciata</i> , <i>Furina ornata</i> , <i>Pseudonaja modesta</i> & <i>V. bimaclata</i>	SNAKE_S2
<i>Psedechis australis</i> & <i>Pseudonaja nuchalis</i>	SNAKE_L1
<i>Ramphotyphlops australis</i> , <i>R. grypus</i> A, <i>R. grypus</i> B, <i>R. grypus</i> C & <i>R. hamatus</i>	SNAKE_BL
<i>Vermicella bertholdi</i> , <i>V. littoralis</i> & <i>V. multifasciata</i>	SNAKE_S3

Classification of the 67 ecological taxa according to similarities in their co-occurrences at the same quadrats yielded similar assemblage boundaries to those derived from the species analysis (Fig. DENT2):

- Assemblage_1: Taxa characteristically restricted to hummock grass communities on the red sandplains and dune fields of the arid zone, with ranges extending across middle and tropical latitudes.

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP03

Record on Sheet

Date 23.10.96

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
E. marginata	✓		✓	Howea charizmiifolia ? (NE)			✓	Cassytha glabelliz	✓		✓
E. calophylla	✓		✓	Lepidosperma angustatum			✓	Cassytha racemosa	✓		✓
				Vulpia sp. - small grass			✓	Agrostocinium 2 sp			✓
SHRUBS							✓	Tet. nuda leafless // scale buds			✓
Howea lissocarpha	✓		✓	Strophanthus sp			✓	Stemodia capillaris (repro + form)			✓
Xan. preissii	✓		✓	Lepidosperma - Smooth, terete			✓	Chorizandra dicksonii			✓
MALLEES							✓	Loxocarpus fasciculata	✓		✓
Dry. nivea	✓		✓				✓	Stylidium piliiforme	✓		✓
Ribbertia hypericoides	✓		✓	GRASSES			✓	Lomandra haemophrodita (male)	✓		✓
Leucopogon pulchellus	✓		✓	Tetraria laevis	✓		✓	Unid grass (hirsute leaves)	✓		✓
Boschia ornata / aiocarpa	✓		✓	Stipa ? otterantissima	✓		✓	Las coll. before	✓		✓
SHRUBS				Briza maxima	✓		✓	Loganiifera huegii			✓
Hibb. commutata	✓		✓	Danthonia sp.	✓		✓	Lachenalia pucilla			✓
Hypocalymma angustifolium	✓		✓	Lenartia presonii	✓		✓	SEDGES			✓
Comesperma virginatum	✓		✓	Tetraria capillaris			✓	Stylidium piliiforme	✓		✓
Acacia pulchella	✓		✓	Briza minor - collected duck			✓				✓
Macropodaria weidlii	✓		✓	Lomandra - narrow leafed - curled -	✓		✓	caespitosa ! Hemaph			✓
Xanth. gracilis	✓		✓	Tetraria ostendia	✓		✓				✓
Dalmanella alata	✓		✓	HERBS			✓	Heavy ligule	✓		✓
Dianella revoluta	✓		✓	Arthropodium capillaris	✓		✓				✓
Tymalium ledifolium	✓		✓	Burchardia longata	✓		✓				✓
Oleracea paucidentata	✓		✓	Cassia micrantha	✓		✓				✓
Petersonia babooides	✓		✓	Stylidium bulbiferum	✓		✓				✓
Astiloma palladium	✓		✓	Haemodorum sp. - stop leaf. - broad	✓		✓				✓
Scaevola sp. - hirsute			✓	Dracopis erythrina	✓		✓				✓
Baeckea	✓		✓	Stylidium bichenianum - brunonianum	✓		✓				✓
Styphelia tenuifolia	✓		✓	Haemodorum - narrow leaf			✓				✓
Styphelia tenuifolia	✓		✓	Opercularia fastigata	✓		✓				✓
				Lechanactia biloba - (NE)	✓		✓				✓
				Lepidosperma - terete, glabrous - shaded areas			✓				✓
				Nurac. la. Neurachne alpecearoides	✓		✓				✓
				Aira calophylla - calophyllae			✓				✓

Loxoc
fascic
↓
fascic

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE	
SITE No	SFRP03
Date	13.12.96
	Rescore

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP03

Date 13.12.96

Restore

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

checkid of Tet. capillaris!

[illegible]

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP. ^{→ Survey track on road ridge} SITE NUMBER SERP04


DATE TRIP 24.10.96 RECORDERS AM DL

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



^{CHECK ODOMETER} Road Location Survey Track - C

Geographic Location Latitude 32° 22' 30.8" S Longitude 116° 00' 10.2" E Altitude 170m
GPS ☒ Accuracy PDOP = 3.5
Map Sheet Savannah

Photograph Name _____ Number #2 Roll # 4

Topographic Position (Circle the correct response)
Upland - Ridge line 4 Lower slope
Upper slope Valley
Midslope Washline

2. SITE DATA Circle the correct response

Slope flat gentle steep Aspect N NE E SE S SW W NW

Surface Soil gitty loam Depth to Rock 0-20 cm Colour DK Brown

Exposed Rock type granite % surface cover 15 size range 5cm - 40cm

Litter 780 % surface cover 2-3 layers Depth (cm) 5-10cm


Bare Ground 0 % surface cover (15% ind. rock)


BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class	2-10%	10-30%	30-70%	over 70%
-------------	-------	--------	--------	----------

TREES			MALLEES	
	over 15m	5-15m	under 5m	
				over 8m
				under 8m
LIFE FORM				
COVER CLASS (%)			2-10%	
DOMINANT SPECIES		<i>E. calophylla</i>	<i>E. calophylla</i> <i>E. marginata</i> <i>L. ligno. taproot</i>	

SHRUBS			SHRUBS	
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m
LIFE FORM				
COVER CLASS (%)	> 70%	> 70%		3 > 70%
DOMINANT SPECIES	<i>X. acanthostachys</i>	<i>X. acanthostachys</i> <i>H. undulata</i> !		<i>H. undulata</i> <i>Melaleuca scutellaria</i> <i>A. pulchellus</i>

GRASSES		HERBS		SEDGES	
				over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)					30-70%
DOMINANT SPECIES		<i>T. sp.</i> <i>Acrid. leaf layer</i>			<i>Tetraria capillaris</i> <i>Mesomelaena tetragynis</i>

4. VEGETATION CONDITION

Comments 4-5/5! Excellent condition. Few weeds (B. minor)
Long unburnt - large stand of *H. undulata* indicates
a 5-8 y.o. hot fire which has germinated dense
thicket of *H. undulata* on ridge.

15

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP04

Record on Sheet

Date 24.10.96

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
E. calophylla	✓		✓	unid small to sprout - styl				Cassutha racemosa	✓		✓
E. marginata	✓		✓	Rhynchospora (Ternstroemia) pycno.				Drosera menziesii			✓
				or G. pilulifera			✓	Trachymene pilosa			✓
SHRUBS				Daviesia pruri decurrens (dood			✓	Hudcotea			✓
Hypo. angustifolium	✓		✓	prob summer drought)			✓	Xanthosia heuglii			✓
Hibbertia spicata	✓		✓	alive!			✓	Tetraria capillaris			✓
MALLEES				D. decurrens			✓	Capillaris			✓
Xanth. aranthostachya	✓		✓					Caesia micrantha			✓
Hibb. commutata	✓		✓	GRASSES				unid Epacrid seedling			✓
Melaleuca scabra	✓		✓	Climbing Drosera				prob L. pulcherrimus			✓
Dry. nuda	✓		✓	D. menziesii?			✓	Isopogon asper	✓		✓
SHRUBS				Drosera pallida				Melaleuca sp.			✓
Petrophile striata	✓		✓	Unid grass sp 1			✓	Caladenia flava			✓
Hakea stenocarpa	✓		✓	Tetraria acutifolia			✓	SEDGES			
Hakea undulata	✓		✓	Ascoloma palidum			✓	Loxocarya fasciculata			✓
Allorauhinia humilis	✓		✓	Homma homma				Lomandra preissii			✓
Macrozamia veitchii	✓		✓	Cassutha racemosa - glabrous??			✓	Lepidosperma tenuis (terre)	✓		✓
Leucophaea pulcherrimus	✓		✓	Briza maritima			✓	Cassutha sp (probably racemosa)			✓
Sphaerobolium medium	✓		✓	HERBS				Maesomeria tetragona	✓		✓
Hibb commutata	✓		✓	numerous leaf bases			✓	Styphelia sp / may be astroloma			✓
Gomph marginatum			✓	Haemodorum paniculatum			✓	Unid epacrid			✓
Hakea lissocarpa	✓		✓	Thysanotus patersonii			✓	Briza minor			✓
Azara pulchella seedling	✓		✓	Agrostocrinum scaberrimum			✓	Burchardia congesta			✓
Lambertia hyperacoides	✓		✓	Stylidium bulbiferum			✓	Lomandra Lepidosperma angustatum			✓
Synaphia petiolaris	✓		✓	stylidium			✓	Tetraria lavis			✓
Acacia drummondii	✓		✓	Stylidium piliferum			✓	Chaenactis Chaenactis			✓
Ranunculus rosea			✓	Stylidium calcitratum			✓	Chaenactis complanata			✓
Vereta leaf Thysanotus			✓	Stylidium brunneum			✓	Tripterococcus sp brunneus	✓		✓
Single leaf Lomandra ??			✓	Thalysandra cincta			✓	Small Euphorb			✓
Dampiera Dampiera			✓	stylidium calosum			✓	Willin bergia Wahlenbergia			✓
Hamamelis - very young leaf	✓		✓	Unid Small herb elliptical leaves			✓	Neuracne elapic	✓		✓
Hypochaeris glabra			✓	Scirpus sp (not flowering)			✓	Stipa sp.			✓
			✓	Short climbing Drosera			✓				

Baekea camphorosmea ✓

Abalastium ? - Agrostocrinum?

Acanthopneum (dood) - Acanthopneum

Actual 370

attracting small insects

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP004

Record on Sheet

Date 13-12-96

- Column 1 plant name

- Column 2 plant number

• **Column 3** **flowering time-** TICK if species flowering

• **Column 4** **identification check**

[illegible]

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP04

Record on Sheet

Date 24.10.96

13.12.96 (R)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES				SHRUBS (cont.)				HERBS (cont.)			
No	FI	ID		No	FI	ID		No	FI	ID	
<i>Eucalyptus wandoo</i>	✓		✓	<i>Haemodorum discolor</i>	✓	✓	✓	<i>Borrichia microphylla</i>	✓	✓	✓
<i>Euc. marginata elegantella</i>	✓		✓	<i>Xanth. acanthostachya</i>	✓	✓	✓	<i>Ten. tripterocarpus brunneus</i>	✓	✓	✓
				<i>Hakea stenocarpa</i>	✓	✓	✓	<i>Hyaloperma rotula</i>	✓		✓
<i>Isopogon scoparius</i>	✓		✓	<i>Macroglossa viedii</i>	✓	✓	✓	<i>Millobia tenuifolia tenuifolia</i>	✓	✓	✓
<i>Hakea undulata</i>	✓		✓	<i>Hakea lissocarpa</i>	✓	✓	✓	<i>Drosera macrantha</i>	✓	✓	✓
<i>Astroloma ciliatum</i>	✓		✓	<i>Thysanotus hirsutius</i>	✓	✓	✓	<i>Cassytha glabella</i>	✓	✓	✓
<i>Acacia pulchella pulchella</i>	✓		✓	<i>Acanthocarpus preissii</i> (dead)	✓	✓	✓	<i>Drosera pinguiculi</i>	✓	✓	✓
<i>Leucopogon pulchellus</i>	✓	✓	✓	<i>Caladenia flava</i>	✓	✓	✓	<i>Stylidium hispidum</i>	✓	✓	✓
<i>Hibbertia commutata</i>	✓	✓	✓	GRASSES				<i>Thysanotus marginatus</i>	✓	✓	✓
<i>Hypocalymma angustifolium</i>	✓		✓	<i>Bipa maxima</i>	✓	✓	✓	<i>Thelymitra cincta</i>	✓	✓	✓
<i>Daboia decurrens</i>	✓		✓	<i>Stipa pycnostachya</i>	✓	✓	✓	<i>Stylidium pycnostachyum</i>	✓		✓
<i>Petrophile striata</i>	✓		✓	<i>Neurachne alpepercurioides</i>	✓	✓	✓	<i>Hydrocotyle ptilifera</i>	✓		✓
<i>Argemone ledifolium var.</i>	✓		✓	<i>Stipa complexa</i>	✓			<i>Homaloscladium homalocarpum</i>	✓		✓
<i>Acacia lateritica</i>	✓		✓	<i>Vulpia muricoides</i>	✓			<i>Hypochaeris glabra</i>	✓	✓	✓
<i>Hibbertia hypericoides</i>	✓	✓	✓	<i>Danthonia setacea</i>	✓	✓	✓	<i>Stylidium calcareatum</i>	✓	✓	✓
<i>Leucopogon capitatus</i>	✓	✓	✓	<i>Bipa minor</i>	✓	✓	✓	<i>Lomandra preissii</i>	✓	✓	✓
<i>Baeckea camphoresmae</i>	✓		✓	<i>Tetrarrhena laevis</i>	✓	✓	✓	<i>Chaenactis coronata</i>	✓	✓	✓
<i>Astroloma pallidum</i>	✓		✓	HERBS				<i>Stylidium brunneum</i>	✓		✓
<i>Metaleuca aff. scabra</i>	✓		✓	<i>Xanthost. luegelii</i>	✓	✓	✓	<i>Haemodorum sp. laet</i>	✓		✓
<i>Allocasuarina humilis</i>	✓		✓	<i>Burchardia longista</i>	✓	✓	✓	<i>Cassytha pinnatifida pinnatifida</i>	✓	✓	✓
<i>Synaphea gracillima</i>	✓		✓	<i>Stylidium pulchellum</i> (dead)	✓	✓	✓	<i>Haemodorum laetum</i>	✓	✓	✓
<i>Synaphea sp. indet</i>	✓		✓	<i>Caesia micrantha</i>	✓	✓	✓	<i>Stylidium carnosum</i>	✓		✓
<i>Hibbertia spicata</i>	✓	✓	✓	<i>Romulea rosea</i>	✓	✓	✓	SEDGES			
<i>Gompholobium marginatum</i>	✓		✓	<i>Scaevola calliptera</i>	✓	✓	✓	<i>Tetaria octandra</i>	✓	✓	✓
<i>Grevillea pilulifera</i>	✓		✓	<i>Agrostocrinum scabrum</i>	✓	✓	✓	<i>Tetaria capillaris</i>	✓	✓	✓
<i>Sphaerolobium medium</i>	✓		✓	<i>Cassytha pinnatifida</i>	✓	✓	✓	<i>Mesomelaena tetragona</i>	✓	✓	✓
<i>Dyandra lindleyana lindleyana</i>	✓		✓	<i>Lomandra caespitosa</i>	✓	✓	✓	<i>Loxocarya fasciculata</i>	✓	✓	✓
<i>Hypocalymma robustum</i> (hike)	✓		✓	<i>Pithocarpa pulchella</i>	✓	✓	✓				
<i>Acacia lasiocarpa</i>	✓		✓								
<i>Acacia pulchella</i>	✓		✓	<i>Trachymene pilosa</i>	✓	✓	✓				
				<i>Hydrocotyle callicarpa</i>	✓	✓	✓				

Wahlenbergia - - -

- Assemblage_2: Taxa of semi-arid to arid temperate areas of the west coast and southern half of Australia, that favour the variety of sandy surfaces (hummock grasslands and shrublands). Some taxa are confined to this coastal strip.
- Assemblage_3: Widespread arid to semi-arid taxa, many of which are confined to finer textured soils, from loamy-sands and sandy-loams to alluvial plains and even samphire pans.
- Assemblage_4: Singleton taxa normally associated with rock surfaces (*Ctenophorus caudicinctus*, *Morelia perthensis* and *Morethia ruficauda*) were previously included in Assemblage_4 (Fig. DENT) because of the ecotonal position of the quadrat GJ1: low on the scree slope of the Pell range, 0.1 km from the floodways and levee banks of the Gascoyne River.
- Assemblage_5: Species restricted to flood plains and riverine flats.
- Six assemblages comprising one or two taxa that were recorded on only one or two quadrats.

In Figure DENT2, the 63 quadrats have been re-ordered in terms of similarities in their taxa-composition. The sorting strategy yielded a quadrat-dendrogram that summarised the herpetofaunal patterns of occurrence. As previously, univariate data on aspects of the quadrat's physical environments were superimposed as a histogram on this dendrogram, attribute-by-attribute, to identify the physical attributes that most closely conformed to the herpetofaunal patterns.

PwetQ and TdiurR provided the best fits, and K(HCO₃), sand, carbon and electrical conductivity (EC) were also significant (**Fig Histo-ecol**). Pseas, PcldQ, Pann and TwarQ showed the same relationship to the herpetofaunal patterns as PcldQ, with which they are strongly intercorrelated (**Table Ecol_KW**). Similarly, TannR and Longitude showed the same relationship to the herpetofaunal patterns as TdiurR, with which they are both strongly inter-correlated. This is precisely the same set of attributes as emerged from the species-based analysis.

Table Ecol-KW. Kruskal-Wallis values for physical attributes that showed significant (<0.05) differences between groups of quadrats that had been clustered in terms of similarities in their ecological-taxa composition (**Fig. Histo-ecol**).

Attribute	Kruskal-Wallis value		Kendall's Tau (prob.) from Appendix 2.
	6-groups	15 groups	
PwetQ	49	56	
PcldQ	46	54	0.95 (***)
Pseas	49	55	0.81 (***)
TwarQ	41	49	- 0.84 (***)
Pann	37	47	0.74 (***)
TdiurRa	35	55	- 0.46 (***)
Longitude	39	51	0.80 (***)
TannRa	33	54	0.95 (***)
Altitude	25	46	0.58 (***)
TcldQ	5 ns ^a	46	- 0.27 (**)
Latitude	11 ns	45	- 0.81 (***)
K(H ₂ CO ₃)	32	40	- 0.34 (***)

(32 22 41.0 116 00 11.7
7 SATS → PDOP = 3.1 ! 210m)

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP: Survey track on western ridge SITE NUMBER SERP05


DATE TRIP 24 10 96 RECORDERS AM DL

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location

Geographic Location Latitude 32 22 41.0 S Longitude 116 00 12.6 E Altitude 180m
GPS ☒ Accuracy (6 SATS)
Map Sheet Jawandalo PDOP 4.2 (4.2)

Photograph Name _____ Number #3 Roll # 4

Topographic Position (Circle the correct response)
Upland _____ Lower slope _____
Upper slope 4 Valley _____
Midslope _____ Washline _____

2. SITE DATA Circle the correct response

Slope flat gentle steep Aspect N NE E SE S SW (W) NW

Surface Soil loamy clay Depth to Rock 10-100 Colour RED

Exposed Rock type granite % surface cover >1 size range 12-3 layers

Litter 60-70% % surface cover _____ Depth (cm) 10-20 cm




Bare Ground 1 % surface cover _____

Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class						2-10%	10-30%	30-70%	over 70%
TREES				MALLEES					
	over 15m	5-15m	under 5m	over 8m	under 8m				
LIFE FORM								30m	10m
	2-10%	10-30%							
	E. wandoo	E. wandoo							
SHRUBS						SHRUBS			
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	under 0.5m				
LIFE FORM								preissi 2m	1m
			2-10%	2-10%	770%				
			H. trifurcata	P. biloba	Phy. cal				
			H. undulata (1° dard)	H. lissocarpa	Hypo. angust				
GRASSES						HERBS			
	SEDGES		over 0.5m		under 0.5m				
LIFE FORM									
			2-10%		2-10%				
			Sty. pander glauca		Loricarya flexuosa				
			Orithosandrus sp						

4. VEGETATION CONDITION

Comments: Reasonable condition. ADS to road but other areas too upland -
4/5th 4/5
Dead H. undulata stand
dying (sensing)
Emergent Alloc. neugliana

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIFIC PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERPOS

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Date 24.10.96

13.12.96(12)

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Eucalyptus wandoo</i>	✓		✓					<i>Monadenia bracteosa</i>	✓	✓	✓
<i>Allocasuarina luzeleana</i>			✓					<i>Haemodorum</i> sp. indet	✓		✓
								<i>Cenosteus aculeata preissii</i>	✓		✓
								<i>Liximia anthemoides</i>	✓	✓	✓
<i>Hakea undulata</i>	✓	—	✓					<i>Romulea rosea</i>	✓	✓	✓
<i>Hakea lisocarpa</i>	✓	—	✓					<i>Stypandra glauca</i>	✓		✓
<i>Hakea v. furcata</i>	✓	—	✓					<i>Orthrosanthus latus</i>	✓	✓	✓
<i>Hibbertia spicata</i>	✓	✓	✓					<i>Galium murale</i>	✓	✓	✓
<i>Hibbertia commutata</i>	✓	✓	✓	GRASSES				<i>Cassytha</i> sp.			
<i>Acacia dentifera</i>	✓	✓	✓	<i>Briza minor</i>	✓	—	✓	<i>Stylidium bulbiferum</i>	✓	✓	✓
<i>Leucopogon pulchellus</i>	✓		✓	<i>Briza maxima</i>	✓	✓	✓	<i>Cheilanthes australis</i>	✓		✓
<i>Hypochilymna angustifolium</i>	✓		✓	<i>Microlaena stipoides</i>	✓		✓	<i>Cassytha perniformis</i>	✓	✓	✓
<i>Petrophile biloba</i>	✓		✓	<i>Vulpia myuros</i>	✓		✓	<i>Wahlenbergia</i>			
<i>Phyllanthus calycinus</i>	✓		✓	<i>Aira canophylla</i>	✓		✓				
<i>Acanthocarpus preissii</i>	✓		✓	<i>Lolium rigidum</i>	✓		✓	<i>Lomandra</i> sp.	✓		✓
<i>Grevillea pilulifera</i>	✓		✓	<i>Bromus hordeaceus</i>	✓		✓				
<i>Hibbertia myrtilloides</i>	✓	—	✓								
<i>Hibbertia ? myrtilloides</i>	✓	—	✓	HERBS							
<i>Xanthorrhoea preissii</i>	✓	—	✓	<i>Centaurea erythraea</i>	✓	✓	✓				
<i>Macropidia viedii</i>	✓	—	✓	<i>Daucus glaberrima</i>	✓	✓	✓				
<i>Grevillea bipinnatifida</i>	✓	—	✓	<i>Scaevola calyptera</i>	✓	✓	✓				
<i>Acacia pulchella pulchella</i>	✓	—	✓	<i>Stachys monogyna</i>	✓	✓	✓				
<i>Antersonia</i> sp. ?	✓	—	✓	<i>Banksia trixago</i>	✓	✓	✓	SEDGES			
<i>Trinailium ledifolium</i> var. <i>ros</i>	✓	—	✓	<i>Hypochaeris glabra</i>	✓	✓	✓	<i>Desmodium aspera</i>	✓	✓	✓
(from SERP4)				<i>Thysanotus angustatus</i>	✓	✓	✓				
				<i>Linum trigynum</i>	✓	✓	✓				
				<i>Tolpis barbata</i>	✓	✓	✓	as in ARSCS 10.12.96			
				<i>Oxalis perennans</i>	✓	✓	✓				
				<i>Drosera rotariifera</i>	✓	✓	✓				
				<i>Anagallis arvensis</i>	✓	✓	✓				
				<i>Sorbus oleraceus</i>	✓	✓	✓				

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

SITE No SERP05

Date 24.10.96

Label each plant with plants number, site code, date and plant's name or working name if required

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
E. laevis	✓		✓					Haemodorum sp. - 1st 1st	✓		✓
Allocas. huegeliana	✓		✓					step back			✓
SHRUBS								Anagallis arvensis	✓		✓
Petrophile biloba	✓		✓					Grass			
Hakea trifurcata	✓		✓					Aira calyptrifolia			
Hakea lissocarpa	✓		✓								
MALLEES											
Hakea undulata	✓		✓								
Acacia sp. long, narrow phyllodes	✓		✓	GRASSES							
↳ globular inflorescence				Briza maxima	✓		✓				
				Briza minor	✓		✓				
SHRUBS				aria grass - wide lamina	✓		✓				
Hibbertia spicata	✓		✓	bleary leaves	✓		✓				
Hypocalymma amplexifolium	✓		✓	hemispherical	✓		✓	SEDGES			
Acanthocarpus preissii	✓		✓	Sonchus oleraceus	✓		✓	Loxocarya flexuosa	✓		✓
Phyllanthus calycinus	✓		✓	Oxalis sp.	✓		✓				
Xanthorrhoea preissii	✓		✓	Small annual grass			✓				
Macrozamia reidii	✓		✓	Brigula-like grass			✓				
Amelites bipinnatifida	✓		✓	HERBS							
Hibbertia commutata	✓		✓	Styphandra glauca	✓		✓				
Hypochaeris glabra			✓	Conostylis aculeata	✓		✓				
Leucopogon pulchellus	✓		✓	Drosera rotundifolia			✓				
Wahlonbergia - ?			✓	Antennaria sp.	✓		✓				
Acacia pulchella	✓		✓	Stylidium bulbiferum	✓		✓				
Andersonia sp.				Thysanotus patersonii	✓		✓				
Trimalium				chrysanth.			✓				
				Helianthus austrotenuifolia	✓		✓				
				ASCINIA anthemoides	✓		✓				
				Cassipoupa racemosa	✓		✓				
				Lichotris - Miconia - Bonadenia			✓				
				Stackhousea	✓		✓				
				Daucus glychidifolius	✓		✓				

Urosia anthemoides

daucus glychidifolius

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No JERP05

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Date 13-10-96

[illegible]

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP SITE NUMBER SERP06


DATE TRIP 24-10-96 RECORDERS AM DL

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location Scrivener Road Flagged Wandoo if Greittler under chinaroo

Geographic Location Latitude 32 23 04.5 S Longitude 116 00 15.8 E Altitude 150m
GPS 03.9 Accuracy PDOP = 2.7
Map Sheet Jannahdale

Photograph Name _____ Number # 4 Roll # 4

Topographic Position (Circle the correct response)
Upland _____ Lower slope _____
Upper slope 4 Valley _____
Midslope _____ Washline _____

2. SITE DATA Circle the correct response

Slope flat gentle steep Aspect N NE E SE S SW W NW

Surface Soil gitty, clayey loam Depth to Rock 1-20 Colour light brown dark brown (chocolate)

Exposed Rock type granite % surface cover 30% size range 2 granite outcrop

Litter 15 % surface cover Depth (cm) 2-5cm

Bare Ground ind. vodes = 35% % surface cover

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class 2-10% 10-30% 30-70% over 70%

TREES				MALLEES	
	over 15m	5-15m	under 5m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)					
DOMINANT SPECIES					

SHRUBS				SHRUBS	
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)			10-30%	70%	
DOMINANT SPECIES			<i>G. enderlicheriana</i>	<i>Melaleuca fulgens</i>	
				<i>Darwinia citriodora</i>	

GRASSES		HERBS		SEDGES	
	over 0.5m		over 0.5m		under 0.5m
LIFE FORM					
COVER CLASS (%)	2-10%	2-10%			
DOMINANT SPECIES	<i>Biza maxima</i>	<i>Stylidium bulbosum</i>			
	<i>Ara canyulize</i>	<i>Medanthes austro-</i>			
	<i>Nulpa sp.</i>	<i>Dioscorea hirsutifolia</i>			
		<i>Scirpus</i>			

4. VEGETATION CONDITION

Comments: Granite heath. Some disturbance → assoc. with roadworks → 4/5. Good condition. Some annual weeds.

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP86

Record on Sheet

Date 13.12.96 (A)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Eucalyptus wandoo</i> (seedling)	✓	—	1/2					<i>Oxalis perennis</i>	✓	✓	
								<i>Galium divaricatum</i>	✓	✓	✓
								<i>Cheilanthes australensis</i>	✓	✓	✓
								<i>Xanthosia candida</i>	✓	✓	1/2
								<i>Sonchus oleraceus</i>	✓	✓	1/2
<i>Pimelea</i> sp. <i>imbricata</i> ?								<i>Trifolium cernuum</i>	✓	✓	1/2
<i>Grevillea endlicheriana</i>	✓	✓	1/2					<i>Anagallis anensis</i>	✓	✓	1/2
<i>Myrtalea ledifolium</i> var.	✓	✓	R					<i>Centaurea erythraea</i>	✓	✓	✓
<i>Labiichea lanceolata</i>	✓	✓	1/2	GRASSES				<i>Potentilla latifolia</i>	✓	✓	1/2
<i>Acacia encinophylla encinophylla</i>	✓	✓	R	<i>Neurachne alopecuroides</i>	✓	✓	1/2	<i>Wahlenbergia pressii</i>	✓	✓	✓
<i>Darwinia citriodora</i>	✓	✓	1/2	<i>Vulpia bromoides</i>	✓	✓	1/2	<i>Lotus</i> sp. sterile	✓	✓	✓
<i>Davasia horrida</i>	✓	✓	1/2	<i>Aira caryophylla</i>	✓	✓	1/2	<i>Centropus drummondiana</i>	✓	✓	✓
<i>Hovea pungens</i>	✓	✓	1/2	<i>Aster sp. barbata</i>	✓	✓	1/2	<i>Kennedia stirlingii</i>	✓	✓	✓
<i>Metaleuca radula</i>	✓	✓	1/2	<i>Biza maxima</i>	✓	✓	1/2	<i>Trachymene pilosa</i>	✓	✓	✓
<i>Myrtalea ledifolium</i> seedling - could be T. led.	✓	✓	1/2	<i>Danthonia setacea</i>	✓	✓	1/2	<i>Dracopis micrantha</i>	✓	✓	✓
<i>Acacia pulchella pulchella</i>	✓	✓	1/2	<i>Vulpia myuros</i> (nilae)	✓	✓	1/2	<i>Monadenia bracteosa</i>	✓	✓	1/2
<i>Diplopeltis huegelii huegelii</i>	✓	✓	1/2	<i>Biza minor</i>	✓	—	1/2				
<i>Hibbertia myrtilloides</i>	✓	✓	1/2	HERBS				<i>Boraginaceae</i>	✓	✓	1/2
<i>Gompholobium marginatum</i>	✓	✓	1/2	<i>Bartsia tixago</i>	✓	✓	1/2	<i>Romulea rosea</i>	✓	✓	1/2
<i>Grevillea bipinnatifida</i>	✓	✓	1/2	<i>Trachymene caerulea</i>	✓	✓	1/2	<i>Stylidium bulbiferum</i>	✓	✓	1/2
<i>Myrtalea ledifolium</i> seedlings	✓	✓	1/2	<i>Hypochaeris glabra</i>	✓	✓	1/2	<i>Daucus glochidiatum</i>	✓	✓	1/2
H				<i>Lithum trigynum</i>	✓	✓	1/2				
				<i>Senecio diascioides</i>	✓	✓	1/2				
				<i>Dioscorea hastifolia</i>	✓	✓	1/2	SEDGES			
				<i>Caesia micrantha</i>	✓	✓	1/2	<i>Pesmodadus aspera</i>	✓	✓	1/2
				<i>Haemodorum discolor</i>	✓	✓	1/2				
				<i>Thysanotus dichotomus</i>	✓	✓	1/2				
				<i>Thysanotus marginatus</i>	✓	✓	1/2				
				<i>Ursinia anthephoides</i>	✓	✓	1/2				
				<i>ADS: Dampiera linearis</i>	✓	✓	1/2				

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP06
 Date 24.10.96

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
E wandoo (seedling)	✓			Trimalium? seedlings -				Adiantum arvensis	✓		///
SHRUBS				stipules on leaves				Nahianbergia? - yellow fl.	✓		✓
Metaleuca? fulgens/radula	✓		///	(Aveilla? seedling?)			///	Unid dicot weed	✓		///
Daviesia? divaricans	✓		///	Borja sp. acocephala	✓		///	superficially like hypochaeris glabra	✓		///
Diplonectis sp.	✓		///					Chidwee - Gaulis			///
Prunella sp.	✓							Romulea - rosea	✓		///
MALLEES				Kennedia prostrata?? (nr)	✓		///	Haemodorum sp.	✓		///
Crevillea endlicheriana	✓		///					Medicago / Trifolium	✓		///
Darwinia citriodora	✓		///	GRASSES							
Gompholobium marginatum	✓		///	Briza maxima	✓		///	Hypochaeris glabra	✓		///
Lupinus? pumila (Shrub)	✓		///	Avena fatua	✓		///	Aira caryophylla	✓		///
SHRUBS → lanceolata				Briza minor	✓		///	Thysanotus patersonii	✓		///
Acacia pulchella (seedling) & adult	✓		///	Romulea rosea	✓		///	SEDGES			
Crevillea bipinnatifida	✓		///	Neurachne alopecuroides // Amphipogon?	✓		///				
(adult)				Micradis sp. (ad in				Unid legume - 5 pinnate / leaflets	✓		///
				Mondani	✓		///	on pinnate leaf	✓		///
				Neurachne alopecuroides				LOTUS			
				Neurachne alopecuroides / Amphipogon	✓		///	Loxocarya flexuosa	✓		///
				HERBS							
				Drosera nemjosi? pallida?				Scroph - small purple corolla	✓		///
				Cheilanthes lausvirensifolia	✓		///				
				Dioscorea hastifolia	✓		///	Aira caryophyllae	✓		///
				Stylidium bulbiferum	✓		///				
				Solan - large pink-white corolla	✓		///	Small, climbing Drosera			///
				Caesia Microcarpa	✓		///	Vulpia sp	✓		///
				Xanthosia candida	✓		///	hepicasperma - sword leaf	✓		///
				Thysanotus dichotomus	✓		///	Unid small asteraceae			
				Dorcus → Dorcus minus glaberrimus	✓		///	Trachymene sp			
Ursinia anthemoides				Vesinia anthemoides	✓		///	Trachymene sp			
				Hemionia pungens → (not Horia)				→ large / Dorcus			
				Sonchus oleraceus - like weed							
				Danthonia sp							

Trachymene coerulea → prob the other large 'Lantana' species

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP06

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

No windos trees in plot

Date 13-12-96

Rescore

[illegible]

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP ^{→ creekline track} SITE NUMBER SERP07


DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location Along Creekline Access Firebreak → to Rangers house

Geographic Location Latitude 32 22 50.3 S Longitude 116 00 24.9 E Altitude 160m (map)
GPS ✓ Accuracy PDOP = 4.14
Map Sheet Jannahdale

Photograph Name _____ Number #5 Roll # 4

Topographic Position (Circle the correct response)
Upland _____
Upper-slope _____
Midslope (circled)
Lower slope (circled)
Valley (circled)
Washline _____
Valley east of 1st Ridge

2. SITE DATA Circle the correct response

Slope flat _____ gentle (circled) steep _____ Aspect N NE E SE S SW W NW

Surface Soil clayey loam Depth to Rock 10 cm → 100 cm Colour orange → dk brown

Exposed Rock type Granite % surface cover 1% size range 2cm → 20cm

Litter 70% % surface cover _____ Depth (cm) 5-15cm

Bare Ground 1 % surface cover _____ (thick leaf litter cover)

Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

With loose rock in the soil profile!

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class						2-10%	10-30%	30-70%	over 70%
1° Tall open wandoo woodland (10-30%)									
TREES						MALLEES			
over 15m		5-15m		under 5m		over 8m		under 8m	
LIFE FORM						30m			
COVER CLASS (%)	(class 10-30) 30-40%	2-10%				10m			
DOMINANT SPECIES	E. wandoo	E. wandoo							
SHRUBS						SHRUBS			
over 2m		2m - 1.5m		1.5 - 1m		1 - 0.5m		under 0.5m	
LIFE FORM						Hypo. angust. E. Banksia. dissocarp. 2m 1m			
COVER CLASS (%)						30-70%			
DOMINANT SPECIES						Hypo. F. E. Can. preissii Hibb. hypoleioides Phyl. cal.			
GRASSES		HERBS		SEDGES		over 0.5m		under 0.5m	
LIFE FORM									1m
COVER CLASS (%)		10-30%							
DOMINANT SPECIES		Stypandria glauca Cheilanthes aust.							

4. VEGETATION CONDITION

Comments x Weed invasion minor.
 ≈ 4/5. Long interfire period. Rotting stumps & signs
 Tall wandoo. Adjacent slope (downslope) has been
 recently burnt & extensive weed invasion has followed
 subsequent to fire

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIFIC PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP87
Date 24.10.96
13.12.96 (R)

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check look-leaf at SERP86

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
Euc. wandoo	✓	—	1/2	Thysanotus sp. <u>thysanotus</u>	✓	—	✓	Arturosanthus huxus	✓	✓	1/2
ADS Allocas. luegeliana	✓	—	1/2	Caladenia sp.	✓	—	✓	Lavenhookia stipitata	✓	✓	1/2
Xanthorrhoea preissii	✓	—	1/2	Dampiera linearis	✓	✓	1/2	Comesperma olympos	✓	✓	1/2
Macroglossia riedlii	✓	—	1/2	Pavementella viscosa	✓	✓	✓	Oxalis permaria	✓	✓	1/2
Hakea dissocarpa	✓	—	1/2	Lomandra integrum	✓	✓	✓	Romulea rosea	✓	—	1/2
Hypocalymma angustifolium	✓	—	1/2	Sonchus oleraceus	✓	✓	1/2	Burchardia congesta	✓	✓	1/2
Gompholobium marginatum	✓	✓	1/2	Stylidium bulbiferum (indd)	✓	✓	1/2	Agrostocrinum sabum	✓	✓	1/2
Leucopogon capitellatus	✓	✓	1/2	Haemodorum discolor	✓	✓	✓	Tricoryne elatior	✓	✓	1/2
Hibbertia lupulicoides	✓	✓	1/2	GRASSES				Dianella revoluta	✓	✓	1/2
Leucopogon propinquus	✓	✓	1/2	Stipa pycnostachya	✓	✓	1/2	Ursinia anthoroides	✓	✓	1/2
Grevillea pilulifera	✓	✓	1/2	Aira canyophylla	✓	✓	1/2	Acanthocarpus preissii	✓	✓	1/2
Hibbertia communata	✓	✓	1/2	Microlaena stipoides	✓	✓	1/2	Xanthosia candida	✓	✓	1/2
Daviesia harrida	✓	✓	1/2	Danthoria sp	✓	✓	1/2	Ptilotis manglesii	✓	✓	1/2
Dyandra lindleyana lindleyana	✓	✓	1/2	Poa drummondiana	✓	✓	1/2	Lomandra hermaphrodita	✓	✓	1/2
Hibbertia spicata	✓	✓	1/2	Dichelanthus crinita	✓	✓	1/2	Thysanotus manglesianus?	✓	✓	1/2
Phyllanthus calycinus	✓	✓	1/2	Briza maxima	✓	✓	1/2	(T. dichotomus)			
Thomasia foliosa	✓	✓	1/2	Neurachne alopecuroides	✓	✓	1/2	Lomandra caespitosa	✓	✓	1/2
Olearia paucidentata	✓	✓	1/2	Briza minor	✓	—	1/2	Lomandra ? odora	✓	✓	1/2
Kunzea micrantha	✓	✓	1/2	HERBS				Pavementella latifolia	✓	✓	1/2
Acacia pulchella	✓	—	1/2	Centaurium erythraea	✓	✓	1/2	T. manglesianus	✓	✓	1/2
Petrophile bulbosa	✓	—	1/2	Conostylis setosa	✓	✓	1/2	Haemodorum simplex	✓	✓	1/2
Grevillea bipinnatifida	✓	—	1/2	Barbosa tixago	✓	✓	1/2	Sowea baea laxiflora	✓	✓	1/2
Xanthorrhoea gracilis	✓	—	1/2	Wahlenbergia preissii	✓	✓	1/2	Styandra glauca	✓	✓	1/2
→ Astroloma pallidum	✓	—	1/2	Arceuthobium glandulifera	✓	✓	1/2	Dioscorea hastifolia	✓	✓	1/2
↖ Drosera sp. ? menziesii	✓	—	1/2	Dampiera pulata	✓	✓	1/2	SEDGES			
Lewinhookia pusilla	✓	—	1/2	Stachysia morongana	✓	✓	1/2	Schoenus unispicatus	✓	✓	1/2
Arthropodium capillipes	✓	—	1/2	Anagallis arvensis	✓	✓	1/2	Desmodadus aspera	✓	✓	1/2
Cheilanthes austrotenuifolia	✓	✓	1/2	Opereularia vaginatum	✓	✓	1/2	Schoenus nanus	✓	✓	1/2
Poaanthera microphylla	✓	✓	1/2	Caesia micrantha	✓	✓	1/2	Tetradlea octandra	✓	—	1/2
Centrolepis drummondiana	✓	✓	1/2	Stylidium hispidum	✓	✓	1/2				
Hydrocotyle (pilifera) / callicarpa	✓	✓	1/2	Senecio diaschides	✓	✓	1/2				
Monadenia bracteosa	✓	✓	1/2	Eragrostium pinnatifidum	✓	✓	1/2				

Astragalus ciliatum ✓

R.

Drosera stolonifera ✓



ADS: Linum trigynum ✓

ADS Lomandra spartea. (Rescore) ✓

ADS: Conostylis aculeata ✓

- Examples of the sand environments that dominate the Ereman (Australia's red centre) are well represented in the study area, as are a variety of the semi-arid woodlands, sand heaths and coastal habitats of the South-west.
- Rich *Lerista* assemblages were associated with the *Eucalypt* and/or *Acacia* litter-patches found on virtually all non-saline quadrats in the study area, even quadrats with only a thin sandy soil A-horison (3.8 ± 1.4 , 56). Seven species were recorded on some quadrats (BB4, CU3, PE4 & ZU3). **Kendrick (19??)** concluded that there has been a significant radiation of *Lerista* taxa in the study area.
- The arid-adapted genera *Ctenophorus*, *Diplodactylus*, *Ctenotus* and *Lerista* are very diverse (10, 11, 18 and 23 species respectively) with high levels of syntopy among *Ctenotus* and *Lerista* species, particularly on the 14 quadrats which represented the red sand-hummock grass environments that dominate Australia's arid zone: 7.8 ± 1.6 (14) versus 4.2 ± 1.9 (49) *Ctenotus* + *Lerista* species per quadrat. In equivalent sand-hummock grass habitats of the Great Victoria Desert, elsewhere in temperate Western Australia, Pianka (1969b) noted up to seven *Ctenotus* species in "ecologic sympatry".

Heatwole and Taylor (1987, p. 215 *et seq.*) reviewed the literature on factors that are known to affect richness in reptile assemblages. A range of factors, including rainfall, amount of sunshine, structural complexity in vegetation and productivity have been shown to have a positive correlation with species richness. In arid regions of Australia, richness has also been explained in terms of environmental heterogeneity, the presence of hummock grass, occupation by reptiles of niches elsewhere occupied by other animal taxa, and low nutrients (poor soils) limiting homeotherms (see Heatwole and Taylor 1987 p. 220, **Morton 19??, Pianka 19??, but also see Rosensweig 199??**).

The 14 quadrats representing these low nutrient, red sand with hummock-grass environments in the Irwin-Carnarvon study area had the richest assemblages we encountered. They yielded an average of 21.5 ± 4.8 (SD) herpetofaunal species (BB4, PE2,5, CU2-5, MR2-5, NE1, KE1-2; Table QR). Rich reptile assemblages have previously been reported from such hummock grasslands on red sand in the Great Victoria Desert (Pianka 1969, Armstrong 1992). For instance, Pianka (1969a) reported 39 species of lizard co-existing in sympatry in a square kilometre of sandridge habitat, and at several sandplain sites he reported up to 30 species. Elsewhere in Central Australia, Fyfe (1985) found that hummock grasslands on red sand surfaces supported a greater number of reptile species than did the Mulga shrublands that occurred on finer textured soils in the same area.

Patterns in the Species Composition of Assemblages

Scale is important in determining which environmental attributes emerge as significant correlates between compositional patterns and environmental attributes (Whitmore 1984, **Also check Dale's paper in A. J. Ecol.**). Regional studies have usually linked compositional patterns to climatic and/or geological categories (Ashton 1976, McKenzie *et al.* 1987b, 1991a, 1992, 1994, McKenzie & Rolfe 1995). In more geographically restricted studies, topographic setting and lithology become important (Whitmore 1984).

in all sites check previous - was on adj - plot -

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP 7

Record on Sheet

Date 24.10.96

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Acanthocarpus preissii

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
E. wandoo	✓			Xanthosia candida	✓		III	Leucnoodia pusilla	✓		III
Allocas. huegeliana (ADS)				Dampiera linearis	✓			Thysanotus? patersonii			III
SHRUBS				Unid small shrub/seedling				Orosia manneissii? (glabrous)			
Daviesia? divergens	✓		III	Trymal ecidifolium				Scaevola sp →	✓		III
Phyllanthus calycinus	✓		III	(cf. as Grevillea or Grevillea?)	III			Black based Schenus	✓		III
Placida pulchella	✓			Walenbergia - hairy leaves			III	Large white Scroph	✓		III
MALLEES				H. spicata? check id			III	Senecio sp	✓		III
Xan. preissii	✓			Small annual Schenus -/Isolopis (small)	III			Lomandra caespitosa	✓		III
Hibbertia hypericoides	✓			GRASSES				Aiza canyphyllae	✓		III
Macrozamia veidtii	✓			Microzamia? / Pra			III	Ursinia antheroides	✓		III
Styphandra glauca	✓			NB. Long-leaved tall grass	✓		III	Haemodorum sp. strapleaf	✓		III
SHRUBS				prob is this sp.				Stachysia sp. huna	✓		III
Hakea lissocarpa	✓			Briza maxima	✓		III	Unid Small rhombic leaf seedling -? Scaevola?			
Hypocalymma argenteifolium	✓		III	Neuradachne alopecuroides	✓		III	SEDGES			
Dry. nivea	✓		III	Stipa elegantissima	✓		III	Lepidosperma large terete smooth			III
Monarda foliosa	✓		III	Briza minor	✓			Loxocarya flexuosa	✓		III
Petrophile biloba	✓			Caesia micrantha	✓		III	Glossostachya scabra Agrostocrinum			III
Macrozamia veidtii	✓			Drosera stolonifera				Diastocum elastifolia Dioscorea hastifolia			III
Hibb. commutata	✓		III	HERBS				Microtis - Microtis - small Asteraceae			III
Kunzea? sp.	✓		III	Stylidium bulbiferum	✓		III	Tetrahrea octandra - tetvaria	✓		III
Tetrahrea capillipes? (rough base) (epidaspema?)				Sowerbaea laxifolia	✓			Lepidosperma - glabrous, terete (Hawii)			III
Phylotus manneissii	✓		III	Styphandra glauca	✓		III	Short Lomandra fine leaves	✓		III
Opercularia echinocarpa	✓		III	Stylidium piliferum	✓		III	Stylidium piliferum			
Ascoloma sp	✓		III	Cheilanthes austrotenuifolia	✓		III	Hydrocotyle sp. (single plant)			
Chorizandra - trilobate leaf -				Orthocentrus sp.	✓		III	Sonchus olerasis			
inflated pods? Hovea?				Conostylis setosa	✓		III	Romulea rosea	✓		
Acanthocarpus preissii (IV)				Arthropodium capillipes	✓			Caledonia sp - single leaf - Spider orchid			III
Styphandra (VI)				Oxalis? corniculata	✓		III	Arthropodium capillipes	✓		III
Grevillea bipinnatifida	✓		III	Trichoryne elaeor	✓		III	Thysanotus - terete leaf (at serax)			III
Grevillea pilulifera - check id.	✓		III	Hairy / ligulate grass	✓		III	Anagallis arvensis	✓		III
Dianella revoluta	✓		III	Burchardia banded congesta	✓		III	terete leaf - Anthrocaceae?			III
Anthrocaceae Danthonia sp	✓		III	Eryngium peritotidum	✓		III	Centrolepis drummondii			

Unid Euphorb
very narrow leaf Haemodorum
Small annual grass
Gompholobium marginatum
Wiry leaved Haemodorum - collected ad inflor. as adj? - check match.
Xanthorrhoea gracilis
Olearia paucidentata
Haemodoraaceae

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP07

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Date 13-12-96

* sighted
Brush-tail
Wallaby

Rescue

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
								Opp-leaved pine fws herb	///		
								<i>Astroloma pallidum</i>			
								Hairy-leaved grass			
								<i>Briza minor</i>			
								<i>Astroloma ciliatum</i>			
MALLEES								Adj. <i>Leandra spartea</i>			
								<i>Ptilothus mangleri</i>		///	
				GRASSES				<i>Dianella revoluta</i>		///	
								<i>Levenhookia pailla</i>		///	
								<i>Burchardia congesta</i>		///	
SHRUBS								<i>Stackhousia</i> sp INF		///	
								SEDGES			
								<i>Pericaria hispida</i>		///	
								<i>bedina?</i>			
								<i>Tetrachena laevis</i>		///	
				HERBS				ADJ - <i>Perostylis aculeata</i>		///	
								ADJ - <i>Linum - yellow fws</i>			

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP SITE NUMBER SERP08


DATE TRIP 25-10-96 RECORDERS AM DL

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location Gravel 4WD track following creekline in

Geographic Location Latitude 32° 22' 25.0" S Longitude 116° 00' 22.4" E Altitude 60m
 GPS Accuracy - From Map
 Map Sheet no fix

Photograph Name _____ Number #6 Roll #4

Topographic Position (Circle the correct response)
 Upland _____
 Upper slope _____
 Midslope _____
 Lower slope _____
 Valley Washline - Creekline in Valley

2. SITE DATA Circle the correct response

Slope flat _____ gentle steep 2 Aspect N NE 52° 52° 2 E SE S SW W NW

Surface Soil rich loam Depth to Rock 0-30cm Colour dk Brown

Exposed Rock type granite % surface cover 2 size range 20-30cm

Litter 280 % surface cover _____ Depth (cm) 10cm

Bare Ground 0 % surface cover _____



Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

Creek at base of steep granite hillsides
 = granite influence
 & granite boulders in creekbed

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class		2-10%	10-30%	30-70%	over 70%
TREES					
	over 15m	5-15m	under 5m	MALLEES over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		2-10%			
DOMINANT SPECIES		<i>E. catenophylla</i>			
SHRUBS					
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	SHRUBS under 0.5m
LIFE FORM					
COVER CLASS (%)	10-30 10-20%	> 70%	10-20%		2-10%
DOMINANT SPECIES	<i>H. trifurcata</i>	<i>Darwinia citriodora</i> <i>Myrsine floribunda</i> <i>Podanea</i> sp.	<i>Thysanotus spatulatus</i>		<i>Acrothosia prostrata</i>
GRASSES					
	HERBS	SEDGES	over 0.5m	under 0.5m	
LIFE FORM					
COVER CLASS (%)		20			
DOMINANT SPECIES		<i>Ten</i>			

4. VEGETATION CONDITION

Comments ^{fire} History of creekline unknown. Tall stands of *Halea pinnatifida*
4 515 L Excellent condition. However, *Leadsoria* invading
from creekline in adjacent areas & vegetation
fringed by road on *Antennaria*. Very thick vegetation!

✓ Lots of *H. nigrivata*

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP08
Date 25.12.96
13.12.96(R)

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Eucalyptus calophylla</i>	✓	—	✓								
<i>Allocasuarina laevis luegeliana</i>	✓	—	✓								
<i>Macrozamia riedlii</i>	✓	—	✓								
<i>Melaleuca radula</i>	✓	✓	✓								
<i>Hakea trifurcata</i>	✓	✓	✓								
<i>Melaleuca aff. scabra</i>	✓	✓	✓								
<i>Calytrix acutifolia</i>	✓	✓	✓								
<i>Tristaniopsis floribundum</i> flor.	✓	✓	✓	GRASSES							
<i>Thomsonia foliosa</i>	✓	✓	✓	<i>Neuradme alopecuroides</i>	✓	✓	✓				
<i>Hibbertia spicata</i> spicata	✓	✓	✓	<i>Brija maxima</i>	✓	✓	✓				
<i>Xanthorrhoea preissii</i> (no material)	✓	✓	✓	<i>Tetrarrhena laevis</i>	✓	✓	R				
<i>Labeclia lanceolata</i>	✓	✓	✓	<i>Amphipogon</i> ? sp. <i>stichus</i>	✓	✓	R				
<i>Baeckea carphorosmae</i>	✓	✓	✓	<i>Stipa</i> ? <i>pycnostachya</i>	✓	✓	✓				
<i>Dodonaea certocarpa</i>	✓	✓	✓	<i>Microseris stipoides</i>	✓	✓	✓				
<i>Darwinia citriodora</i>	✓	✓	✓								
<i>Hypocalymma angustifolium</i>	✓	✓	✓								
<i>Phyllanthus calycinus</i>	✓	✓	✓	HERBS							
<i>Acacia pulchella pulchella</i>	✓	✓	✓	<i>Anagallis arvensis</i>	✓	✓	✓				
<i>Daviesia horrida</i>	✓	✓	✓	<i>Araucario carpus preissii</i>	✓	✓	✓				
<i>Dryandra lindleyana</i> lindleyana	✓	✓	✓	<i>Tricornis elatior</i>	✓	✓	✓				
<i>Conostylis</i>	✓	✓	✓	<i>Conostylis aculeata preissii</i>	✓	✓	✓				
<i>Xanthorrhoea preissii</i> (no material)	✓	✓	✓	<i>Burchardia congesta</i>	✓	✓	✓				
<i>Ashodora pallidum</i>	✓	✓	R	<i>Thysanotus dichotomus</i>	✓	✓	✓	SEDGES			
<i>Acaenocarpus preissii</i>	✓	✓	✓	<i>Haemodorum</i> sp. <i>indet</i>	✓	✓	✓	<i>Desmodium aspera</i>	✓	✓	✓
<i>Xanth. gracilis</i> ?? (no material)	✓	✓	✓	<i>Chamaecilla corymbosa</i>	✓	✓	✓	<i>Tetraria capillaris</i> ?	✓	✓	✓
				<i>Caesia micrantha</i> (sterile)	✓	✓	✓				
				<i>Lomandra micrantha</i>	✓	✓	✓				
				<i>Lomandra nigricans</i>	✓	✓	R				
				<i>Cheilanthes austrotenuifolia</i>	✓	✓	✓				
				<i>Thysanotus mangelicus</i>	✓	✓	✓				

Although climatic attributes provided the best fits, a wide array of climatic and substrate gradients were strongly correlated with the compositional patterns revealed by our numerical analysis of the total data-base. **Many of these environmental gradients were not inter-correlated**, and several distinct stepwise structures dominated both the quadrat and the species classification dendrograms, so we concluded that the component species were responding to different environmental gradients and/or to the same gradients differently.

On the reasoning that species which usually co-occur are more likely to be responding to the same environmental gradient/s than are species with different patterns of co-occurrence, we sought to untangle the biogeographical processes by treating the four species assemblages derived from the overall analysis as independent data-sets, and analysing them separately (cf. McKenzie *et al.* 1989, 1991a, McKenzie & Belbin 1991). Strong relationships between herpetofaunal patterns and soil-geomorphic attributes emerged for species assemblage_1 (soil pH), assemblage_3 (soil texture) and assemblage_4 (over-bank stream flow). Assemblage_2 conformed with a gradient in winter rainfall.

These analyses of the individual assemblages revealed that patterns in species composition of the herpetofaunal assemblages were being influenced by environmental processes operating at two geographical scales:

- A biogeographical scale related to the differences between the Eremean and South-western faunas, and corresponding to the study area's arid-to-mesic and tropical-to-temperate climatic gradients.
- An ecological scale related to geomorphic and soil attributes.

Localised patterns of allopatry among closely related species belonging to the more diverse genera was an additional source of variation in assemblages_2 and _3. Our ecologically re-defined taxa virtually eliminated this variation without losing ecological discrimination, indicating that the reserve system will need to sample the geographical extent of the winter-rainfall and soil texture gradients, if evolutionary processes are to be protected.

The classification structures derived from a matrix of frog, gecko, dragon and skink trap-records (Figure TRAP.2wa) were virtually identical to the structures derived from the total data-set (Figure DENT). Snakes, pygopids and goannas were poorly sampled on our quadrats, perhaps because of their mobility and low densities. Pragmatically, these three taxa could have been ignored, and the labour-intensive hand-foraging methods deleted, without sacrificing ecological or biogeographical discrimination or changing the outcomes of the study.

REFERENCES

- Armstrong, G.P. (1992). Reptiles. pp 163-179. In: A Biological Survey of the Yellabinnia Region, South Australia, in October 1987. (Ed by P.G. Copley & C.M. Kemper). National Parks and Wildlife Service. South Australian Department of Environment and Land Management.

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

SITE No	SERP 08
Date	25-10-96

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>E. calophylla</i>	✓			<i>Hibbertia spicata</i>				<i>Haemodorum</i> sp.	✓		
				<i>Melaleuca</i> aff. <i>scabra</i>	✓			<i>Chenopodium</i> sp.			
SHRUBS				<i>Melaleuca radula</i> // <i>fulgens</i>	✓			<i>Anagallis arvensis</i> (seedling)			
<i>Allocasuarina kuoriana</i>	✓			<i>Tricoryne elatior</i>	✓						
<i>Dodonaea</i> sp. <i>ceratocarpa</i>				<i>Xanthorrhoea gracilis</i>	✓			<i>Chenopodium</i> sp. <i>corymbosa</i>			
<i>Hakea trifurcata</i>	✓			<i>Hypocalymma angustifolium</i>							
MALLEES				small, unid seedling <i>Scaevola</i> ?							
<i>Darwinia citriodora</i>	✓			? <i>L. Strobil.</i> <i>Diplocladia</i> ?							
<i>Trimalium robustum</i>	✓			GRASSES							
<i>Druandrea nuda</i>	✓			<i>Chloris-lamina</i> grass							
<i>Halimolobos</i> sp. (tall shrub)	✓			<i>Halimolobos</i> grass							
SHRUBS				<i>Amphipogon</i> // <i>Neurachne</i>	✓			SEDGES			
<i>Calytrix acutifolia</i>	✓							<i>Lepidosperma</i> sp. - small - edged	✓		
<i>Acanthocarpus preissii</i>	✓			<i>Stipa</i> sp. - v. hairy stem & lamina	✓			<i>Lepidosperma terete</i> , <i>scabrous</i>			
<i>Thamnia</i> sp. <i>foliosa</i>	✓			<i>Briza maxima</i>	✓			<i>Loxocarpus flexuosus</i>	✓		
<i>Daveya</i> ? <i>divergens</i>	✓							<i>Lomandra</i> ? <i>preissii</i>	✓		
(1° broadleaved form)								<i>Tetraria capillaris</i> ?			
<i>Macrozamia verdii</i>	✓			HERBS							
<i>Phyllanthus calycinus</i> (NE)	✓			<i>Thysanotus</i> sp? <i>spatus</i> ??							
<i>Baeoba camphorosmae</i>	✓			<i>Conostylis acutata</i>	✓						
<i>Acacia pulchella</i>	✓			<i>Burchardia</i> or <i>Caesia</i>							
				<i>L. uncinata</i> - white bases							
<i>Stipa elegantissima</i>	✓			& brown pappi scale - <i>Andropogon</i>							
→ prob. some <i>Andropogon</i>				<i>Cheilanthes tenuifolia</i>	✓						
other species				<i>Thysanotus patersonii</i>	✓						

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP08

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 Identification check

Date 13.12.96

[illegible]

check in specimens if
sparse in fl or tall grass.

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP. Cheltenham Rd Access **SITE NUMBER** SERP09 (Reocore)


DATE TRIP 29.10.96 **RECORDERS** AM

DATE TRIP _____ **RECORDERS** _____

DATE TRIP _____ **RECORDERS** _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location 36.3 - Cheltenham Rd

Geographic Location **Latitude** 32 22 **S Longitude** 115 59 43.8 **E Altitude** 60m **GPS Accuracy** PDOP=3.3 6SAT

Photograph **Name** _____ **Number** #8 + #9 Rd #4

Topographic Position **Upland** _____ **Lower slope** → Flats at base of hills **Upper slope** _____ **Valley** _____ **Midslope** _____ **Washline** _____

2. SITE DATA Circle the correct response

Slope flat **gentle** **steep** **Aspect** N **NE** **E** **SE** **S** **SW** **W** **NW**

Surface Soil Sandy loam **Depth to Rock** >1m **Colour** Dark Brown to grey

Exposed Rock **type** _____ **% surface cover** _____ **size range** _____


Litter 30 **% surface cover** _____ **Depth (cm)** 2-3cm

Bare Ground 2 **% surface cover** _____

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class					
	2-10%	10-30%	30-70%	over 70%	
TREES					
	over 15m	5-15m	under 5m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		2-10%			
DOMINANT SPECIES		E. lane-poolii			
SHRUBS					
	over 2m	2m - 1.5m	1.5 - 1m	1-0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)				10-30%	10-30%
DOMINANT SPECIES				Stylidium latifolia Mel. aff. scabra	Dasyopogon breviflorus Patersonia occidentalis
GRASSES		HERBS	SEDGES	over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)					30-70% -
DOMINANT SPECIES					Unid sedges - small Restio - Large Gahnia - Mel. Mesorelaena foliagosa

4. VEGETATION CONDITION

Comments

S/S - Some weeds. Benched sandplain at base of foothills. Rescore. E. lane-poolii x Poorly defined winter wet depression. has Sedge community.

1.0

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No **SERP09**

Record on Sheet

Date **29.12.96**

13.12.96(R)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

-Rescore Bronwyn SERP09

TREES				SHRUBS (cont.)				HERBS (cont.)			
No	FI	ID		No	FI	ID		No	FI	ID	
<i>Eucalyptus laevis</i>	✓	✓	✓	<i>Caesia occidentalis</i>	✓	✓	✓	<i>Drosera rotundifolia</i>	✓	✓	✓
<i>Eucalyptus calophylla</i>	✓	✓	✓	<i>Lomandra caespitosa</i>	✓	✓	✓	<i>Lolus angustissimus</i>	✓	✓	✓
				<i>Triconia elatior</i> (Not on sheet)				<i>Centropogon ovatus</i>	✓	✓	✓
				<i>Leucanthodia pusilla</i>	✓	✓	✓	<i>Hemiliosciadium hemiliosciadium</i>	✓	✓	✓
<i>Xanthorrhoea preissii</i>	✓	✓	✓	<i>Stylidium hamiltonianum</i>	✓	✓	✓	<i>Dampiera alata</i>	✓	✓	✓
<i>Kingia australis</i>	✓	✓	✓	<i>B. brya scirpoides</i>				<i>Stylidium calcitrans</i>	✓	✓	✓
<i>Kunzea sp. micrantha</i>	✓	✓	✓	<i>Previous record B. scirpoides</i>				<i>Trachymene pilosa</i>	✓	✓	✓
<i>Lechenostictis biloba</i>	✓	✓	✓	<i>Heb det R sphaeroph</i>				<i>Pterochaeta paniculata</i>	✓	✓	✓
<i>Nannia capitatum</i>	✓	✓	✓	GRASSES				<i>Hydrocotyle calycarpa</i>	✓	✓	✓
<i>Acacia laterifolia</i>	✓	✓	✓	<i>Neurachne abepecuridea</i>	✓	✓	✓	<i>Brya sphaerocephala</i>	✓	✓	✓
<i>Jacksonia vestrioides</i>	✓	✓	✓	<i>Brya maxima</i>	✓	✓	✓	<i>Sonchella laxiflora</i>	✓	✓	✓
<i>Possiaea ariocarpa</i>	✓	✓	✓	<i>Brya minor</i>	✓	✓	✓	<i>Ranunculus rosea</i>	✓	✓	✓
<i>Baeckea camphorosma</i>	✓	✓	✓	<i>Danthonia sp.</i>				<i>Lomandra hemisphaerica</i>	✓	✓	✓
<i>Hakea ceratophylla</i>	✓	✓	✓	<i>Vulpia bromoides</i>	✓	✓	✓	<i>Dasypogon bromeliifolius</i>	✓	✓	✓
<i>Melaleuca aff. scabra</i>	✓	✓	✓	<i>Danthonia setacea</i>	✓	✓	✓	<i>Drosera sp. n. n. n.</i>	✓	✓	✓
<i>Stirlingia latifolia</i>	✓	✓	✓					<i>Stirlingia latifolia</i>			
<i>Dyandra lindleyi</i>	✓	✓	✓					<i>Lomandra sericea</i>			
<i>Dasypogon bromelioides</i>	✓	✓	✓					<i>Thysanotus manglesianus</i>	✓	✓	✓
ADS: <i>Dodonaea ceratocarpa</i>	✓	✓	✓	HERBS				<i>Casuarina glauca</i>	✓	✓	✓
(W) <i>Hypocalymma angustifolium</i>	✓	✓	✓	<i>Hypochaeris glabra</i>	✓	✓	✓	<i>Haemodorum laxum</i>	✓	✓	✓
<i>Euc. marginata</i>	✓	✓	✓	<i>Ursinia anthemoides</i>	✓	✓	✓	<i>Lomandra brittanii</i>	✓	✓	✓
				<i>Xanthoxia huegelii</i>	✓	✓	✓	<i>Lomandra sericea</i>	✓	✓	✓
				<i>Stylidium bulbiferum</i>	✓	✓	✓	<i>Brachyotum ciliaris</i>	✓	✓	✓
				<i>Opercularia vaginata</i>	✓	✓	✓	SEDGES			
				<i>Dampiera linearis</i>	✓	✓	✓	<i>Cyathochaeta avenacea</i>	✓	✓	✓
				<i>Podolepis gracilis</i>	✓	✓	✓	<i>Hypolaena exsulca</i>	✓	✓	✓
				<i>Siloxerus humifusus</i>	✓	✓	✓	<i>Mesomelaena tetragona</i>	✓	✓	✓
				<i>(W) Patersonia occidentalis</i>				<i>Loxocarya fasciculata</i>	✓	✓	✓
				<i>Anigozanthus manglesii</i>	✓	✓	✓	<i>Schoenus sculptus</i>	✓	✓	✓
				<i>Philyrella pygmaea</i>	✓	✓	✓	<i>Schoenus nanus</i>	✓	✓	✓
				<i>Casuarina penifloris</i>	✓	✓	✓	<i>Isotria marginata</i>	✓	✓	✓
				<i>Chaenactis lanuginosa</i>	✓	✓	✓	<i>Tetraria aelandra</i>	✓	✓	✓

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE Label each plant with plants number, site code, date and plant's name or working name if required

SITE No	SERP09	Record on Sheet	• Column 1	plant name
---------	--------	-----------------	------------	------------

Date	29.10.96	• Column 2	plant number
------	----------	------------	--------------

SITE No	SERP09	Record on Sheet	• Column 1	plant name
---------	--------	-----------------	------------	------------

- | | |
|----------------------|---|
| Date <u>29.10.96</u> | <ul style="list-style-type: none"> • Column 2 plant number • Column 3 flowering time- TICK if species flowering • Column 4 identification check |
|----------------------|---|

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
E. lane-paulii	✓		✓	Borassia ? aridophylla			✓	Haemodorum sp. (M. p. p.)			✓
E. calophylla (Chrysophyllum?)	✓		✓	Labichea punctata	✓			Broad			✓
SHRUBS				Sonchella laxiflora	✓		✓	Phyllodrella pygmaea			✓
Xanthorrhoea sp. peisii ?	✓			Aracia drummondii seedling	✓		✓	Cassutha sp.	✓		✓
Kingia australis	✓		✓	drdc Nemcia	✓		✓	Unid small white Asteraceae			✓
Melaleuca scabra	✓		✓	Homna homna	✓		✓	Bony sp. (Asteraceae)	✓		✓
MALLEES				Hydrocotyle			✓	Podotheca sp.	✓		✓
Skullingia latifolia	✓		✓	Levenhockia			✓	Waltia paniculatum	✓		✓
Baeckea campylocladia	✓		✓	GRASSES				Tiny Stylidium pusillum			✓
Pterocarya occidentalis	✓		✓	Briza maxima	✓		✓	Trachymene pilosa			✓
Haem				Neurochne alpestrum	✓		✓	Annual schaefferia / Isopis			✓
SHRUBS				Unid grass - Dactyloctenium	✓		✓	Centropogon aristata			✓
Unid small Danthonia galls								SEDGES			
Dryandra nivea	✓		✓	Chromola corymbosa	✓		✓	Mesembryanthemum tetragona	✓		✓
→ Kunzea sp.	✓		✓	Small annual grass				Loxocarya fasciculata	✓		✓
				- Flowers on peduncles				Unid Restio -	✓		✓
Lechenaultia biloba	✓		✓	Drosera ? (Myrtaceae) (red)				seicea			
Opuntia echinocarpa	✓		✓	HERBS				Leonardra ? (Myrtaceae)	✓		✓
Dactyloctenium aegyptium	✓		✓	Caesalpinia sp. (Myrtaceae) (large)	✓		✓	Unid sedge with fuzzy inflorescence			✓
Xanthosia aegyptium	✓		✓	Drosera stolonifera	✓		✓	galls			
6-leaflet tiny legume pea			✓	Pampiera linearis	✓		✓	Gahnia // Tetradlea orbicula	✓		✓
Anigozanthus nargens	✓		✓	Romulea rosea	✓		✓	sp. Loxocarya			
ADS (Podonaea)				Stylidium calceatum			✓	Curled-leaf Leonardra	✓		✓
Tricoryne elatior	✓		✓	Musinia anthemoides			✓	(L. angustata-like) L. hermanni			✓
				Hypochaeris glabra	✓		✓	L. angustatum	✓		✓
				Stylidium brunneum			✓	Halimolobos (varia) (ADS)			✓
				Briza minor	✓		✓	incrasata			✓

ADJ- Hypo. angustifolium
- S. marginata.

lollidos leaves

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP09

Record on Sheet

Date 13.12.96

- Column 1 plant name

- Column 2 plant number

• Column 3 flowering time- TICK if species flowering

- **Column 4** **identification** **check**

2nd Rescue
of Boko plot

[illegible]

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA SERPENTINE NP SITE NUMBER SERP10


DATE TRIP 29.10.96 RECORDERS AM

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location 32 22 31.2 Chatfield Rd

Geographic Location Latitude 32 22 S Longitude 115 59 E Altitude 60m
GPS ☒ Accuracy _____
Map Sheet Savannah PDOP = 2.4 (65A)

Photograph Name _____ Number #10 + #11 Roll #4

Topographic Position (Circle the correct response)
Upland _____ Lower slope Foot hills of scarp
Upper slope _____ Valley _____
Midslope _____ Washline _____

2. SITE DATA Circle the correct response

Slope flat _____ gentle 2 steep _____ Aspect N NE E SE S SW (W) NW

Surface Soil gritty, clayey gravel Depth to Rock 0-30cm Colour orange-brown

Exposed Rock type laterite % surface cover 40% size range 1-3cm

Litter 40 % surface cover Depth (cm) 2-5cm

Bare Ground 10 % surface cover




Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

colluvial deposits from upper slope
granite influenced sands
& laterite + dolerite clays

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class		2-10%	10-30%	30-70%	over 70%
TREES					
	over 15m	5-15m	under 5m	MALLEES over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		2-10%			
DOMINANT SPECIES		<i>E. marginata</i>			
SHRUBS					
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	SHRUBS under 0.5m
LIFE FORM					
COVER CLASS (%)			10-30% 2-10%	10-30%	
DOMINANT SPECIES			<i>X. acanthostachys</i> <i>H. crassifolia</i> <i>H. cucullata</i>	<i>L. multiflora</i> <i>H. lissocarpa</i> <i>Dry. amata</i>	<i>Mel. aff. scabra</i> <i>Calo. tomentosus</i> <i>Banksia camphorosma</i>
GRASSES					
	GRASSES	HERBS	SEDGES	over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)					
DOMINANT SPECIES					

4. VEGETATION CONDITION

Comments

Stunted

Mallee - Form jarrah

S/S: Good/Excellent condition. Above grazing

10

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP 10
Date 29.10.96
13.12.96(R)

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Eucalyptus marginata</i> ssp <i>deglutissima</i>	✓			<i>Drosera menziesii</i>	✓		1/2	<i>Lomandra sericea</i>	✓	✓	1/2
				<i>Proserpinaca stolonifera</i>	✓		1/2	<i>Lomandra hermaphrodita</i>	✓		✓
				<i>Gompholobium marginatum</i>	✓		1/2	<i>Homalosciadium leucocarpium</i>	✓		1/2
<i>Xanthorrhoea acanthostachya</i>	✓		1/2					<i>Laxmannia squarrosa</i>	✓		1/2
<i>Halkea undulata</i>	✓		1/2					<i>Trachypogon pilosa</i>	✓		1/2
<i>Verticordia acerosa</i> acerosa	✓		0					<i>Cassipouia pauciflora</i>	✓	✓	1/2
<i>Calamagrostis torulosa</i>	✓		1/2					<i>Lewyhookia stipitata</i>	✓		
<i>Leucopogon pulchellus</i>	✓	✓	1/2					<i>Actinotus leucocephalus</i>	✓		1/2
<i>Hypocalymma angustifolium</i>	✓	✓	1/2	GRASSES	✓		1/2	<i>Senecio decurcatus</i>	✓		1/2
<i>Pinetia imbricata peligera</i>	✓	✓	1/2	<i>Poa drummondiana</i>	✓		1/2	<i>Chaenactis corymbosa</i>	✓		1/2
<i>Acacia coarctata</i>	✓		1/2	<i>Briza maxima</i>	✓		1/2	<i>Hypochaeris glabra</i>	✓		1/2
<i>Gompholobium polydium</i>	✓		1/2	<i>Danthonia</i> sp.		✓		<i>Cassia micrantha</i> (NEW)	✓		1/2
<i>Andersonia lehmanniana</i>	✓		1/2	<i>Stipa</i> sp.		✓		<i>Dampiera alata</i>	✓		1/2
<i>Leuzea micrantha</i>	✓		0	<i>Poa canyophylla</i>	✓		1/2	<i>Lomandra nigricans</i>	✓		1/2
<i>Gompholobium knightianum</i>	✓		1/2	<i>Danthonia acerosa</i>	✓		1/2	<i>Goodenia caerulea</i>	✓		0
<i>Acacia laterifolia</i>	✓		1/2	<i>Stipa pycnostachya</i>	✓	✓	1/2	<i>Tetramea huda</i>	✓		1/2
<i>Asplenium pallidum</i>	✓		1/2	<i>Danthonia setacea</i>	✓		1/2	<i>Stylidium humifundum</i>	✓		1/2
<i>Boronia eriocarpa</i>	✓		0	<i>Neraxia aloepecuroides</i>	✓		1/2	<i>Stylidium bulbiferum</i> (NEW)	✓		1/2
<i>Hibbertia hypericoides</i>	✓		1/2	HERBS				<i>Tripterococcus brunis</i>	✓		1/2
<i>Baeckea campyrosmae</i>	✓		1/2	<i>Pterochaeta paniculata</i>	✓	✓	1/2	<i>Haemodorum laxum</i>	✓		✓
<i>Melaleuca aff. scarpa</i>	✓		1/2	<i>Lewyhookia pusilla</i>	✓	✓	1/2	<i>Polygonum aviculare</i>	✓		✓
<i>Hibbertia commutata</i>	✓		1/2	<i>Hypochaeris glabra</i>	✓	✓	1/2	<i>Stylidium carnosum</i>	✓		1/2
<i>Lambertia multiflora</i>	✓		1/2	<i>Xanthosia huegelii</i>	✓	✓	1/2				
<i>Thomasia fetosa glutinosa</i>	✓		1/2	<i>Hydrocotyle callicarpa</i>	✓	✓	1/2				
<i>Isopogon asper</i>	✓		1/2	<i>Millotia tenuifolia</i> - sterile	✓		1/2	SEDGES			
<i>Dryandra armata</i>	✓		1/2	<i>Patersonia gracilis</i>	✓	✓	1/2	<i>Schoenus grammophyllus</i>	✓		1/2
<i>Synaphea gracillima</i>	✓		1/2	<i>Patersonia pygmaea</i>	✓	✓	1/2	<i>Loxocarya fasciculata</i>	✓		1/2
<i>Dryandra linleyana linleyana</i>	✓		1/2	<i>Olearia paucidentata</i>	✓	✓	1/2	<i>Tetralix octandra</i>	✓		1/2
<i>Halkea cyclocarpa</i>	✓		1/2	<i>Patersonia occidentalis</i>	✓	✓	1/2	<i>Schoenus aff. brevisetis</i>	✓		1/2
<i>Halkea luscarnpa</i>	✓		1/2	<i>Lobelia gibbosa</i>	✓	✓	1/2				
<i>Hibbertia acerosa</i>	✓		1/2	<i>Pithecoloba pulchella</i>	✓	✓	1/2				
<i>Petrophile striata</i>	✓		1/2	<i>Stylidium hispidum</i>	✓	✓	1/2				

Not seen on previous

✓ SERP 3 / SERP 4

Entered as S. brevisetis

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP10

Record on Sheet

Date 29.10.96

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

NR: SERP00

LB. minor

Lepidospermum angustatum

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>E. marginata</i>	✓		✓	<i>Tetrathraea</i> sp - scabrous		✓	✓	<i>Haemodorum</i> sp.	✓		✓
				<i>Xanthoxia huaghi</i> seedlings	✓		✓	<i>Levenhookia pusilla</i>	✓		✓
<i>Hakea undulata</i>	✓			<i>Acacia</i> sp? drummondii seedling	✓			<i>Trachymene pilosa</i>	✓		✓
SHRUBS				6-leaflet Yellow Pea				<i>Malva penniculatum</i>	✓		✓
<i>Xanthorrhoea acanthostachya</i>	✓			Small Trifoliate seedling		✓	✓	<i>Chaenactis coronata</i>			✓
<i>Calorhamnus torulosis</i>	✓		✓	= <i>Gomph marginatum</i>	✓		✓	<i>Cassipouia racemosa</i>	✓		✓
MALLEES				<i>Pterocarya pinnata</i> ? (NF)	✓		✓	<i>Hypochaeris glabra</i>	✓		✓
<i>Lambertia multiflora</i>	✓		✓	<i>Acacia</i> sp - undulate leaves	✓		✓	<i>Hydrocotyle</i> sp.	✓		✓
<i>Melaleuca aff. scabra</i>	✓		✓	GRASSES				<i>Hemalia homalia</i>	✓		✓
<i>Dryandra nivea</i>	✓		✓	Climbing Drosera		✓	✓	- <i>Stipa</i> sp.	✓		✓
<i>Pimelia ciliatum</i>	✓		✓	leaves // whorl.				<i>Aiza caryophyllae</i>	✓		✓
SHRUBS								<i>Stachys</i> ? (NF)	✓		✓
<i>Hakea cyclocarpa</i>	✓							Tiny Annual grass (long twisted awn)			
<i>Hakea lissocarpa</i>	✓			<i>Astragalus/Leucopogon</i>				SEDGES			
<i>Hibbertia acerosa</i>	✓			↳ <i>A. lemniscata</i>		✓	✓	<i>Lepidospermum</i> sp. scabrous			✓
<i>Dryandra armata</i>	✓		✓	Marked leaves Twirling				<i>Loricaria fasciculata</i>	✓		✓
<i>Leucopogon pulcherrimus</i>	✓		✓	<i>Neurachne absepecuroides</i>	✓		✓	<i>Lomandra</i> sp. scabrous	✓		✓
<i>Hibbertia commutata</i>	✓		✓	<i>Briza maxima</i>	✓		✓	<i>Tetralix octandra</i>	✓		✓
<i>Isopogon asper</i>	✓		✓	HERBS				Small, white Asteraceae?			
<i>Hibbertia hypericoides</i>	✓		✓	<i>Larmanis squamosa</i>	✓		✓	<i>Olearia pauciflora</i>	✓		✓
<i>Baeckea cantharomasmæ</i>	✓		✓	<i>Stylidium brunonianum</i>	✓		✓	<i>Sandus olaraceus</i>			✓
<i>Petrophile swara</i>	✓			<i>Stylidium piliferum</i>	✓		✓	Black-base Schoenus	✓		✓
Andrit				<i>Stylidium carnosum</i>	✓		✓	<i>Astroloma pallida</i>	✓		✓
<i>Synedrella sp.</i>			✓	<i>caesia</i> sp? - NF //			✓	<i>Danthonia</i> sp.			✓
				<i>Drosera stolonifera</i>			✓	<i>Briza minor</i>			✓
<i>Hypocalymma angustifolium</i>			✓					<i>Gompholobium knightianum</i>			✓
				<i>Tripterocarococcus</i> sp	✓		✓	<i>Stylidium piliferum</i>	✓		✓
White-hirsute Asteraceae	✓		✓	<i>Thysanotus</i>				<i>Lomandra caespitosa</i>			✓
↳ large herb <i>Ptilotheca</i>				<i>Drosera menziesii</i>			✓	<i>Poa</i> sp.			✓
<i>caesia micrantha</i>				Small, viney Schoenus			✓	<i>Orthroschus/Paterosia</i> ?			✓
<i>Dampiera elata</i> // <i>Platysaceae</i>			✓					Mid Direct Seedling in vas-soo			✓
<i>Gompholobium</i> sp. - narrow, trifoliate			✓	<i>Thomasia</i> // <i>Lasiopetalum</i>			✓				

leaf

Schoenus? sp? - Tall
↳ black bases & Papery leaf margins

↳ Bollga??

lots of r. pygmaea

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP10

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Low. Pasivation
in SERP99 ✓

Date Resore

13.12.94

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
								Stipa sp.			
								Wild asteraceae			
								L. white tomentose			
								indumentum (NF)			
								Lomandra hermaphrodite			
								(as in dot SERP69)			
MALLEES								Paterosmia pygmaea			
				GRASSES				Lep. terete, smooth, large			
								Melolita			
SHRUBS								Nira sp.			
								Hydrocotyle			
								SEDGES			
								Triptero caracas brunonis			
								Hypocalymma angustifolium			
								Andersonia lehmanniana			
								Brachycola ?			
				HERBS				Pinellia imbricata			
								Compholobium knightianum			
								Clede			
								id H. acerosa			
								Not seen on vesicle			
								(NF)			
								Purple Lobelia			

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA -Upper Scrivener Rd Serpentine NP. SITE NUMBER SERP 11


DATE TRIP 30.10.96 RECORDERS AM

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location

Geographic Location Latitude 46-7 32 22 S Longitude 116 00 58.0 E Altitude 250m
GPS Accuracy 7 DOP = ±1000m
Map Sheet Jarrahdale

Photograph Name Number #12 = mistake
#13 Roll #4

Topographic Position (Circle the correct response)
Upland Upper slope Lower slope
Valley
Midslope 5 (see map) Washline

2. SITE DATA Circle the correct response

Slope flat gentle steep 4 Aspect 138° SE SE N NE E S SW W NW

Surface Soil loamy gravel Depth to Rock 0-30cm Colour orange-brown

Exposed Rock type latelite % surface cover 20 size range 1-3cm

Litter 40 % surface cover 5-10cm! Depth (cm)

Bare Ground 5 % surface cover)





Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Muir)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class	2-10%	10-30%	30-70%	over 70%
-------------	-------	--------	--------	----------

TREES				MALLEES	
	over 15m	5-15m	under 5m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)	10-30%	2-10%			
DOMINANT SPECIES	<i>E. marginata</i>	<i>E. marginata</i>			
SHRUBS			SHRUBS		
	over 2m	2m - 1.5m	1.5 - 1m	1-0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)				2-10%	77%
DOMINANT SPECIES				<i>M. hissarapha</i> <i>Xen. preissii</i>	<i>H. liso</i> <i>Dry. nivera</i> <i>Hib. hypericoides</i>
GRASSES		HERBS		SEDGES	
	over 0.5m			over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)		2-10%			2-10%
DOMINANT SPECIES		<i>Sc. nevola</i> sp. <i>Conostylis setosa</i> <i>Anigozanthus</i> sp.			<i>Loxocarya fasciculata</i> <i>Tetradlea celandina</i> <i>Lepidosperma</i> sp.
					<i>Lomandra homophylla</i> <i>Lomandra caespitosa</i>

4. VEGETATION CONDITION

Comments	BURNT 1993-1994. History of logging - A
Few weeds	4-4 1/2

2

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP - Lewis Rd SITE NUMBER SERP13


DATE TRIP 30.10.96 RECORDERS AM

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



900 from start gravel end of Lewis Rd

Road Location 39.0 29.3

Geographic Location Latitude 32 21 39.0 S Longitude 116 00 29.3 E Altitude 200m
 GPS ☒ Accuracy 5-1 - PDOP
 Map Sheet Jarrahdale

Photograph Name _____ Number # 14 Roll # 4

Topographic Position (Circle the correct response)
 Upland _____ Lower slope _____
 Upper slope (circled) Valley _____
 Midslope (circled) MIDSLOPE 3 Washline _____

2. SITE DATA Circle the correct response

Slope flat gentle steep Aspect N NE E SE S SW W NW

Surface Soil _____ Depth to Rock _____ Colour _____

Exposed Rock type _____ % surface cover _____ size range _____

Litter _____ % surface cover _____ Depth (cm) _____

Bare Ground _____ % surface cover _____

Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

SERP 12
 300m from 1st gate
 on Scindler Road
 HEAD EAST

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class					
	2-10%	10-30%	30-70%	over 70%	
TREES					
	over 15m	5-15m	under 5m	MALLEES over 8m	under 8m
LIFE FORM					
COVER CLASS (%)					
DOMINANT SPECIES					
SHRUBS					
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	SHRUBS under 0.5m
LIFE FORM					
COVER CLASS (%)					
DOMINANT SPECIES					
GRASSES		HERBS	SEDGES	over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)					
DOMINANT SPECIES					

4. VEGETATION CONDITION

Comments

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP11

Record on Sheet

Date 30.12.96

13.12.96(R)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES			No	FI	ID	SHRUBS (cont.)			No	FI	ID	HERBS (cont.)			No	FI	ID
<i>Eucalyptus marginata</i>	✓				R	<i>Thelymitra crinita</i>	✓	✓			R	<i>Trachymene pilosa</i>	✓	✓			R
Dy. Linden Linden					R	<i>Lomandra brittanii</i>	✓	✓			R	<i>Pterodroma paniculata</i>	✓	✓			R
<i>Hakea undulata</i> ADS	✓					<i>Cassytha poriformis</i>	✓	✓	✓			<i>Hydrocotyle calicarpa</i>	✓	✓			R
<i>Dryandra lindleyana</i> lindleyana		✓				<i>Trichostema spathulatum</i>	✓	✓			R	<i>Pentapeltis peltigera</i>	✓	✓			R
<i>Leucopogon capitellatus</i>	✓	✓	✓		R	<i>Drosera pallida</i>	✓	✓		✓		<i>Xanthosia candida</i>	✓	✓			R
<i>Hypocalymma robustum</i>	✓	✓			R	<i>Anigozanthus manglesii</i>	✓	✓		✓	R	<i>Crassedia varifolia</i> ^{var. albifolia}	✓	✓			R
<i>Bossiaea ornata</i>	✓	✓			R	<i>Drosera manglesii</i>	✓	✓				<i>Hyalosperma cotula</i>	✓	✓	✓		R
<i>Korea dioriximifolia</i>	✓	✓			R	<i>Lolus</i> sp.	✓			✓		<i>Agrostocrinum scabrum</i>	✓	✓	✓		R
<i>Phyllanthus calycinus</i>	✓	✓			R	GRASSES						<i>Goodenia caerulea</i>	✓	✓			R
<i>Grevillea pilulifera</i>	✓	✓	✓		R	<i>Stipa pycnostachya</i>	✓	✓	✓		R	ADS: <i>Thysanotus thysoides</i>	✓	✓			R
<i>Hibbertia commutata</i>	✓	✓	✓		R	<i>Neurachne alopecuroides</i>	✓	✓		✓	R	<i>Chaenodescilla corymbosa</i>	✓	✓	✓		R
<i>Hibbertia hypericoides</i>	✓	✓	✓		R	<i>Danthonia setacea</i>	✓	✓	✓		R	<i>Hydrocotyle calicarpa</i>	✓	✓			R
<i>Nymphalobium marginatum</i>	✓	✓			R	<i>Poa</i> sp. <i>drummondiana</i>	✓	✓	✓		R	<i>Phalangium paradoxum</i>	✓	✓	✓		R
<i>Lechenaurtia biloba</i>	✓	✓			R	<i>Vulpia muiroides</i> ^{myces}	✓	✓	✓		R	<i>Stylidium hispidum</i>	✓	✓	✓		R
<i>Pronaya fraseri fraseri</i>	✓	✓	✓		R	<i>Tetrarhena laevis</i>	✓			✓	R	<i>Crasspedia variaeflora</i>	✓	✓			R
<i>Hibbertia acares</i>	✓	✓	✓		R	<i>Aira caryophyllaea</i>	✓			✓	R	<i>Stylidium junceum</i>	✓	✓			R
<i>Acacia lateritica</i>	✓	✓	✓		R	HERBS						<i>Caesia microcarpa</i>	✓	✓			R
<i>Acacia teretifolia</i>	✓	✓	✓		R	<i>Burchardia congesta</i>	✓	✓			R	<i>Opercularia apiciflora</i>	✓	✓			R
<i>Pronaya fraseri fraseri</i>	✓	✓			R	<i>Dampiera alata</i>	✓	✓			R	NC <i>Anigozanthus?</i> ^{stille}	✓	✓			R
<i>Astroloma pallidum</i>	✓	✓			R	<i>Dampiera linearis</i>	✓	✓	✓		R	NC <i>Paterosia juncea?</i> ^{stille}	✓	✓			R
<i>Pimelea rubicilla</i> (not ciliata)	✓	✓			R	<i>Gonocarpus cordiger</i>	✓	✓	✓		R	<i>Centaurium euthyale</i>	✓	✓			R
<i>Sphaerolobium medium</i>	✓	✓			R	<i>Xanthosia huegelii</i>	✓	✓			R	<i>Linum trigynum</i>	✓	✓			R
<i>Acacia nervosa</i>	✓	✓			R	<i>Paterosia pugnax</i>	✓	✓			R	<i>Lomandra nigricans</i> (old)	✓	✓	✓		R
						<i>Corosyllis setosa</i>	✓	✓			R	<i>Lomandra micrantha</i>	✓	✓			R
<i>Astroloma ciliatum</i>	✓	✓			R	<i>Stylidium prunellatum</i>	✓	✓			R	SEDGES					
<i>Daveia decurrens</i>	✓	✓			R	<i>Senecio diaschides</i>	✓	✓	✓		R	<i>Tetaria capillaris</i>	✓	✓	✓		R
<i>Synplea gracillima</i>	✓				R	<i>Olearia paucidentata</i>	✓	✓			R	<i>Ochlopus unispiculatus</i>	✓	✓			R
<i>Hypochaeris glabra</i>	✓				R	<i>Opercularia echinocephala</i>	✓	✓	✓		R	<i>Tetaria octandra</i>	✓	✓			R
<i>Stylidium carnosum</i>	✓				R	<i>Gonocarpus cordiger</i>	✓	✓			R	<i>Mesomelaena gracilicarpa</i>	✓	✓			R
<i>Ptilotus manglesii</i>	✓				R	<i>Acaevola calliptera</i>	✓	✓	✓		R	<i>Loxocarya fasciculata</i>	✓	✓	✓		R
												NC <i>Schoenus</i> sp.	✓	✓			R

Ptilotus

Figure Assem1. 2wa. Assemblage_1 re-ordered according to species co-occurrences at the same quadrats and similarities in the overall composition of the quadrats. Quadrat codes are printed vertically.

	BBBCE	CCCC	G	KKMMMM	K
	BBBUL	UUUU	J	EERRRR	E
	24561	2345	2	122534	3
<i>Acanthophis pyrrhus</i>				*	
<i>Lerista aff. muelleri</i> C				*	
<i>Ctenotus saxatilis</i>				**	
<i>Diporiphora winnecke</i> i				**	
<i>Notoscincus ornatus</i>				**	
<i>Lerista kennedyensis</i>				*	
-----+-----+-----+-----+-----					
<i>Aprasia smithi</i>				*	
<i>Cyclodomorphus melanops</i>				*	
<i>Ramphotyphlops aff. grypus</i> C				*	
<i>Ctenotus piankai</i>				***	
<i>Menetia aff. greyii</i> C				****	
<i>Lerista bipes</i>				*	
<i>Ctenophorus femoralis</i>				* *	
<i>Ctenophorus clayi</i>				* * *	
<i>Ctenophorus rufescens</i>				* ***	
<i>Menetia aff. surda</i> A				* **	
<i>Varanus brevicauda</i>				** **	
<i>Ctenotus calurus</i>				*****	
<i>Ctenotus aff. hanloni</i> B				*****	*
<i>Ctenophorus aff. rubens</i> B			*	*****	
<i>Ctenotus pantherinus</i>		*		*****	
<i>Ctenotus iapetus</i>		****		*****	
<i>Cyclorana platycephalus</i>				*	
<i>Lerista planiventralis</i>	*****	**		* *	

Burchardie = prob (oesia (NF!))

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP17

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Date 30.10.96

g name if required
Stylidium pycnostachyum? ?? →
Stylidium juncum ✓ Basal rosette
 2 sp. *Cleistanthus* - Amplexed leaves

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>E. marginata</i>	✓			<i>Xanthosia hueglinii</i>				<i>Cassytha</i> sp (smooth)	✓		
								<i>Poa</i> sp	✓		
SHRUBS				<i>Gonocarpus</i> // <i>Fraxinaria</i>	✓			<i>Billardiera</i> sp	✓		
<i>Dryandra nivea</i>	✓			<i>Synaphaea</i> sp. petiolaris				<i>Ericid</i> → <i>Astroloma</i> ? <i>Leuc. cap.</i>			
<i>Hibb. hypericoides</i>	✓			<i>Gomph. marginatum</i>	✓			<i>Stipa elegantissima</i> pycno.	✓		
<i>Phyllanthus calycinus</i>	✓			<i>Acacia</i> sp. - pungent apices	✓			<i>Petersonia pygmaea</i>	✓		
MALLEES				6-leaflet thin Ped				<i>Burchardia longista</i>	✓		
<i>Hibb. commutata</i>	✓			<i>Chaemosila</i> ? <i>corymbosa</i>	✓			<i>Olearia paucidentata</i>	✓		
<i>Hibb. aserosa</i>	✓			GRASSES				<i>Angiozanthus mangleriana</i>	✓		
<i>Hakea pycnantha</i>	✓			<i>Acacia nervosa</i>	✓			<i>Crospedia</i> sp.			
<i>Xan. dreisii</i>	✓			<i>Astroloma pallidum</i>	✓			<i>Hydrocotyle</i> sp	✓		
SHRUBS				<i>Stylidium carnosum</i>	✓			<i>Trachymene pilosa</i>			
<i>Boscipia ornata</i>	✓			<i>Gonocarpus cordiger</i>	✓			<i>Wailia paniculata</i>			
<i>Conostylis setosa</i>	✓			<i>Drosera pallida</i>				SEDGES			
<i>Sphaerolobium medium</i>	✓			<i>Thelymitra crinita</i> (MATA) IN	✓			<i>Loxocarya fasciculata</i>	✓		
<i>Daviesia dreisii</i> ?	?			<i>Hypochaeris glabra</i>				<i>Lepidosperma angustatum</i>			
<i>Grevillea pilulifera</i>	✓			<i>Xanthosia</i> ? <i>candida</i>	✓			<i>Tetraria capillaris</i>	✓		
<i>Exco. Hovea</i> ? <i>charnifolia</i>				<i>Acacia drummondii</i>	✓			<i>Lepidosperma</i> thick smooth	✓		
<i>Pinelia</i> sp. - <i>sharpeana</i> ?	✓			HERBS							
<i>Hypocalymma rotundum</i>	✓			<i>Stylidium bulbosum</i>	✓			<i>Loxandora</i> sp. - basal inflor - heart shaped			
<i>Pentapeltis peltigera</i>	✓			<i>Scaevola</i> ? <i>platyphylla</i>	✓			<i>Tetraria octandra</i>	✓		
Unid grass - rough wide <i>Pennis</i> NF				<i>Trichodine spatulatum</i>	✓			<i>Lepidosperma</i> - rough terete	✓		
<i>Opercularia hispida</i>	?			<i>Small Helipterum</i> // <i>Rodanthe</i>				<i>Loxandora caespitosa</i> 'britanni'	✓		
<i>Lechenaultia biloba</i>	✓			<i>Leptura</i>				<i>Unid</i> sp. <i>Euphorb</i>			
<i>Loxandora</i> sp. - papery margins	✓			<i>Drosera menziesii</i>				<i>Cassia micrantha</i> (PF)			
<i>Dampiera</i> sp.	✓			<i>Dampiera linearis</i>	✓			<i>Laxmannia</i> or <i>wirey</i> <i>Schoenus</i>			
<i>Opercularia</i> - <i>edimocarpa</i>	✓			<i>Stylidium piluliferum</i>	✓						
<i>Tiny everlastina</i>	✓			<i>Agrostocrinum scabrum</i>	✓			ADS → <i>Hakea undulata</i>			
not <i>Helip. corbata</i>				<i>Sanigo</i> sp.	✓			<i>Loxandora dreisii</i>			
<i>Dampiera alata</i> // <i>Acacia wilsoniana</i>				<i>Vulpia</i> sp.	✓			<i>Astroloma ciliatum</i> ?			
<i>D. alata</i>				<i>Nauvacho alpeceuroides</i>	✓						

prob
answers

→ alate NE form.

→ Phallarium paradoxum (Nithasane)

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP 17

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Date 13/12.96

Rascare

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
								Goodenia caerulea			✓
								Conocarpus sp			✓
								Danthonia sp			✓
								Daviesia decurrens			✓
								Thelymitra sp			✓
								Lomandra ? rigida			✓
MALLEES								Acacia terete folia			✓
								Lobelia ?			
				GRASSES				Stylidium sp			✓
								Phlox maritima			✓
								Astroloma sp - tall weed			✓
SHRUBS								Stylidium canadense			✓
								canadense			
								SEDGES			
								Chamaecrista corymbosa			✓
								Argemone pulchella seedling?			✓
								Argemone pulchella			✓
								Senecio sp			✓
				HERBS				Lomandra prostrata (IN)			✓
								Tetrachena laevis			
								Astroloma sp			
								4 curled leaves			
								prob same as before			
								Aira caespitosa			✓
								Androsace			

73+13+6

= 92 (3-ppt)

Area
100 by
SUNDAY!

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA SERPENTINE NP ^{-Scrivener Rd} SITE NUMER SERP12

DATE TRIP 30-10-96 RECORDERS AM

DATE TRIP _____ RECORDERS _____


DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

In E. Iaelbe!

to 1st firebreak gate.
ODOMETER -

Mud Map Draw a sketch of the location of the site below



Road Location 32 22 57.0 116 00 41.4

Geographic Location Latitude 32 22 S Longitude 116 00 41.8 E Altitude 205m
GPS ☒ Accuracy 5m
Map Sheet Jarrahdale PDOP 4.6 2.7

Photograph Name _____ Number # 13 + # 14 Roll # 4

Topographic Position (Circle the correct response)
Upland _____ Lower slope _____
Upper slope ✓ Valley _____
Midslope _____ Washline _____

2. SITE DATA Circle the correct response

Slope flat gentle steep Aspect N NE E SE S SW W NW

Surface Soil loamy clay Depth to Rock 30-40 Colour grey-brown

Exposed Rock type granite % surface cover 10 size range 60cm - 150cm

Litter 80 % surface cover Depth (cm) 20-30cm

Bare Ground 2% % surface cover




incl. rocks = 12%

Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Muir)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class					
	2-10%	10-30%	30-70%	over 70%	
TREES					
	over 15m	5-15m	under 5m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		30-70%			
DOMINANT SPECIES		<i>E. laevis</i> <i>E. marginalis</i>			
SHRUBS					
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	SHRUBS under 0.5m
LIFE FORM					
COVER CLASS (%)				10-30	10-30%
DOMINANT SPECIES				<i>H. stenocarpa</i> <i>Nerax cuneatum</i> <i>Boswellia brevifolia</i>	<i>Melaleuca scabra</i> <i>Leucop. Hibb. hypericoides</i>
GRASSES		HERBS	SEDGES	over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)					30-70%
DOMINANT SPECIES					<i>Melaleuca tetragona</i>

4. VEGETATION CONDITION

Comments

4/5. No track & heavy voo grazing. All long unburnt. *E. laevis* conspicuous in landscape. Part of band of *E. laevis* running into valley. Emergent *X. preissii* & influence from adjacent granite v. sedge understory.

ADS. Thicket of *H. bifurcata*

V. tall thicket of *H. bifurcata* adjacent

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP12

Record on Sheet

Date 30.10.76

13.12.76(R)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
Eucalyptus calophylla	✓		YR	Dry. lindle... lindle	✓		YR	Paterisoria pygmaea	✓	✓	YR
Euc. laeliae	✓		YR	Hakea amplexicaulis	✓		YR	Billardiera parviflora gutatta	✓	✓	YR
Darwinia humnoides	✓	✓	R	Hakea lussorephta	✓		R	Cassutha paniciformis	✓	✓	✓
Allocasuarina humilis	✓	✓	YR	Gompholobium marginatum	✓		R	Cassutha racemosa	✓	✓	✓
Eriostemon spicata	✓	✓	YR					Cassutha glabella	✓	✓	YR
Hakea drooparpa	✓	✓	YR					Lomandra brittanii	✓	✓	✓
Pinalea preissii	✓	✓	YR					Lomandra purpurea	✓	✓	✓
Hypocalymma angustifolium	✓	✓	YR					Lomandra preissii (narrow)	✓	✓	✓
Hibbertia communata	✓	✓	YR	GRASSES				Wahlenbergia preissii	✓	✓	
Hibbertia hypericoides	✓	✓		Danthoria setacea	✓	✓	YR	Goodenia caerulea	✓	✓	R
				Stipa pycnostachya	✓	✓	YR	Haemodorum discolor	✓	✓	YR
Hakea trifurcata	✓	✓	YR	Poa drummondiana	✓	✓	YR	Drosera menziesii	✓	✓	YR
Olearia pulcherrima	✓	✓	YR	Tetraulhera laevis	✓	✓	YR	Trichostema spatulatum	✓		YR
ADS: Olearia bipinnatifida			R	Danthoria acerosa	✓	✓	YR	Stylidium bulbiferum	✓		YR
Melaleuca aff. scrubra	✓	✓	YR	Amphipogon sp.	✓	✓	YR	Xanthosia huegelii	✓		R
Dryandra bipinnatifida	✓	✓	YR	Poa campophylla	✓	✓	R				
Anderssonia lehmanniana	✓	✓	YR	Neracis abequeiroidea	✓		YR				
Petrophile striata	✓	✓	YR								
Acacia pulchella pulchella	✓	✓	YR	HERBS							
Bessidea sp. Nawaona (id=)	✓	✓	YR	Dampiera alata	✓	✓	YR				
Baedeia campyrosmae	✓	✓	YR	Dampiera linearis	✓	✓	YR				
Kunzea micrantha	✓	✓	YR	Agrostovinum scabrum	✓	✓	YR				
Leucopogon pulchellus	✓	✓	YR	Scaevola calliptera	✓	✓	YR	JEDGES			
Olax benthamiana	✓	✓	YR	Burchardia congesta	✓	✓	YR	Mesomelaena tetragona	✓	✓	YR
				Caesia micrantha	✓	✓	YR	Tetralia capillaris	✓	✓	YR
Nemcia dilatata	✓	✓	YR	Arthropodium capillipes	✓	✓	YR	Tetralia octandra	✓	✓	YR
Acacia alata alata	✓	✓	YR	Stylidium brunonianum	✓	✓	YR	Mesomelaena graciliceps	✓	✓	
Lechenaultia biloba	✓	✓	R	Caesia micrantha	✓	✓	YR	ADS: Cyathodactylus aeneus	✓		
Templetonia kilorn	✓	✓	R	Operscularia apiciflora	✓	✓	YR				
Xanthorrhoea preissii	✓		YR	Gonocarpus cordiger	✓	✓	YR				
Dryandra ovata	✓		YR	Stylidium hispidum	✓	✓	YR				
Xanthorrhoea gracilis	✓		YR	Conesperma virgatum	✓	✓	YR				

Figure Assem2. 2wa. Assemblage_2 re-ordered according to species co-occurrences at the same quadrats and similarities in the overall composition of the quadrats. Quadrat codes are printed vertically.

	BBNMMM	BCCCCCEEZZZZPPPP	NNNNNNNZ	B	BMPN	CMM	BKM
	BBERRR	BUUUUUULLLUUUUEEEE	AAAAEEUU	O	OREE	UDD	OED
	254132	4453621221342534	23451235	3	2415	112	524
<i>Aprasia fusca</i>		* *					
<i>Lerista haroldi</i>		*					
<i>Aprasia haroldi</i>			*				
<i>Ramphotyphlops australis</i>			*				
<i>Pygopus lepidopodus</i>			*				
<i>Tympanocryptis adelaidensis</i>			*				
<i>Vermicella bimaculata</i>			*				
<i>Ctenophorus aff. rubensA</i>			*				
<i>Neobatrachus pelobatoides</i>				*			
<i>Crenadactylus ocellatus</i>		*	*	*****			
<i>Ctenotus alleni</i>				** ***			
<i>Cryptoblepharus carnabyi</i>			*		*		
<i>Ctenotus lesueurii</i>			*		*		
<i>Lerista humphriesi</i>			*				
<i>Lerista decora</i>			*** *		*		
<i>Ctenophorus maculatus</i>			**	**	*		
<i>Lerista aff. kendrickiA</i>			*	** *			
<i>Delma aff. australisA</i>			*	*			
<i>Vermicella multifasciata</i>			*				
<i>Lerista aff. praepeditaB</i>		** *	****		*		
<i>Lerista varia</i>		*	****	*			
<i>Tiliqua rugosa</i>			****		*		
<i>Menetia aff. surdaB</i>			* **	* ****			
<i>Diplodactylus ornatus</i>		*** **	***				
<i>Lialis burtonis</i>	*	*	** **				***
<i>Ctenotus fallens</i>		** ** *	**	***			
<i>Diplodactylus aff. alboguttatusA</i>		* * ** *	***				

Grass in 200. 22-

10 10 10 10 NB also found in SERP 11

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP 12

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Date 13-10-96

13.12.96

resoure

[illegible]

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Serpentine NP ^{→ Lewis Rd} SITE NUMBER SERP13


DATE TRIP 30-10-96 RECORDERS AM

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



Road Location Lewis Rd, Serpentine

Geographic Location	Latitude	S Longitude	E Altitude
	GPS	Accuracy	- see other sheet
	Map Sheet	Jarrahdale	

Photograph	Name	Number	Roll
		#114	Roll #4

Topographic Position	(Circle the correct response)
Upland	Lower slope
Upper slope	Valley
<u>Midslope</u>	Washline

2. SITE DATA Circle the correct response

Slope flat gentle steep Aspect N NE E SE S SW W NW

Surface Soil loamy clay ^{day loam} Depth to Rock 10-30cm Colour orange - dk brown

Exposed Rock type granite % surface cover 1% size range 5-20cm

Litter 80 % surface cover Depth (cm) 20-30cm


Bare Ground | % surface cover

Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class					
	2-10%	10-30%	30-70%	over 70%	
TREES					
	over 15m	5-15m	under 5m	MALLEES over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		10-30%			
DOMINANT SPECIES		<i>E. laevis</i> <i>E. calophylla</i>			
SHRUBS					
	over 2m	2m - 1.5m	1.5 - 1m	1-0.5m	SHRUBS under 0.5m
LIFE FORM					
			2-1.5m		<i>Racoma capensis</i> <i>Astragalus</i> <i>H. lissocarpus</i>
COVER CLASS (%)		10-30%	10-30%	10-30%	30-70%
DOMINANT SPECIES		<i>H. trifurcata</i>	<i>Xan. acanthostachya</i> <i>H. undulata</i>	<i>Melaleuca scabra</i> <i>Hypo angustifolium</i>	<i>Hibb. hypoleucoides</i> <i>Dag. nivalis</i> <i>Leucophaea pulcherrima</i>
GRASSES		HERBS	SEDGES	over 0.5m	
LIFE FORM					
COVER CLASS (%)	2-10%	MIXED HERBS 2-10%		10-30%	
DOMINANT SPECIES	<i>Briza maxima</i>	<i>Lagenifera longii</i> <i>Stylidium bulbiferum</i> <i>Caesia burchardii</i>		<i>Lepidosperma</i> <i>Tetradlea</i> <i>Tetradlea capillaris</i>	

4. VEGETATION CONDITION

Comments 3 to 3 1/2 / 5. V. weedy - influx from adjacent pasture
Long time unburnt - senescent *H. undulata*

3-2.5.

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIFIC PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP 13

Record on Sheet

Date 20.10.96

14.12.96(R)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES				No	FI	ID	SHRUBS (cont.)				No	FI	ID	HERBS (cont.)				No	FI	ID
<i>Eucalyptus calophylla</i>	✓	✓	✓				<i>Haemodorum? laxum</i>	✓	✓	✓				<i>Romulea rosea</i>	✓	✓	✓			✓/R
<i>Euc. laetiae</i>	✓	✓	✓				<i>Haemodorum discolor</i>	✓	✓	✓										
<i>Macrotanais viedlii</i>	✓	✓	✓				<i>Drosera menziesii</i>	✓	✓	✓				<i>Centrolepis drummondiana</i>	✓	✓	✓			✓
<i>Xanthorrhoea acanthostachya</i>	✓	✓	✓				<i>Drosera erythrorhiza</i>	✓	✓	✓				<i>Levenhookia pusilla</i>	✓	✓	✓			✓/R
<i>Leucopogon pulchellus</i>	✓	✓	✓				<i>Drosera pulchella???</i>							<i>Homaloscladum homalocarpum</i>	✓	✓	✓			✓
<i>Leucopogon capitellatus</i>	✓	✓	✓											<i>Wahlenbergia preissii</i>	✓	✓	✓			✓
<i>Melaleuca aff. scabra</i>	✓	✓	✓				<i>Hypochaeris glabra</i>	✓	✓	✓				<i>Crassulca pedicellata</i>	✓	✓	✓			✓
<i>Allocasuarina humilis</i>	✓	✓	✓				<i>Centrolepis aristata</i>	✓	✓	✓				<i>Boya sphaerocephala</i>	✓	✓	✓			✓/R
<i>Brevillea pilulifera</i>	✓	✓	✓				GRASSES							<i>Palafoxia pygmaea</i>	✓	✓	✓			✓/R
<i>Hibbertia hypoleucoides</i>	✓	✓	✓				<i>Briza maxima</i>	✓	✓	✓				<i>Isotria hypocrateriformis</i>	✓	✓	✓			✓
<i>Baeckea campochloasmae</i>	✓	✓	✓				<i>Briza caryophylla</i>	✓	✓	✓				<i>Trifolium cernuum</i>	✓	✓	✓			✓/R
<i>Astragalus pallidus</i>	✓	✓	✓				<i>Briza minor</i>	✓	✓	✓				<i>Comesperma ciliatum</i>	✓	✓	✓			✓/R
<i>Dryandra lindleyana lindleyana</i>	✓	✓	✓				<i>Stipa pycnostachya</i>	✓	✓	✓				<i>Cassipoua porriiformis</i>	✓	✓	✓			✓/R
<i>Hypocalymma angustifolium</i>	✓	✓	✓				<i>Danthonia setacea</i>	✓	✓	✓				<i>Lomandra brittanii</i>	✓	✓	✓			✓
<i>Hakea linearis</i>	✓	✓	✓				<i>Avena barbata</i>	✓	✓	✓				<i>Lomandra purpurea</i>	✓	✓	✓			✓/R
<i>Hakea trifurcata</i>	✓	✓	✓				<i>Poa drummondiana</i>	✓	✓	✓				<i>Lomandra micrantha</i>	✓	✓	✓			✓/R
<i>Hypocalymma angustifolium</i>	✓	✓	✓				<i>Neuvachne alopecuroides</i>	✓	✓	✓				<i>Mona denia bracteosa</i>	✓	✓	✓			✓
<i>Hibbertia communata</i>	✓	✓	✓				<i>Tetrathera laevis</i>	✓	✓	✓				<i>Thelymitra crinita</i>	✓	✓	✓			✓
<i>Lechenaultia biloba</i>	✓	✓	✓				HERBS							<i>Caladenia marginatum</i>	✓	✓	✓			✓
<i>Acacia pulchella pulchella</i>	✓	✓	✓				<i>Dampiera linearis</i>	✓	✓	✓				<i>Microris media media</i>	✓	✓	✓			✓
<i>Acacia lateriticola</i>	✓	✓	✓				<i>Stylidium humionum</i>	✓	✓	✓				<i>Crassipedia sp. variabilis</i>	✓	✓	✓			✓
<i>Synaphea gracillima</i>	✓	✓	✓				<i>Stylidium bulbiferum</i> (undet)	✓	✓	✓				<i>Galium sp.</i>	✓	✓	✓			✓
<i>Hakea undulata</i>	✓	✓	✓				<i>Tetrathera nuda</i>	✓	✓	✓				<i>Levenhookia stipitata</i>	✓	✓	✓			✓
<i>Petrophile striata</i>	✓	✓	✓				<i>Lagenifera huerfii</i>	✓	✓	✓				SEDGES						
<i>Gompholobium marginatum</i>	✓	✓	✓				<i>Opercularia apiciflora</i>	✓	✓	✓				<i>Tetraria capillaris</i>	✓	✓	✓			✓/R
<i>Eriostema spicatum</i>	✓	✓	✓				<i>Centaurium erythraea</i>	✓	✓	✓				<i>Tetraria octandra</i>	✓	✓	✓			✓/R
	✓	✓	✓				<i>Chaenactis corymbosa</i>	✓	✓	✓				<i>Schoenus grammaphyllus</i>	✓	✓	✓			✓/R
<i>Ursinia altheroides</i>	✓	✓	✓				<i>Thysanotus manglianus</i> (strig)	✓	✓	✓				<i>Juncus caespiticius</i>	✓	✓	✓			✓
<i>Stylidium comosum</i>	✓	✓	✓				<i>Burchardia congesta</i>	✓	✓	✓				<i>Loxocarya fasciculata</i>	✓	✓	✓			✓/R
<i>Trachymene pilosa</i>	✓	✓	✓				<i>Lomandra opartea</i>	✓	✓	✓				<i>Meconella tetragona</i>	✓	✓	✓			✓/R
<i>Isopogon asper</i>	✓	✓	✓				<i>Cardenia caerulea</i>	✓	✓	✓										
							<i>Cacia micrantha</i>	✓	✓	✓										

<i>Vermicella littoralis</i>		**	*****						
<i>Cyclodomorphus celatus</i>		*	** *		***				
<i>Ctenophorus badius</i>	*	*****	*		* *			***	*
<i>Lerista elegans</i>	*****	*****	***		*				
<i>Morethia lineoocellata</i>	**	****	*****				****		
<i>Lerista lineopunctulata</i>	***	*****	* *****		***				
<i>Lerista aff. praepedita</i> A		*****	****		***** *				
<i>Menetia amaura</i>	*	* **	*** * **		****		*		
<i>Arenophryne rotunda</i>			* ****		*				
<i>Diplodactylus spinigerus</i>			*****						
<i>Pletholax gracilis</i>			* * **						
<i>Tympanocryptis parviceps</i>			* ***						
-----+-----+-----+-----+-----+-----									
<i>Ctenotus aff. hanloni</i> A		*** *							
<i>Ramphotyphlops aff. grypus</i> B		*							
<i>Delma butleri</i>		** *	*						
<i>Menetia aff. greyii</i> B		****							
<i>Pseudonaja nuchalis</i>		*** *							
<i>Diplodactylus rankini</i>		* *							
<i>Demansia psammophis</i>		*							
<i>Delma nasuta</i>		*	*						
<i>Diplodactylus michaelsoni</i>			*						
<i>Pseudechis australis</i>			*						

COND T = 3 1/2

Anagallis-like weed - 10" leaves // C. eschscholae

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP13

Record on Sheet

Date 30/10/96

14.12.96 (2)

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

Tiny annual grass


Dampiera elata (ADS)

C. drummondii

No plants in plot

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
Acacia				Synsphaera sp.	✓		///	Haemodorum sp.	✓		///
E. izabaei	✓		///	Macropidia reidii	✓			Cassutha sp.	✓		///
E. calophylla			///	Drosera spaldii				Melaleuca aff. scabra	✓		///
SHRUBS				Hibbertia commutata	✓		///	Tetrarrhena laevis	✓		
Xen. acanthostachya	✓		///	Neurochne dioscoreoides	✓			Avena sp.			///
Hypocalymma angustifolium	✓		///	Lagenaria hughii	✓		///	Poa sp.			///
MALLEES				Caesia nicaeana	✓		///	Danthonia sp.	✓		///
Azorella / Sturtia	✓			Burchardia congesta	✓		///	Drosera menziesii			///
Comesperma virgatum			///	GRASSES				Schoenus - blackbase	✓		
Hakea undulata	✓			Buzia maximae	✓		///	Calladenia flava (NFI)			///
Hakea lucasiorum	✓		///	Stipa sp.	✓		///	Acacia drummondii	✓		///
Hakea trifurcata	✓			Maize - hirsute grass → Poa?				Levenhookia pusilla			///
SHRUBS				Thelypodium crinita	✓			Loxocarya fasciculata	✓		///
Oreocarya echinophylla	✓		///	Dryandra nivea	✓		///	SEDGES			
Barkeria cantherosmae	✓		///	Drosera erythrorhiza			///	Lomandra dracunculifolia	✓		///
Hibbertia commutata	✓		///	Romulea rosea	✓		///	Tetraria capillaris	✓		///
Dampiera linearis	✓		///	Hypochaeris glabra	✓			Lepidosperma hirsuta	✓		///
Leucopogon pulcherrimus	✓		///	Unknown → Centropogon drummondii OK			///	Tetraria octandra scabra	✓		///
Grevillea seedling	✓		///	HERBS				Lomandra hirsutissima	✓		///
Psidium				Caesia / Burchardia?				NA = Nahrbergia sp.	///		
Hibbertia hypericoides	✓		///	Arctostaphylos	✓		///	Tiny Levenhookia with			
Allocas humilis			///	Stylidium brunonianum	✓		///	Borgia	✓		///
1 seedling			///	Stylidium bulbiferum	✓		///	phaceloides	✓		///
Acacia pulchella	✓		///					Pink Nat. lily			
Grassodia sp.				Tetrarrhena? - spaldii? (nuda)			///	Astroloma sp.	///		
Levenhookia	✓		///	Centropogon aristata				Trachymene pilosa			
Gaulth (chickweed)			///	Ursinia antheroides				Chamaecrista polytricha	✓		///
Buzia minor	✓		///	Thysanotus patersonii			///	Lomandra caespitosa	✓		
Ara canophyllae	✓		///	Dampiera linearis	✓		///	Petrophile striata	✓		
Homophyllum marginatum	✓		///	Stylidium carnosum				Patersonia pugnaea	✓		///
Stachnoidia (NFI)								Hydrocotyle			
H. trifurcata	✓		///								

Stipa
Stipa



BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA SERPENTINE National Park SITE NUMBER SERP14


DATE TRIP 3/10/96 RECORDERS AM.

DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Mud Map Draw a sketch of the location of the site below



x In remote location on mountain side. WEST FACE OF RIDGE NEAR SUMMIT!

Road Location off Chatfield Rd

Geographic Location Latitude 32 22 S Longitude 116 00 2.5 E Altitude 130m
GPS Accuracy PDOP = 7.0
Map Sheet Jaural date

Photograph Name _____ Number #19 + #20 / Rd #4

Topographic Position (Circle the correct response)
Upland _____ Lower slope _____
Upper slope ✓ Valley _____
Midslope _____ Washline _____

2. SITE DATA Circle the correct response

Slope flat gentle steep Aspect N NE E SE S SW (W) NW

Surface Soil gr Depth to Rock 0-20cm Colour pk-MED BROWN

Exposed Rock type granite % surface cover 45% size range 60-150cm

Litter 65 % surface cover Depth (cm) 10cm

Bare Ground 0 % surface cover

with Rock 40%

↳ Allocas. hugeliana needles


BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class	2-10%	10-30%	30 - 70%	over 70%
-------------	-------	--------	----------	----------

TREES				MALLEES	
	over 15m	5 - 15m	under 5m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)			10-30%		
DOMINANT SPECIES			<i>Allocas. huegliana</i>		

SHRUBS				SHRUBS	
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)			2-10%	30-70%	
DOMINANT SPECIES			<i>Grevillea pteridifolia</i>	<i>Mel. viminalis</i> <i>Dodonaea sp.</i>	

GRASSES		HERBS	SEDGES	over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)		10%-20%			
DOMINANT SPECIES		<i>Tachypogon sp.</i> <i>Xanthosia candida</i> <i>Dioscorea hastifolia</i>			

4. VEGETATION CONDITION

Comments

some annual weeds 1-5

515

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIFIC PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP14

Record on Sheet

Date 14.12.96 (R)

31.10.96

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Alchornea lucida</i>	✓		R					<i>Thysanotus marginatus</i>	✓		✓
								<i>Cheilanthes austrotenuifolia</i>	✓		R
								<i>Sonchus oleraceus</i>	✓		R
<i>ADP. Xanthorrhoea preissii</i>	✓		R					<i>Tragalis arvensis</i>	✓		R
<i>Grevillea erdlicheriana</i>	✓		R					<i>Lechenicea pusilla</i>	✓		R
<i>Dodonaea ceratocarpa</i>	✓		R					<i>Daucus glochidiatus</i>	✓		✓
<i>Calyptrix acutifolia</i>	✓		R	<i>Briza minor</i>	✓		R	<i>Hypochaeris glabra</i>	✓		R
<i>Hibbertia spicata</i>	✓		R	GRASSES				<i>Borja sphaerocephala</i>	✓		R
<i>Verticordia pennigera</i>	✓		R	<i>Briza maxima</i>	✓		R	<i>Wahlenbergia preissii</i>	✓		✓
<i>Thymalium ledifolium</i> var. <i>var.</i>	✓		R	<i>Danthonia sp.</i>	✓		✓	<i>Burchardia congesta</i>	✓		R
<i>Mirbelia spinosa</i>	✓		R	<i>Mira canyodactyla</i>	✓		✓	<i>Artropodium capillipes</i>	✓		R
<i>Darwinia citriodora</i>	✓		R	<i>Danthonia setacea</i>	✓		R	<i>Lobelia heterophylla</i>	✓		R
<i>Hovea pungens</i>	✓		R	<i>Danthonia acerosa</i>	✓		R	<i>Haemodorum laxum</i>	✓		R
<i>Mitras. laegeliana</i>	✓		R	<i>Erihartia longifolia</i>	✓		✓	<i>Drosera macrantha</i>	✓		✓
<i>Metaleuca radula</i>	✓		R	<i>Stipa compressa</i>	✓		✓	<i>Trachymene pilosa</i>	✓		✓
<i>Arven</i>				<i>Neurachne alopecuroides</i>	✓		R	<i>Drosera patida</i>	✓		✓
<i>Baedeia camphorosmea</i>			R	<i>Avena sp.</i>			✓	<i>Drosera menziesii</i>	✓		✓
<i>Baedeia camphorosmea</i>			R	HERBS				<i>Craspedia variabilis</i>	✓		✓
				<i>Pterochaeta pauciculata</i>	✓		R	<i>Lotus sp.</i>	✓		✓
				<i>Stylidium bulbiferum</i> (uncf)	✓		R				
				<i>Ursinia anthemoides</i>	✓		R				
				<i>Tricoryne elatior</i>	✓		R				
				<i>Caesia micrantha</i>	✓		R				
				<i>Bartsia trixago</i>	✓		R	SEDGES			
				<i>Scaevola glandulifera</i>	✓		R	<i>Schoenus unispiculatus</i>	✓		R
				<i>Actinotus leucocephalus</i>	✓		R	<i>Desmodium aspera</i>	✓		R
				<i>Trachymene coerulea</i>	✓		R	<i>Schoenus nanus</i>	✓		✓
				<i>Rhodanthe corymbosa</i>	✓		✓				
				<i>Haemodorum discolor</i>	✓		R				
				<i>Dioscorea hastifolia</i>	✓		R				
				<i>Xanthosia candida</i>	✓		R				

Figure Assem3. 2wa. Assemblage_3 re-ordered according to species co-occurrences at the same quadrats and similarities in the overall composition of the quadrats. Quadrat codes are printed vertically.

	BWMMMMWGGGGMMMW	BBBB	GNPNNNWWNN	BBBBCNNCCPNPPKMM	CECZZZ	EZZ	KKK	BCN	P
	BORRDROJJJJDDDD	OOOO	JAEEEEEOEE	BBBOUAAUUEAEEEEERR	ULUUUU	LUU	EEE	BUA	E
	1225314235412455	1342	1242151334	24555342335521234	416453	221	345	311	1
<i>Cryptoblepharus plagiocephalus</i>									
<i>Egernia depressa</i>	**	*		**					
<i>Denisonia fasciata</i>		*							
<i>Gemmatophora longirostris</i>	*				*				
<i>Pseudophryne guentheri</i>	*					*			
<i>Furina ornata</i>			*						
<i>Morethia butleri</i>			*						
<i>Morethia obscura</i>			*						
<i>Delma aff. australisB</i>		*	*	*					
<i>Egernia stokesii</i>			*						
<i>Ctenotus uber</i>	*		*				***		
<i>Lerista aff. kendrickiB</i>							*		
<i>Lerista aff. muelleriD</i>				*			***		
<i>Diplodactylus aff. pulcherB</i>	**	**	*					*	
<i>Neobatrachus wilsmorei</i>	*****			**	**				
<i>Ctenotus schomburgkii</i>	**	*****	*	***	**	*	*		
<i>Rhynchoedura ornata</i>	*	*****	*	*					
<i>Varanus caudolineatus</i>	**	*	*	*			*		
<i>Pseudonaja modesta</i>	*	**	*					*	
<i>Ctenophorus inermis</i>	*	*	*****	*	*	*		***	
<i>Eremiascincus fasciolatus</i>		*	*		*	**			
<i>Varanus gouldii</i>	**	*	*	*	*	*		*	
<i>Diplodactylus conspicillatus</i>	*	*	*	*	**	*			
<i>Ramphotyphlops hamatus</i>	*	*	*	*	*	*			
<i>Ramphotyphlops aff. grypusA</i>	*	****	*	*					
<i>Diplodactylus aff. alboguttatusB</i>	**	*	*	*	*	*			
<i>Cyclorana maini</i>	*	*	****		*	*			

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

SITE No	SERP14
Date	31-10-96

Record on Sheet

- 5-leaflet pea

Loxocarya flexuosa ✓ //

TREES				SHRUBS (cont.)				HERBS (cont.)			
No	FI	ID		No	FI	ID		No	FI	ID	
			<i>Allocasuarina huegiana</i>	✓			<i>Stylidium bulbiferum</i>	✓			<i>Haemodorum</i> sp.
							<i>Aragalis arvensis</i>	✓			<i>Trachymene</i> p. sp. LARGE
			SHRUBS				<i>Xanthosia candida</i>	✓			Small white Asteraceae
			<i>Melaleuca radula</i> / <i>Boerhaavia</i>	✓			<i>Dioscorea pallida</i>				metroid
			<i>Grevillea enderlicheriana</i>	✓			<i>Dioscorea purgens</i> ?				White Scroph. (Large)
							<i>Hovea purgens</i> ↓	✓			<i>Hippocrepis glabra</i>
			MALLEES				<i>Sandus oblanceolatus</i>				<i>Trichogone elatior</i>
			<i>Mirbelia purgens</i> Spinescent	✓			<i>Thysanotus patersonii</i> (climbing)	✓			<i>Lepidosperma angustatum</i>
			<i>Levenhookia pusilla</i>	✓			GRASSES				
			<i>Burchardia congesta</i>	✓			<i>Avena</i> sp.	✓			<i>Grevillea?</i> // <i>Trymellum?</i>
							<i>Briza maxima</i>	✓			
			SHRUBS				<i>Briza minor</i>	✓			<i>Scaevola</i> sp.
							<i>Borja sphaerocarpa</i>	✓			<i>Wahlenbergia</i> sp.
			<i>Baeckea camphorosmae</i>	✓							SEDGES
			<i>Hibbertia spirata</i>	✓			<i>Dianthonia</i> sp.	✓			Rough sedge // Sedge (Hort)
			<i>Tetete leaf</i> <i>Haemodorum</i>	✓			<i>Dioscorea monjessii</i>				
			NEW ACHNE <i>Alopecuroides</i>	✓							<i>Waitia pinniculatum</i>
			<i>Ursinia anthemoides</i>	✓							<i>Trachymene pilosa</i>
			<i>Calytrix acutifolia</i>	✓			HERBS				
							<i>Crassedia</i> sp.				<i>Aragalis arvensis</i>
			<i>Darwinia citriodora</i>	✓			<i>Archipogon capillipes</i>	✓			
							<i>Dioscorea nasrifolia</i>	✓			<i>Aira caryophyllae</i>
							<i>Austrotetrarrhena</i>				<i>Vulpin</i> sp.
							<i>Cheilanthes austrotenuifolia</i>	✓			<i>Stylidium bulbiferum</i>
							<i>Cassia micrantha</i>	✓			
							<i>Ursinia anthemoides</i>	✓			<i>Centaurium</i> / <i>Isoplepis</i> (small)

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP14

Record on Sheet

Date 14.12.96

Rescore

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
								Actinotus sp	/		
								Lantana-like herb	// T. coerulea		
								Haemadion sp	//		
								lthin strapleaf			
								lcs noted before on			
								1st score)	//		
MALLEES								Hibbertia spicata	///		
								Loxocarya flexuosa	//		
				GRASSES				Tynantrum sp (seedling)	///		
								Lavenderia pumila	///		
								Borya sphaerocarpa	///		
SHRUBS								Black-base Schoenus	//		
								SEDGES			
								Dodonaea - Caratophylla	///		
								Lobelia? NF			
								in bud.	//		
				HERBS				Tyrcorne dafior	///		
								Naitzia panniculatum	///		
								Xanth. prekli (ADS)			
								Verticordia penigera			
								Burchardia congesta			

BUSHLAND PLANT SURVEY RECORDING SHEET 1 - use pencil only

BUSHLAND AREA Falls Road Serpentine N.P. SITE NUMBER SERP15

DATE TRIP 8.12.96 RECORDERS AM


DATE TRIP _____ RECORDERS _____

DATE TRIP _____ RECORDERS _____

1. LOCATION of the QUADRAT

Creekside Vegetation

Mud Map Draw a sketch of the location of the site below



No Rescon
N
E

Road Location Falls Rd Serpentine N.P.

Geographic Location Latitude 32° 22' 11.21" S Longitude 116° 00' 32.23" E Altitude 55m (map)
GPS Accuracy NO FIX!!
Map Sheet

Photograph Name Number Rd1 #5 #19, 20, 21, 22

Topographic Position (Circle the correct response)
Upland Lower slope
Upper slope Valley
Midslope Washline creekside flat 10

2. SITE DATA Circle the correct response

Slope flat gentle steep Aspect N NE E SE S SW W NW

Surface Soil loam Depth to Rock 7m Colour Dk Brown

Exposed Rock type granite % surface cover 0.5% size range 60cm

Litter 80% % surface cover Depth (cm) 10-20cm

Bare Ground 0 % surface cover

Adapted from 'Bushland Plant Survey' written by B. Keighery (1994) and published by Wildflower Society of WA (Inc). PO Box 64 Nedlands WA 6008

Panaivama
Postings 406988
Uth: 6418350


Dk Br-Bk (peat)


BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Mulr)- use pencil only

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class 2-10% 10-30% 30 - 70% over 70%

TREES				MALLEES	
	over 15m	5 - 15m	under 5m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)		30-70%			
DOMINANT SPECIES		<i>E. rudis</i>			
		<i>E. calophylla</i>			

SHRUBS			SHRUBS		
	over 2m	2m - 1.5m	1.5 - 1m	1 - 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)	30-70%				
DOMINANT SPECIES	<i>Agonis sp.</i>				
	<i>Typha floribunda</i>				
	<i>Acacia sp.</i>				
	<i>Acacia nemoralis</i>				

GRASSES		HERBS	SEDGES	over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)		10-30%	10-30%		
DOMINANT SPECIES		<i>Watsonia sp.</i>	<i>Gahnia sp.</i>		

4. VEGETATION CONDITION

Comments 2/5! V. poor condition. Heavily invaded by *Watsonia* + *Blackberry*. Remnant creekline vegetation with *E. rudis*.

lx

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESCENCE

SITE No SERP15

Date 8.12.89

Label each plant with plants number, site code, date and plant's name or working name if required

Record on Sheet

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

[illegible]

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Label each plant with plants number, site code, date and plant's name or working name if required

SITE No SERP 15

Record on Sheet

Date 8.12.96

NO RECORD

- Column 1 plant name
- Column 2 plant number
- Column 3 flowering time- TICK if species flowering
- Column 4 identification check

TREES	No	FI	ID	SHRUBS (cont.)	No	FI	ID	HERBS (cont.)	No	FI	ID
<i>Melaleuca rhaphinophylla</i>	✓	✓	✓								
<i>Eucalyptus rudis</i>	✓	✓	✓								
<i>Eucalyptus calophylla</i>	✓	✓	✓								
<i>Beyeria lechenaultii</i>	✓	✓	✓								
<i>Astavea aff. fascicularis</i>	✓	✓	✓	GRASSES							
<i>Grevillea manglii</i> sp. manglii	✓	✓	✓	<i>Ehrharta longiflora</i>	✓	✓	✓				
<i>Acacia saligna</i>	✓	✓	✓	<i>Hemarthria uncinata</i>	✓	✓	✓				
				<i>Microstachya stipoides</i>	✓	✓	✓				
<i>Tymonium flavibundum</i> flor	✓	✓	✓	? <i>Paspalidium</i> sp. sterile iddet	✓	✓	✓				
<i>Acacia urophylla</i>	✓	✓	✓	<i>Danthonia setacea</i>	✓	✓	✓	SEDGES			
<i>Agave linearifolia</i>	✓	✓	✓	<i>Briza maxima</i>	✓	✓	✓				
<i>Labeckia lanceolata</i>	✓	✓	✓	<i>Paspalum dilatatum</i>	✓	✓	✓				
<i>Lasianthum pterocarpum</i>	✓	✓	✓								
Blackberry = <i>Rubus</i> aff. <i>Selmeri</i>	✓	✓	✓	HERBS							
ADS: <i>Darwinia citriodora</i>	✓	✓	✓	<i>Gladiolus undulatus</i>	✓	✓	✓				
				<i>Xanthosia candida</i>	✓	✓	✓				
				<i>Coniza albida</i>	✓	✓	✓				
				<i>Lotelia alata</i>	✓	✓	○				
								<i>Baumea vaginalis</i>	✓	✓	✓
				<i>Sonchus oleraceus</i>	✓	✓	✓	<i>Baumea juncea</i>	✓	✓	✓
				<i>Stylidium calcaratum</i>	✓	✓	○	<i>Lepyrodia glauca</i>	✓	✓	✓
				<i>Trachymene pilosa</i>	✓	✓	○				
				<i>Watsonia</i> sp.	✓	✓	✓				
				IE <i>Scaevola</i> sp. - not <i>Lotelia</i>							
				<i>Anagallis arvensis</i>	✓	✓	✓				
				<i>Pelargonium australe</i>	✓	✓	✓				

<i>Ctenophorus reticulatus</i>	* *	*****		****			*			*					**		*
<i>Diplodactylus squarrosus</i>	*****	*****		****			**		*	**		*				**	
<i>Lerista uniduo</i>	*****	*****		****		***	****		**	*		*	*	**			
<i>Lerista aff. muelleri</i> B	*****	*****		****		*	*****		*			*					
<i>Ctenophorus scutulatus</i>		*****		**		*****	****										
<i>Lerista connivens</i>	***	*		*		*****	*		*	*							
<i>Heteronotia binoei</i>	*	**		***		****	*		*	*		*					
<i>Diplodactylus aff. pulcher</i> A	***			*		**	**		*	**		**			*		
<i>Menetia aff. greyii</i> A	*	**		***		*	**		**	**		*	*	*			
<i>Gehyra variegata</i>	*****	*****		*		*****	*****		*****	*****		*****			**		
<i>Lerista macropisthopus</i>	*****	*		***		*****	*****		*	*****					**		
<i>Diplodactylus strophurus</i>		**		*		**			*****	*****					**		
<i>Nephrurus levis</i>	*****	***		**		*****			*****	*****		***	*	*			
<i>Pogona minor</i>	**	*		**		**	***		*****	*****		*	***	***	***		
<i>Varanus eremius</i>	***			**		****	**		*****	*		*	****				
<i>Vermicella bertholdi</i>		***		***		***	*		**	*		*	****				
<i>Pygopus nigriceps</i>	*			*		*	*										
<i>Demansia calodera</i>				****					***	*		*					
<i>Moloch horridus</i>	*			***		*	*		*	***		*					
<i>Ctenotus mimetes</i>		*		*													
<i>Tiliqua occipitalis</i>				**													
<i>Neobatrachus sp</i> C				*			*										
<i>Tiliqua multifasciata</i>				*													