

Hi Tony,

Enclosed are data sheets from our sites. Are a few more plants to work out but they are not critical. Thought it probably more important you got these.

A few of the geology codes need checking. We did not have the correct legend to go with a couple of the geol. map sections and I can't locate the appropriate geology maps here. I've included the site locations on topo's, hope this will help. Give me a call if any queries.

If you have already entered the mammal data could I get a copy of it please?

Hope all's going well down there.

It's fairly cool here again, everyone walking around rugged up (low 20°)

Cheers,

A handwritten signature in dark ink, appearing to be 'T. M.' or similar, with a horizontal line above it.

20/7/2004

Kimberley Mammals -- Summary of Spotlight Effort

Team <u>AB, TH</u>	Location <u>MORNINGTON</u>
--------------------	----------------------------

Date <u>21/5/04</u>	Time started <u>1745</u>	Time finished <u>1850</u>	Total time <u>65 min</u>
Temperature <u>30</u>	Wind <u>NIL</u>	Cloud <u>NIL</u>	Moon <u>NIL</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>MRN #3</u>	<u>100</u>	<u>NIL</u>	

Date 22/5/04	Time started 1735	Time finished 1905	Total time 90
Temperature 30	Wind NIL	Cloud 100 50%	Moon NJ
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
MRA # 4	100%	NIL	

Date	24/5/04	Time started	1740 1700	Time finished	1910	Total time	20
Temperature	29	Wind	N/E	Cloud	50%	Moon	N/E
Trap line (name or similar habitat)		% total time		Mammals seen (Species and number)			
MRN #1		80%		N/E			
MRN #4		20%		N/E			

Date 26/5/04	Time started 1900	Time finished 2020	Total time 80
Temperature	Wind	Cloud	Moon
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
I6 4	20%	FELIS CATUS (2)	
I6 5	80%		
		Vehicle spotlight, along Mt Hatt Tr	
Saw 3 Boobies, 4 Frigatebirds, 3 Stone-curlews, 1 spotted nightjar			

INGLS
GAP

Date 27/5/04	Time started 1240	Time finished 1940	Total time
Temperature	Wind	Cloud	Moon
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
155	100%	Nk (2 Frognaths)	

Kimberley Mammals -- Summary of Spotlight Effort

Team <u>AB / TH</u>		Location	
Date <u>28/5/04</u>	Time started <u>1800</u>	Time finished <u>1910</u>	Total time <u>70</u>
Temperature	Wind <u>—</u>	Cloud <u>100%</u>	Moon <u>—</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>ILS / IBA</u>	<u>80</u>	<u>NIL</u>	
<u>IB4</u>	<u>20</u>	<u>NIL</u> <u>6 Boobies</u>	
		<u>1 Frogmouth</u>	
<u>vehicle with ably Mt Hart Rd</u>			

Date <u>31/5/04</u>	Time started <u>1750</u>	Time finished <u>1840</u>	Total time <u>50</u>
Temperature <u>cool</u>	Wind <u>NIL</u>	Cloud <u>100%</u>	Moon <u>—</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>MG3/1</u>	<u>100</u>	<u>P. brachyotis (2)</u>	

MT HART

Date <u>2/6/04</u>	Time started <u>1715</u>	Time finished <u>1830</u>	Total time <u>75</u>
Temperature <u>cool</u>	Wind <u>NIL</u>	Cloud <u>50%</u>	Moon <u>Full</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>Rebrachyot</u>			
<u>MG3/1</u>	<u>100</u>	<u>P. brachyotis (5)</u>	
		<u>Pseudochelone dahlia (1)</u>	

P.T.O.

Date <u>6/6/04</u>	Time started <u>1720</u>	Time finished <u>1910</u>	Total time <u>110</u>
Temperature <u>warm</u>	Wind <u>NIL</u>	Cloud <u>NIL</u>	Moon <u>NIL</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>MF 4 + 2</u>	<u>100</u>	<u>Petrogale brachyotis (1)</u>	
		<u>Pseudochelone dahlia (3)</u>	

MORGAN FALLS

P.T.O.

Date <u>7/6/04</u>	Time started <u>1725</u>	Time finished <u>1930</u>	Total time <u>125</u>
Temperature <u>warm</u>	Wind <u>NIL</u>	Cloud <u>NIL</u>	Moon <u>NIL</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>MF2</u>	<u>65%</u>	<u>P. brachyotis (3)</u>	
<u>MF4</u>	<u>10%</u>	<u>Felis catus (1), P. brachyotis (1)</u>	
<u>MF5</u>	<u>25%</u>	<u>NIL</u>	

1

2

3

Possum seen briefly moving towards shelter in rocks. Tricia had a good clear, but brief view & was sure it was *P. dahlia*. Andrew only saw very briefly & unable to identify.

4

Fairly group of 3 *P. dahlia* - 2 large one small. watched in full spot for 1755-1820. Then at 1755 was just emerged from shelter as first seen at back of tree. Then all climbed into *Planchonia*, then into *Ricinus virens* & commenced to feed. Seemed totally unphased by spotlight. Got good look at tail of two of the possums. Collected droppings at

5

this site later.

We first saw them about 60-70 m away, then approached to 40-50 m. Couldn't get closer as they were the other side of gorge & falls. Watched them with binoculars for duration of 25 mins

Kimberley Mammals -- Summary of Spotlight Effort

Team <u>AB/TH</u>	Location <u>MORGAN FALLS</u>
-------------------	------------------------------

Date <u>8/6/04</u>	Time started <u>1756</u>	Time finished <u>1910</u>	Total time <u>80</u>
Temperature <u>warm</u>	Wind <u>nil</u>	Cloud <u>nil</u>	Moon <u>nil</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>MF1</u>	<u>100</u>	<u>nil (3 Froemouths)</u>	
		<u>looking for gliders but 2 didn't see</u>	

Date <u>9/6/04</u>	Time started <u>1730</u>	Time finished <u>1900</u>	Total time <u>90</u>
Temperature <u>warm</u>	Wind <u>nil</u>	Cloud <u>nil</u>	Moon <u>nil</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>MF2</u>	<u>40%</u>	<u>P. brachyotis (1), P. dahl (3)</u>	
<u>MF4</u>	<u>40%</u>	<u>P. brachyotis (2)</u>	
<u>MF3</u>	<u>20%</u>	<u>Macropus robustus (1)</u>	

Date	Time started	Time finished	Total time
Temperature	Wind	Cloud	Moon
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	

Date	Time started	Time finished	Total time
Temperature	Wind	Cloud	Moon
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	

Date	Time started	Time finished	Total time
Temperature	Wind	Cloud	Moon
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	

Kimberley Mammals -- Traplines

Check code for Geol.

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location (Mt Hart Mts) Ynglis Gap Site		Trapline name and/o I	
Lat 17° 03' 01"	Date traps set 26/5/04	Cages 8	Big Ell 10	14	
Long 125° 08' 53"	Date traps lifted 30/5/04				
Topography and aspect Reasonably steep slope facing NE. Top of slope with large area of 4000 exposed black rock (in places forming fingers down slope).					
Rock type, nature and extent of exposure 25. Hart Dolomite Hill 2/3 of slope with 9000 exposed rock + 40-50% grass cover (rocks largely < 2cm) Black dolomite boulder area at top of slope - 100% exposed rock (largely 3 m) + no very exact record of exposed rock area					
Soil type, origin and depth Skeletal red brown loam.					Sample Y/N
VEGETATION (Muir system over page) L-Hr (except for Dolomite rock area without veg.)					
Main trees On slope (below regless black boulder) Euc. teretifolia + Euc. teretifolia greeniana					
Main Shrubs Slope - Greivillia Edge of boulder Greivillia retusifolia, Strachan.					
Main grasses Slope - Sorghum plumosum, Chrysopogon pallidus sp.					
Others Around edge of boulder + between boulder fingers dense vegetation (90%) as well as above included Greivillia brevifolia, Tinospora, Tardodroma, Melastroma amaranthum, Lonicera 2128, 2128					
Vines Melastroma oblongifolia, Albizzia lebbekii (juv), Buchnera sp., Asclepias x 2, Premna cumarata					
Main grasses form & height	hummock	Perennial tussock 0.5m	Perennial mats 1213	Annual	Brachyotum viridiflorum
FIRE					
When was last fire	1993	1992	< 3 yrs	3-10 yrs	> 10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Burnt Sept. 03 (T.A. Abbotts) charred but fairly vigorous growth. On slope numerous 1.2m Greivillias + 1m Euc. teretifolia. Base of Schima 2.5m Greivillias have been burnt but regrowth. Possibly 2 mts since fire on early burn + thick. lot of vegetative growth in 'valleys' between black boulder areas - possibly only 1 mts since fire. maybe wasn't burnt with rest.					
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) Tally Abbotts says this area was 'cooked' in Sept. last year but had green shoot a few weeks later.					
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Lower slope very few crevices (+ dense grass cover would mean hot fire) Black boulder area with large gaps + crevices - big enough for med. size mammals (however low veg. diversity + quantity)					
OTHER DISTURBERS (e.g. cattle, pigs, weeds) Cattle droppings (1 lot) seen 1/3 of the way up the slope. Hardly likely to be a problem due to ruggedness of slope.					
HABITAT CHANGE Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage					
PTO for more notes space		Photos Y/N		Photo re-takes Y/N	

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE d 70-100%	MID-DENSE c 30-70%	SPARSE i 10-30%	VERY SPARSE r 2-10%
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Birds

G.W. Cisticola
 R. Lorikeet
 R.W. Parrot
 Little Cuckoo Shrike
 W.W. Triller
 Double Bar Finch
 W. Wagtail
 Frogmouth
 Boobook
 Curlew
 Peaceful Dove
 Cuckoo
 Crimson Finch
 Mynah
 Pied Butcherbird
 Magpie
 Jackie Winter
 B.F. Cuckoo Shrike
 Peewee
 B.F. Woodswallow
 Galah
 Little Corella
 " Crow
 Variegated Lark
 Grey Butcherbird
 S.C. Thrushbird
 Ravenbird
 Kookaburra

Additional Notes

Rufous Whistler
 Long-tailed Finch
 Brown Falcon
 D. Faced Honeyeater
 Brown Songlark
 Rainbow Bird
 R. Lorikeet
 Fairy Martin
 Wedge tail eagle
 Night Heron
 Bar-shouldered Dove
 Brown Quail
 Little Woodswallow
 Great Egret
 W. Necked Heron
 Black cockatoo
 Bustard (off site)

Kimberley Mammals -- Traplines

Check
Geology
Code

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location 'Ingliss Gap Site'	Trapline name and/or code 2
--	--------------------------------	--------------------------------

Lat 17 06 54 125 10 06	Date traps set 26/5/04	Cages 8	Big Elliott 10	me 1
Long 17 06 49 125 10 07	Date traps lifted 30/5/04			

Topography and aspect Rugged sandstone (PSM-check geol map) range.
South facing slope.

Rock type, nature and extent of exposure Pgd p - lower slope with mainly smaller boulders, becoming large (5m+) boulders + slab rock higher up. Top of ridge rock.

Soil type, origin and depth Skeletal sandy loam amongst rocks
Sample ☒ Y/N

VEGETATION (Muir system over page) h A/c	Grows covers 60-70% on lower slope, up to 50% higher up where there is more exposed rock. GTC			
Main trees	Eucalyptus, Corymbia cadophora, Hakea orborens, Xanthostemon sp, Brachion. viridiflorus (= 212 ?), Erythrophloeum, Fern			
Main Shrubs	Grevillia pyramidalis, Cechlospermum fraseri			
Main grasses	Dwarf slope Thysan. bitextura then mainly Schima racemosa, Het. capensis, with Chrysopygon pallidus, Sorghum plumosum, Syntherisma bombycinus, Themeda triandra			
Others	Allysiacae, Vaginallis, Catalpa montana, Grevillea sp, Rhus sp, Erythroxylum			
Vines	Urosia cylindrica, Spermocoe 2145, Grewia reticulata, Crotalaria melleocarpa, Pterocaulon sp, Phytocora minor, Lycopodium lineari, Terminalia canescens			
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual (2217) Grewia, brom. Terminalia

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs?	>10 yrs
------	--------------------	------	------	---------	-----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) 2 yr? or older
Not much evidence of recent charring. Thysan. bitextura seeding. Grasses robust growth
1.7m Erythrophloeum chlorostachys + juv. Euc. Grevillia pyramidalis, Fern
Cechlospermum have been burnt previously but have regrown + all are approx. 1m tall.
Terminalia canescens regrow on burnt branches.

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)
Large boulders + rock areas provide shelter for small-mid size mammals at top of slope + also down slope in some places. (Further up slope from trapline shelter looks better).

OTHER DISTURBERS (e.g. cattle, pigs, weeds)
Passiflora foetida on trapline.

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	Photos ☒ Y/N
	Photo re-takes Y/N

Chickadee

Lat 17 06 35 125 10 43	Date traps set 26/5/04	Cages	Big Elliott	m
Long 17 06 46 125 10 43	Date traps lifted 30/5/04			

Rock type, nature and extent of exposure Scattered small rocks in vicinity of creek, grading into small (generally $< 2^{25}$) gravel between creek + range. CZS - (checked geol. map)

Sample ☒ Y ☐ N

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

1.5m *Euc. mirrata* + *Erythrophloeum* + *Im-Bachnanassa* Santalum + *TH2155* (2)
present over site. *Chrysogonum* has ~~cherry~~ charred bark but 2+ years growth,
since. Very dense *Heterocarpus*, + *TH2155* + sedge growth ^{along} creekline.

Acacias)) Only protection would be if dense growth along creek was too green to burn.

OTHER DISTURBERS (e.g cattle, pigs weeds)

Passiflora foetida + *Clitoria* present on site

Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

Photos ~~YIN~~

Photo re-takes Y/N

Both ends of trapline.

[illegible]

Kimberley Mammals -- Traplines

Check Geol. map.

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location 'Inglis Gap Site'	Trapline name and/or 5
--	-------------------------------	---------------------------

Lat 17° 08' 07" N 125° 09' 17" E	Date traps set 26/5/04	Cages 8	Big Ellic 10	14
Long 17° 02' 58" S 125° 09' 12" E	Date traps lifted 30/5/04			

+ 100m below.

Topography and aspect	Creekline.
-----------------------	------------

Rock type, nature and extent of exposure	Varied. Some slabby rock along creekline but not a lot. Atterbury
--	--

Soil type, origin and depth	Black clay soil. Alluvial.	Sample ☒ Y/N
-----------------------------	----------------------------	--

Mud of bank
G.D.

VEGETATION (Muir system over page) Varied. Some patches of closed canopy LAD. Other h.A.				
Main trees	Terminalia platyphylla, Lophocarpus, Serbania, also Planchonina, Pandanus sp.			
Main Shrubs	Patches of 1.2m fruty Thespesia thersites.			
Main grasses	Heteropogon, Dic. panicum, Arundinella nepalensis (also Mnesithea rotundifolia)			
Others	Sida (2/62), Abolmoschus, Erythrophloeum elongatum, Erythrina, resp. lig.			
Vines	Sida (2/62), Corchorus, also Buchnera, Serbania canabina.			
Main grasses form & height	hummocky	Perennial tussock	Perennial mats	Annual Cyperus vaginatus (2/62)

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)	Was this burnt Sept. 03? Could Thespesia get to fruit in one year? Given veg density
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))	Old Serbania along creek, badly affected by fire, some killed others have regrow. Patch where Serbania has previously extended from back from creek line (one now in poor condition) also big clump of Pandanus in this patch. Quite a lot of juv (1m) Serbania. (also) Terminalia platyphylla in creekline at one spot; also scattered along creek bank.

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

OTHER DISTURBERS (e.g. cattle, pigs, weeds)	Clitoria on site. (Check Chas. Corchorus sp.) Corchorus alitorius.
---	--

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	Photos ☒ Y/N
	Photo re-takes Y/N

Blistering on Euc. sp.
Thespesia thersites, Ficus vaceolata + Timonius timonius

End 16° 47' 00" Kimberley Mammals -- Traplines
124° 55' 18"

Team	Location	Trapline name and/or code
(AB/TH) (NM/JR) (TS/AT) (CP +)	Matthew George site.	1+3

Start 16 47 11	Date traps set 30/5/04	Cages	Big Elliott	med Elliott
Lat. 124 55 16				
End	Date traps lifted 3/6/04	8	10	14
Long				

Topography and aspect Topography runs along each side of rugged dolomite gorge (1 on east, 3 on west). Sides of gorge quite steep but all very slopes (with some minimal patches of vegetation - more than high Cap Site). The N. N. W. riparian strip of quite diverse species, particularly on west side, patchily along water edge.

Rock type, nature and extent of exposure ~~Hard Bolerik (Rich on Geology map)~~. Dark brown sandy loam (more sandy in some places, more loamy in others). Alluvial.

Soil type, origin and depth *As hard Dolomite (path on Geology map). limited soil in places along riparian strip, but above this boundary area of 100%*

Sample *YN*

VEGETATION (Muir system over page) LAc (narrow riparian strip)

Main trees *Melaleuca leucadendron*, *Lophystemon*, *Timonius timon*, *Subekhin*, *Pernambuco*, *Bauhinia*, *Celtis*, *Pouteria*, *sericea* (also *Grevillea* *intermedia*), *Acacia* *newberrana*, *Albizia* *lebbek*

Main Shrubs *Strachan lucida*, *Tabernaemontana*, *Ervatamia*, *orientalis* (TH2167) ✓

Main grasses *Ischaemum guineense* more within wet beds; & some *Echinochloa polystachya* present. Also *Eriochloa polystachya* & *Brachiaria repens* at start of TLP.

Others *Ficus opposita*, *Cochlospermum koseri* (not straight limb), *Melia*, *Babul*, *Tinospora*, *Albizia heterophylla*, *Croton*, *Claoxylon*, *Urtica*, *Prasia plectocarpa*, *Helicteres* spp.

Vines Buchananania, Ficus ramosa, Ficus hispida (Fr 2169), Planchonia, Carallia, Greuterella, Miconia, Marantaceae, Ipomoea (photo). Glucosmos to Polrato. Jodinium didymum

Main grasses form & height	hummock	Perennial tussock 1.5-2m 1.5-2m	Perennial mats	Annual Nephrolepis Brachystachya
-------------------------------	---------	---------------------------------------	----------------	--

1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398</
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--------

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Burnt Sept. 03! came down over Mt. Mathew - Tolly Roberts. Intensity of fire in gorge limited by lack of undergrowth + rocky nature of gorge. Churn + numerous trees + some big m. Leucodermis have been killed by fire. ~~the~~ Ischaemum rustical resprouted from burnt base. Start of TH - entrance to gorge has carried more fire - m...

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) Rubbley rock has protected some *Acacia namao* from fire.

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

OTHER DISTURBERS (e.g cattle, pigs weeds)

Passiflora on trellises.

HABITAT CHANGE Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest thickens

shrinkage	收缩率
-----------	-----

PTO for more notes space	Photos ☒ Y/N	Photo re-takes Y/N
--------------------------	--	--------------------

[illegible]

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE i 10-30%	VERY SPARSE r 2-10%
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Birds
 Bush shouldered Dove
 Peaceful Dove
 W. G. Rock Pigeon
 W. Gaped Honeyeater
 Bowerbird
 Leaden? Flycatcher
 Little Corella
 Little Woodswallow
 Tree Creeper
 Rainbow Lorikeet
 B. F. Cuckoo Shrike
 W. Wagtail
 Double bar Finch
 Little Cuckoo Shrike
 Grey Shrike Thrush
 Grey Butcherbird
 Pied "
 Brown Chantrel
 S. C. Cockatoo
 R. W. Parrot
 Restless Flycatcher
 Crimson Finch
 Tree Creeper
 Rufous Whistler
 Kookaburra
 Miners
 War. Wren

Azure Kingfisher
 S. C. Thrushbird

Kimberley Mammals -- Traplines

Clark
Geology

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Matthew Gorge site	Trapline name and/or code 2
--	--------------------------------	--------------------------------

Start 16 47 12 Lat 124 54 59	Date traps set 31/5/04	Cages 8	Big Elliott 10	m 1
End 16 47 01 Long 124 54 59	Date traps lifted 4/6/04			

Topography and aspect Trapline runs along slope parallel to small rocky creek, then drops down into creekline + up through 'gorgy' area.

Rock type, nature and extent of exposure Sandstone. Slope 80-90% (Pentecost Sandstone? check)
Creekline 90%+

Soil type, origin and depth Slope - dark brown clayey loam.

Sample Y/N

VEGETATION (Muir system over page) L B c, G T d (on slope - persistence on creek banks)

Main trees Slope - Euc. tectifica, Corymbia cadophora
Creekline - Lycopodium, Buchanania obliquata, Acacia pleurocarpa, Celtis, Vitis glaberrima/acuminata
Main Shrubs Slope - Stachyris lucida, Grewia breviflora, Ealytrix extipulata + Eriahne sulcata
Main grasses West end of trapline - Scribneria nemoralis
East end of trapline + ridge above creek/gorge - Sorghum stiaoidem with some Met. contorta
Others Ficus opposita, Ficus, Eriahne saligna, Dichiptera, Cochlospermum fraxi (not straight line)
Creekline - Adenium heterophyllum, Cynratia trifoliata, Passiflora, Cymbopogon? dependens
Vines Timonium timon, Aeschynomene aspera, Owenia venusta (juv)

Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual
		Scribneria nemoralis 1m	Met. contorta 1.2	Scribneria nemoralis 1.7m

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Burnt Sept. last year (2003) (Haffy Abbotts)
Grewias have reshot from burnt bases
Corymbia cadophora charred to 3m - shot from epicormic buds.
Juv. Euc., Buchanania, Eriahne saligna - reshot (after fire).

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Bouldery area on slope would provide small amount of protection for small mammals as would bouldery sections of creekline
Some but limited fire shelter in gorge walls for medium size mammals.

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

Passiflora present on site.

HABITAT CHANGE

Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N

Photo re-takes Y/N

Acum
~~Este~~ valida
?

~~Este~~ subiect

~~12. Substantive~~

5

posed rock
5 cm across

1. barium
 2. night
 3. the
 4. the

>10 yrs	
1/2003 but app	

g most

--	--

Kimberley Mammals -- Traps

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Mothan Gorge Site		Trapline name and/or code 5					
Lat 16°47'13"N 124°55'16"W		Date traps set 31/5/04		Cages 8		Big Elliott 10		med Elliott 14	
Long 16°47'27"N 124°55'15"W		Date traps lifted 4/6/04							
Topography and aspect Trapline lies along top of bank of Barker River (parallel with river on west side) then crosses rocky floodout riverbed.									
Rock type, nature and extent of exposure Hard dolomite. Banks 10-40% exposed rock - rocks < 25cm. Riverbed: 90% exposed boulders 50-70% + 90% exposed rock (varies)									
Soil type, origin and depth Dark brown sandy loam / loam - alluvial									
Sample Y/N									
VEGETATION (Muir system over page) LAc. Gtd (on bank) GLr in floodout/creekbed.									
Main trees <i>Eucalyptus</i> , <i>Albizia</i> , <i>Pisonia</i> , <i>Brachycheilus diversifolia</i> , <i>Lophocarpus</i> , <i>Bauhinia</i> , <i>Melaleuca</i> , <i>Casuarina</i> , <i>Leucaena</i> , <i>Acacia</i> , <i>Sonneratia</i> , <i>Portulaca</i> , <i>Memecylon</i> .									
Main Shrubs <i>Tephrosia</i> , <i>Zizania</i> , <i>Grewia</i> , <i>Reticularia</i> , <i>Hibiscus</i> , <i>Microcarpha</i> , <i>Portulaca</i> , <i>Memecylon</i> .									
Main grasses <i>Sorghum</i> , <i>Stipagrostis</i> , <i>Heteropogon</i> , <i>Chloris</i> , <i>Setaria</i> , <i>Syntherisma</i> , <i>Themar</i> .									
Others <i>Ficus</i> , <i>Artocarpus</i> , <i>Clusia</i> , <i>Planchonella</i> , <i>Boerhaavia</i> , <i>Ipomoea</i> , <i>Convolvulus</i> , <i>Centropogon</i> , <i>Psychotria</i> , <i>Passiflora</i> , <i>Maranta</i> , <i>Azadirachta</i> , <i>Albizia</i> , <i>Leucaena</i> , <i>Acacia</i> , <i>Sonneratia</i> , <i>Portulaca</i> , <i>Memecylon</i> .									
Vines <i>Boerhaavia</i> , <i>Artocarpus</i> , <i>Clusia</i> , <i>Planchonella</i> , <i>Boerhaavia</i> , <i>Ipomoea</i> , <i>Convolvulus</i> , <i>Centropogon</i> , <i>Psychotria</i> , <i>Passiflora</i> , <i>Maranta</i> , <i>Azadirachta</i> , <i>Albizia</i> , <i>Leucaena</i> , <i>Acacia</i> , <i>Sonneratia</i> , <i>Portulaca</i> , <i>Memecylon</i> .									
Main grasses form & height hummock Perennial tussock Annual <i>Sorghum</i> 1-7m Others to 185m Perennial mats <i>sulcata</i> Annual <i>Albizia procera</i>									
FIRE When was last fire 1993 1992 < 3 yrs 3-10 yrs > 10 yrs									
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Burnt Sept. 2003. (Tally Abbot Charming on Sonneratia + Melaleuca in creek bed (top m). Some Euc. sp. have fallen, others Fire may have swept around a bit. <i>Atalaya hanceana</i> , <i>Santalum lanceolatum</i> , <i>Planch. repens</i> from burnt base as have juv. <i>Terminalia caribaea</i> , <i>Bauhinia</i> + <i>Melia</i> .									
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) 1 x 2m Cypress pine on river bank									
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Rocky river bed would offer some protection for small mammals									
OTHER DISTURBERS (e.g. cattle, pigs, weeds) Evidence of cattle along trapline. <i>Passiflora pruriens</i> (more so in creek riverbed) <i>Euphorbia hirta</i>									
HABITAT CHANGE Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage									
PTO for more notes space									
Photos Y/N Photo re-takes Y/N									

Kimberley Mammals -- Traplines

Camp: 15° 01' 42", 126° 39' 51"

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Morgan Falls	Trapline name and/or code 1
--	--------------------------	--------------------------------

Start Lat 15° 01' 44" End Lat 15° 01' 59" Long 126° 39' 45"	Date traps set 6/6/04 Date traps lifted 10/6/04	Cages 8	Big Elliott 10	med Elliott 14
---	--	------------	-------------------	-------------------

Topography and aspect	Flat (woodland)
-----------------------	-----------------

Rock type, nature and extent of exposure	Beginning of Trapline laterite with occasional sandstone. By end of trapline is largely sand with occasional visible sandstone
--	--

Soil type, origin and depth	Brown loamy sand with laterite pebbles to start then pale red-brown sand	Sample Y/N
-----------------------------	---	------------

VEGETATION (Muir system over page) LAr, GTi + Hi				
Main trees	Euc. tetradonta mainly with occasional Euc. minor (in blossom), Ter. ? (latip), Erythrophloeum			
Main Shrubs	Petalostigma pubescens, Bossiaea bossiaeioides, Grevillea argentea, Pteritocarpus (2215), Callitris (2215), archaeta			
Main grasses	Sorghum stipoides, Tricaria sp. (probably T. bitextura - sterile)			
Others	Planchonia Duchingii, ananias, Persoonia tomentosa, Grevillea robusta, Wrightia saligna, Acacia -			
Vines	Polygona, Crotalaria, Grevillea, strobilifera, Denhamia obscura, Gossypium populifolium.			
Main grasses form & height	hummock < 5m Tricaria sp. bitextura	Perennial tussock	Perennial mats	Annual Seed heads to 2m Sorghum stipoides

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)	for 2 'web' age. 2002 → 2003. Tricaria rehot from burnt base + not very 'strong' plants Crotalaria to 50m tall - rehot from base, Im Bossiaea, Petalostigma Im responded from base. Charring on base of older Petalostigma's & 105m Ter. ? latip rehot from base. Fire may have been patchy some areas with denser grass + more litter. Has others with very little litter. Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) Several Callitris at end of site (also a few dead Eucyren stages) no juv. Callitris seen
--	---

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)	No real shelter on site from fire.
--	------------------------------------

OTHER DISTURBERS (e.g. cattle, pigs, weeds)	Evidence of cattle on site
---	----------------------------

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	Photos Y/N
	Photo re-takes Y/N

Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Birds

Additional Notes

Red tailed Black
 Crimson Finch
 W. Crested Honeyeater
 Brown Honeyeater
 Crow little
 Brush cuckoo
 Little Corella
 Rainbow bird
 D.B. finches
 Bowerbird
 Broad Cuckoo Shrike
 St. Friarbird
 Black bittern
 W. Quilled Rock pigeon
 Striped Pardalote
 Bar shouldered Dove
 Brown goshawk
 Wood thrush
 Red winged Parrot
 Rainbow lorikeet
 Cooey
 Mudlark
 Reston bluetongue

? Red Goshawk
 Little Cuckoo shrike
 Brown Owl
 Fairy tern
 Kestrel
 Banded Honeyeater
 White fronted Honeyeater
 White lined? Honeyeater
 Mistake bird

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Morgan Falls		Trapline name and/or code 2									
Start 15° 01' 41" Lat 126° 39' 52"		Date traps set 6/6/04		Cages 8		Big Elliott 10		med Elliott 14					
End 15° 01' 49" Long 126° 39' 59"		Date traps lifted 10/6/04											
Topography and aspect		Trapline runs close to edge of rugged KL Sandstone Crags - Flippers westward exposed											
Rock type, nature and extent of exposure		Most of Trapline 90-100% exposed rock (end of site gorge wall becomes slope + woodland but still quite rocky - 80% exposed.) King Leopold S											
Soil type, origin and depth		Brown loamy sand (skeletal) with quite a bit of humus in place											
		Sample Y/N											
VEGETATION (Muir system over page)		Patchy L-B/L-B; in places kind of site L-B/L-B and G-L.											
Main trees		Ficus (2189), Eucalyptus, (Alep Pouteria sericea, Rhytidophloeum chloro-											
Main Shrubs		Duchanania chinensis, Duchanania venicosa, Vanthostemon parviflorus, (2172)											
Main grasses		Triodia (2206) in rocky/gorge area. Sorghum stipoides towards end of site											
Others		Triumphetta brachystylis, Convolvulus, Planchonia, Dioscorea transversa											
Vines		Adiantum heterophyllum, Tinospora, Thunbergia, Parsonia velutina, Ichneumon											
Main grasses form & height		(hummock) Triodia 0.7m		Perennial tussock		Perennial mats		(Annual) Sor. stipoides 1.02m					
FIRE		When was last fire		1993		1992		< 3 yrs		3-10 yrs		> 10 yrs	
Comment (e.g. indicators used to assess age, longer term effects, intensity etc)		At least 1 'wet' possibly more. Not a lot of charring evident on trees. A few dead trees possibly killed by fire impact 1.5m Duchanania and Bossia. In Oenocarpus vinnosus responding at 1.5m. Rocky nature of area has protected some veg from fire.											
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))													
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)		Rocks + gorge face offer good crevices + shelter for mammals (+ veg.)											
OTHER DISTURBERS (e.g. cattle, pigs weeds)													
HABITAT CHANGE		Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage											
PTO for more notes space		Photos Y/N						Photo re-takes Y/N					

Stachys,
Syntherisma,
Capparis
pubescens,
Beady white tridentata. (2176),
Acaia humifusa (2184)
Capparis jacobii (2182)

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Morgan Falls	Trapline name and/or code 3
--	--------------------------	--------------------------------

Start 150° 01' 44" Lat 126° 39' 50"	Date traps set 6/6/04	Cages 8	Big Elliott 10	med Elliott 14
End 150° 01' 47" Long 126° 39' 36"	Date traps lifted 10/6/04			

Topography and aspect Trapline follows south side of Palmcondora Cr. Very beginning of site rocky slab creek bed, then approx. 100 m of damp grassland. Becomes rocky creek bed again mixed with damp grassland areas. (Creek is flowing permanent pool with some braiding).

Rock type, nature and extent of exposure King Leopold Sandstone - Rocky sections of Trophe creekline 80% exposed rock. Grassy low lying areas - almost no rock.

Soil type, origin and depth Dark brown clay/loam (except for rocky creekline which is coarse riverine sand.) Sample ☒ Y/N

VEGETATION (Muir system over page) Very patchy. Edge of creekline associated with grassland over, has narrow band of 2-5m Pandanus almost continuous creek + some emergent (12m) Mel. - Leucodendron - 15m tall. Patches of rocky creek (gravel) has mainly immature trees to 2-5m or patchy scrub - mainly Calytrix.

Main trees Mel. 2 windflowers, Calytrix ad brownii

Main Shrubs Germainia frutescens, Sargham plumosum, Metopogon contractus, Botriochloa sp. Eriachne sp.

Main grasses Muesithea rattusoides, Annual Sargham, Bruniella nipaensis, Pseudopogon sp.

Others Acacia plectocarpa, Buchanania decurva, Euc. tetradonta, Grevillea pteridifolia, Grevillea reginae, Terminalia platyphyla (juv.), Acacia divaricata, D. indica, Utricularia sp, sedges.

Vines Acacia holosericea, Antidesma ghaesambola

Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual
		Cler. trams, + Met. con. 1-2m		

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
-------------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) 2 years plus since fire? Green grass, low lying area would discourage low intensity fire. Quite a few, < 30cm, Euc. + Melaleuca seedlings (growing in grassland area). Very first section of site 1-3m Calytrix and Pandanus. West side of trapline Pandanus have regrow to 2m (after fire) - also 3m Mel. 2 windflowers common here. Drier slope above trapline has more than 1 year growth + some litter.

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Rocky creekline + rocky area in drier country above trapline would offer some fire shelter for small mammals. As would greenness of veg. (discouraging fire).

OTHER DISTURBERS (e.g. cattle, pigs weeds) Some grass areas flattened by cattle. Some grazing of area by cattle.

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	Photos ☒ Y/N Photo re-takes Y/N

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Morgan Falls	Trapline name and/or code L+
--	--------------------------	---------------------------------

Start 15 01 38 Lat 12 0 39 54	Date traps set 6/6/04	Cages 8	Big Elliott 10	med Elliott 14
End 15 01 38 Long 12 0 40 03	Date traps lifted 10/6/04			

Topography and aspect King Leopold Scudston Edge of rugged rocky gorge. First section a bit flatter prior to waterfall, then trapline follows the edge of the gorge.

Rock type, nature and extent of exposure King Leopold scudston. First section 95% exposed rock (big 'slabby' areas). Becomes boulder stacks + slabby rock, around edge of gorge. ~ 80% exposed rock.

Soil type, origin and depth Skeletal loamy sand (with quite a bit of humus in places)

Sample ☒ Y/N

VEGETATION (Muir system over page) Patches of L-B d/c then tree cover largely gives way to shrubs then

Main trees Ficus (22) Term? latipes, Litor glabra, acuminata back to patches of L-Bc (occasional encrup 75m)

Main Shrubs Styphnolobos lucida, Calytrix brownii, Grevillea agnifolia

Main grasses Triodia (2206) on rocky area (Sarcostyphnum intruding from above in places).

Others Stenandrium thurifolia, Sarcostemma australe, Brachychiton tridentatus, Acacia dunnii, Xanthorrhoea

Vines Tinospora, Cissampelos? (monanth), Triodia Garden

Main grasses hummock Perennial tussock Perennial mats Annual

form & height triodia - 7m

Acacia deltoidea subsp. deltoidea

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Way along gorge line is has quite a bit of protection from fire by rock slabs. Appears to be more leaf litter + denser green cover in woodland above edge of gorge (possibly 2 years since fire? - Cordonia responded to 1-2m after fire).

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Calytrix on edge of gorge toward end of trapline trunk has charred but tree looks healthy (we also a couple of dead Calytrix 'stags' on ground in same area).

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

'Slabby' bouldery nature of gorge offers good fire protection for mammals (+ veg)

OTHER DISTURBERS (e.g cattle, pigs weeds)

HABITAT CHANGE Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos ☒ Y/N

Photo re-takes Y/N

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Morgan Falls	Trapline name and/or code 5
--	--------------------------	--------------------------------

End Lat 15° 01' 20" N Start Lat 12° 39' 52" S	Date traps set 6/6/04	Cages 2	Big Elliott 10	med Elliott 14
End Long 115° 01' 34" E Start Long 126° 39' 52" E	Date traps lifted 10/6/04			

Topography and aspect	Flat (woodland)
-----------------------	-----------------

Rock type, nature and extent of exposure	Laterite at beginning of site becoming sandstone toward end
--	---

Soil type, origin and depth	Beginning laterite brown sand with laterite pebbles becoming pale red-brown sand.	Sample Y/N
-----------------------------	---	------------

VEGETATION (Muir system over page) LRA GTC + Hr.			
Main trees	<i>Euc. tetradonta</i> , <i>Euc. sp.</i> , <i>Acacia</i> , <i>Banksia</i> , <i>Callitris</i> , <i>Allocasuarina</i> , <i>Podocarpus</i> , <i>Myrtus</i> , <i>Leptospermum</i> , <i>Prostanthera</i> , <i>Leptosiphon</i>		
Main Shrubs	<i>Callitris</i> , <i>Allocasuarina</i> , <i>Podocarpus</i> , <i>Myrtus</i> , <i>Leptospermum</i> , <i>Prostanthera</i> , <i>Leptosiphon</i>		
Main grasses	Annual <i>Sorghum</i> , <i>Tridax</i> sp.		
Others			
Vines			
Main grasses form & height	Hummock <i>Tridax</i> sp. 0.5m	Perennial tussock	Perennial mats <i>Annual</i> <i>Sor. stridoidum</i> 1.7m tall

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)	Sparse growth of <i>Tridax</i> + Annual <i>Sorghum</i>. <i>Tridax</i> rehot from burnt base. <i>Tridax</i> suggests only one season since burn. However, are patches of excellent juv. <i>Euc</i> (1-3m) not burnt at base. Suggests last fire did not have a lot of fuel or was not very intense. Charring to 6m on <i>Euc</i>. <i>Petalostigma pubescens</i> rehotting. Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) A number of large (6-12m) Cypress in 2nd half of site but no medium size or juvenile (a lot of open spaces on fire).
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)	
No real shelter from fire.	

OTHER DISTURBERS (e.g. cattle, pigs, weeds)	Evidence of cattle on site
---	----------------------------

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	Photos Y/N
	Photo re-takes Y/N

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Mornington	Trapline name and/or code I
--	------------------------	--------------------------------

Lat 17° 33' 57" S Long 125° 59' 57" E	Date traps set 21/5/04	Cages 8	Big Elliott 10	med Elliott 14
Lat 17° 33' 54" S Long 125° 59' 55" E	Date traps lifted 25/5/04			

Topography and aspect Gorge heading nth. ~~east~~ into south facing range. Trapline runs along basal edge of ~~SW~~ then ~~SW~~ facing wall then crosses creek + heads back along basal edge of ~~SW~~ facing range. ~~At end of gorge, road with some vegetation.~~

Rock type, nature and extent of exposure King Leopold Sandstone gorge. All West facing wall steep highly exposed SW + east faces with some skeletal soils (above trapline). Creekline is delta formed in crevices. Gorge faces with some broken boulders - good for small mammals.

Soil type, origin and depth Alluvial loam with quite a bit of humus. (Shallow on slopes almost peat in valley floor). Sample ☒ Y/N

VEGETATION (Muir system over page) LAC at + patchy Qd-i in delta at mouth of gorge. Stripes, variable LAC in places.			
Main trees Delta - Euc. phyllorrhoea, Mel. leucodonta, Leucodonta sp., Timonius timon, Rigid above delta Teri. Kimberlyana , Brachychiton V. scutellaria , Buchanania dentata , Kimberlyana , Eucalyptus, Pandanus sp.			
Main Shrubs Delta - Ischnomimus austroale , Sermonia truncatula , Mimus , thea reticulata procerus			
Main grasses Rich poor scattered Cymbopogon sp + Tridax sp. Enneapogon purpurascens			
Others Floor of delta Scoparia cover - Ochradia austroale , Mib. hypellii (20%), schinoplectus microstus + mel cautibai sage (20%), 210 in pool in delta . 210 scattered sage			
Vines Others Trichostema regianum , Crotalaria verrucosa , Acacia glochitica (check name) Melaleuca viridis from above creek Brachia neuroranga in scyrium austroale			
Main grasses form & height	hummock	Perennial tussock	Perennial mats
		Delta - 1-5m	Annual
		Back faces 0.5m/1m	

FIRE	When was last fire	1993	1992	< 3 yrs	(3-10 yrs)	>10 yrs
------	--------------------	------	------	---------	-----------------------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Burnt left up (must have come over range) Charring on Melaleuca to 6m (Leucodonta charred whole of trunk) Dense grass cover in delta area + robust growth in other veg. in delta + gorge faces - fire appears to have not burnt some over, probably due to wetness or greening of veg. Lot of juv. ~~Buchanania~~ + ~~Mibiscus~~ on NW. facing slope (more intensive burn here)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Rocky gorge offers some fire protection, as does permanently wet delta area.

OTHER DISTURBERS (e.g. cattle, pigs, weeds) Old cattle dropping on floor of gorge (delta area) Donkeys seen close to gorge (juv + weaner) ~~Euphorbia~~ ~~hista~~ present (1 plant seen)

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	Photos ☒ Y/N
	Photo re-takes Y/N

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Birds

Brown Honeyeater
 Little Corella
 Pied Butcherbird
 Grey "
 Peaceful Dove
 Striated Pardalote
 Black-breasted buzzard
 Brown Goshawk
 Wedge-tailed Eagle - (breeding
 Fitz. Bluff
 off site)
 Kestrel
 Black Cockatoo
 Northern Rosella
 Red-wing parrot (feeding fledgling)
 Rainbow Lorikeet
 Little Corella
 Boobies
 Sulphur-crested Cockatoo
 Babbler
 Bearded Eater
 Silver-crowned Friarbird
 Yellow-billed Honeyeater

Additional Notes

Long-tailed Finch
 Doubtful
 Black-faced Woodswallow
 Little Woodswallow
 Rusten Flycatcher
 Willie Wagtail
 Mistake bird
 Little Crow
 Torresian Crow
 Grey Fantail
 White Fantail
 Kookaburra
 Singing bushlark
 Bowerbird
 W.W. Warbler
 Red-backed Wren
 B.F. Cuckoo Shrike
 Little "
 Diamond Dove
 W. Quilled Rock Pigeon (Fitz. Bluff
 off site)
 Weebill
 Whistly Kite
 Brown Quail
 Blue-faced Honeyeater
 Red-bellied Kingfisher
 Magpie

Kimberley Mammals -- Traplines

Team	Location	Trapline name and/or code
(AB/TH) (NM/JR) (TS/AT) (CP +)	Mammation site	2

(West)	Lat 17° 34' 07"	Date traps set	21/5/04	Cages	Big Elliott	med Elliott
(East)	Long 126° 59' 53"	Date traps lifted	24/5/04			
				8	10	14

Topography and aspect Trapline runs in straight line parallel to a stoney ridge (just below ridge). Sth. west aspect.

Rock type, nature and extent of exposure *Carsen volcanics. lot of small <20cm, loose rocks.
90% not covered by soil. ~~50%~~ 50% not covered by veg.*

Soil type, origin and depth	Skeletal sandy loam with lot of rock. Fluvial.	Sample (Y/N)
-----------------------------	--	--------------

VEGETATION (Muir system over page) LAR (<i>Tr. intermedia</i> + <i>Sch. nervosum</i> - approx 50% cover)				
Main trees <i>Euc. terminalis</i> , <i>Euc. brevifolia</i> , <i>Banksia cunninghamii</i>				
Main Shrubs <i>Crochlospermum fraseri</i> (glabrous long straight branch variety - s. ericksonii - 2/20)				
Main grasses <i>Triodia intermedia</i> (most of trapline), <i>Schima nervosum</i> on ridge just above (both) of trapline.				
Others <i>Gossypium australe</i> , <i>Banksia</i> ^{form <i>australis</i>} <i>sp.</i> <i>sp.</i> <i>Hakea carborescens</i> , <i>Eriodroma</i>				
Vines <i>ciliata</i> , <i>Sorghum plumosum</i> , <i>Flueggea virosa</i> , <i>Racoma</i> (2/20), <i>Rhynchospora minima</i> , <i>Triglochin</i>				
Main grasses form & height	hummock <i>Tr. int.</i> <i>Schima</i> - 1m	Perennial tussock SEMINER - 1m.	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
-------------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Robust *Trochiloides intermedius* hummingbirds.
Evidence of past fire - old charcoal to 4m up trunk. *Euc. breviflora* has sprouted from epicormic buds along branches (Branchlets 70cm long approx.)
Many *Cochlospermum* (to 3m) assume they would be burnt killed in fire - check.

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))
Georgium australe resprouting - assume after fire?

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)
Some larger rock areas on ridge would offer shelter for small mammals.

OTHER DISTURBERS (e.g cattle, pigs weeds)

Donkeys (Jenny & Fred) seen at start of Trampoline.

HABITAT CHANGE		Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space		Photos Y/N Photo re-takes Y/N

Kimberley Mammals -- Traplines

Team (AB/TH)(NM/JR) (TS/AT) (CP +)	Location Mornington	Trapline name and/or code Trapline 3
---------------------------------------	------------------------	---

Start Lat End 17° 34' 30" 44"	Date traps set 21/5/04	Cages 8	Big Elliott 10	med Elliott 14
Long 125° 59' 44" 53"	Date traps lifted 25/5/04			

Topography and aspect
Creekline along valley floor - pools some of the way + sometimes dry creekline.

Rock type, nature and extent of exposure (Rocks on levee scattered <10m; Rocky creekline)
Qa (alluvium) - in creekline - either side of volcanic (valley) ridges, either side of valley

Soil type, origin and depth
Alluvial heavy loam on levees. Some alluvial sand in creek Sample Y/N

VEGETATION (Muir system over page) L A c (along creek). L B c (surrounding country) + G T / G L i
Patches of Along creek Lophospermum, Boab + Bauhinia. Drier country surrounding creek -
Main trees Bauhinia, Acacia suberosa (occasional Eucalyptus) Wakea, Eucalyptus, Ficus, species to
Main Shrubs Carissa, Hib. = 2103, Melaleuca 2004, in creek: - Eriachne portulacae, Schima nervosum, Eriachne obtusa.
Main grasses Heteropogon contortus, Dicanthium aegyptium, Above levee: - Aristida
Others Sporobolus, 2004, 1.5m Melaleuca, Cyrtosperma, Tephrosia, Ficus (2003, 2004),
Vines Cordalana retusa, A. helioscopia, 2004, A. helioscopia, rufescens, Theophrasta, Bridelia, 2004,
Main grasses hummock! Perennial tussock 1m. Perennial mats Annual Melaleuca montana
form & height M. T. CDN, 1m. Dic. FRC, 1m ARI, ERI FES Acacia galeatricha

FIRE	When was last fire	2003 1993	2002 1992	< 3 yrs	3-10 yrs	> 10 yrs
------	--------------------	-----------	-----------	---------	----------	----------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)
Dead Acacia (bunt) ≈ 3m. Juv. unbunt Acacia: 0.7 - 1.5m. 1.5m Melaleuca sp.
Charring on Wakea suberosa, blistering on Boab trunks (3m)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) As above.

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)
In places fire has been stopped by creek, are 3m acacias on west side, although further north acacias on west side are burnt.

OTHER DISTURBERS (e.g. cattle, pigs, weeds)
Donkeys sighted + donkey tracks seen. Cattle tracks seen.
Some cattle traps on site and Lophospermum hirtum.

HABITAT CHANGE Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space	Photos Y/N	Photo re-takes Y/N
--------------------------	------------	--------------------

valley
H.S.
Teprosia DT 772
Pantest River
Acacia humida var. humida
Sida fibulifera

Kimberley Mammals -- Traps

Team	Location	Trapline name and/or code
(AB/TR) (NM/JR) (TS/AT) (CP +)	Mornington site	4

sth

(124)

Start Lat 17 34 28 125 59 48	Date traps set 21/5/04	Cages 8	Big Elliott 10	med Elliott 14
End Lat 17 34 16 125 59 49	Date traps lifted 25/5/04			

Topography and aspect Trapline runs along east side of nth/sth flowing creek.
Heavy rocky pools divided by ~~shallow~~^{broken} rocky areas

Rock type, nature and extent of exposure Carson Volcanic. loose rock largely $\approx$ 20cm in creekline

Soil type, origin and depth	Sandy loam (levee of creek). Alluvial.
	Sample ☒ Y ☐ N

[illegible]

FIRE	When was last fire	2004 1993	2002 1992	<u>< 3 yrs</u>	3-10 yrs	>10 yrs
-------------	--------------------	----------------------	----------------------	-------------------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) 2-3yr

~~3m Acacia coleii~~
1-2m Acacia coleii. Very little charring on old trees. (west side of creek with 3m Acacia coleii, ~~not~~ burnt > 3yr).

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) As above.

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Creek has stopped fire between east + west banks > 2 years ago. (limited amount of rocky/woody in creek between pools)

OTHER DISTURBERS (e.g cattle, pigs weeds)

None + Cattle evidence in area but not seen on site.

HABITAT CHANGE PTO for more notes space	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage	
	Photos ☒ Y/N	Photo re-takes Y/N

19150 = *Mimosa* affinis, *Crotalaria* sp. H, *Sporobolus* sp., *Crotalaria retusa*,
Indigofera tinctoria, *Indigofera tinctoria*, *Phaseolus vulgaris*, *Tridax* sp.,
~~*Desmodium*~~ *Desmodium* (sp.), *Leucaena* (sp.), *Eucalyptus* sp., *Rhus* *mespilifera*,
Desmodium *mespilifera*, *Leucaena* *mespilifera*, *Crotalaria* *mespilifera*, *Ficus* *mespilifera*,
Cassipouira *mespilifera*, *Bauhinia* *mespilifera* (2106)

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Morrington site	Trapline name and/or code 5
--	-----------------------------	--------------------------------

(NW) Stat 17 34 52 Lat 12 00 32	Date traps set 21/5/04	Cages 8	Big Elliott 10	med Elliott 14
(SE) End 17 35 02 Long 126 00 48	Date traps lifted 25/5/04			

Topography and aspect	Flat black-soil plain.
-----------------------	------------------------

Rock type, nature and extent of exposure	Carson Volcanic. Some areas with 10% exposed scattered rock; other areas less than this (rocks largely < 20cm).
--	---

Soil type, origin and depth	Cracking black-soil with some lightly cracking red-soil intrusions. Eluvial.
Sample	YN


VEGETATION (Muir system over page) LA, (90% grass cover over much of trapline - patches with < 30% reg. cover GT d (GTi))				
Main trees	Baobab + Euc. terminalis			
Main Shrubs	Carissa lanceolata (very scattered)			
Main grasses	Dianthus, Paspalum, Chrysopsis fallax, Aristida latifolia (some Panicum decompositum + Sahm. nervosum)			
Others	Flemingia pumila, Abutilon sp, Rhynchosia minima			
Vines				
Main grasses form & height	hummock	Perennial tussock Dieffenb. Chr. fol Aristida	Perennial mats	Annual


FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	> 10 yrs
------	--------------------	------	------	---------	----------	----------


Comment (e.g. indicators used to assess age, longer term effects, intensity etc)	Dense robust grass cover of mainly mature plants. Charring on Eucalypts + Baobabs several years old. Number of young (1m) Baobabs present.
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))	
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)	Cracks in self-mulching clay (planigale shelter)

OTHER DISTURBERS (e.g. cattle, pigs, weeds)	Some grazing evident (minimal) Cattle Cattle manure on site
---	---

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	Photos Y/N
	Photo re-takes Y/N







Morgan Falls

MF5

MF4

MF3

MF Camp

MF2


MF1

Falls

284-

Falls

50



Trap Layout for Mornington, Inglis Gap, Mathew Gorge, Morgan Falls.
Mammal Survey 20/5/2004 – 10/6/2004

2 x med. Elliot + 1 x lge. Elliot

30m

Cage trap

30m

2 x med. Elliot + 1 x lge. Elliot

30m

Cage trap

30m

1 x med. Elliot + 1 x lge. Elliot

30m

Cage trap

30m

1 x med. Elliot + 1 x lge. Elliot

30m

Cage trap

30m

1 x med. Elliot + 1 x lge. Elliot

30m

Cage trap

30m

1 x med. Elliot + 1 x lge. Elliot

30m

Cage trap

30m

1 x med. Elliot + 1 x lge. Elliot

30m

Cage trap

30m

1 x med. Elliot + 1 x lge. Elliot

30m

Cage trap

30m

2 x med. Elliot + 1 x lge. Elliot

30m

2 x med. Elliot + 1 x lge. Elliot

(30m as paced and according to terrain.)

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <u>MORNINGTON</u>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
1	<i>Rattus traneys</i>	22/5/04	LE	M 54839	Muscle DNA Museum DNA	145	140	172	M	A	v. large testes	MRN 1
2	<i>Zyromys argurus</i>	22/5	ME	M 54840	Muscle DNA Museum DNA	26	85-5	112	M	SA	Nil	MRN 3
3	<i>Equus asinus</i>	22/5	tracks									3, 4, 5
4	<i>Canis lupus</i>	22/5	tracks									3, 4, 5
5	? <i>M. robustus</i>	22/5	scats									4
6	<i>Z. argurus</i>	23/5	LE									1
7	"	"	ME									1
8	"	"	LE									1
9	"	"	LE	R	-	44	-	-	F	A	Nil	1
10	? <i>P. brachyotis</i>	"	scats	-	-	-						1

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)				Location <i>MORNINGTON (MRN) + INGIS GAP</i>								
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
11	<i>Z. argurus</i>	24/5/04	ME	R		47			♀	A	? pregnant	MRN 1
12	<i>R. tunneyi</i>	25/5	ME	R						SA	-	MRN 1
13	<i>R. tunneyi</i>	23/5	LE	R								MRN 3
14	<i>Equus asinus</i>	23/5	Obs									MRN 2
15	<i>Bos taurus</i>	23/5	tracks									MRN 1
16	<i>Bos taurus</i>	23/5	tracks									MRN 3
17	<i>Felis catus</i>	26/5	S	S								IG 4
18	<i>Z. argurus</i>	27/5	ME	M 54841	MURDOCH MUSEUM A3	33	98	101	♀	SA	Nil	IG 1
19	<i>R. tunneyi</i>	27/5	ME	M 54842	MURDOCH MUSEUM A4	20	135	105	♂	A	enlarged testes	IG 5
20	<i>Bos taurus</i>	27/5	tracks									IG 5

INGIS GAP
(IG)

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

INGIS GAP

Kimberley Mammals -- RECORDS

102 - 3
93 - 34 = 29
64 - 39 = 25
7

Team (AB/TH) (NM/JR) (TS/AT)					Location							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
1	R. tunneyi	27/5	ME	R		72			M	A	—	
2	"	1.	ME	R		65			F	A	—	
3	R. tunneyi	"	LE	R		49			F	SA		
4	"	"	ME	R		25			M	J		
5	"	"	ME	R		71			F	A	—	
6												
7												
8												
9												
0												

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Inglis Gap</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
21	<i>Bos taurus</i>	26/5/04	D	S								IG 3
22	<i>Rattus tunneyi</i>	28/5	ME	R		75g			♂	SA		IG 5
23	<i>Z. argurus</i>	28/5	ME	M 54843	WURDOCK MUSEUM	26	94	damaged	♂	J		IG 2
24	<i>Z. argurus</i>	28/5	ME	R		39			♀	A	? pregnant	IG 4
25	<i>Isodon macrourus</i>	28/5	C	R					♀	A	see over	IG 3
26	<i>Isodon macrourus</i>	28/5	C	R					♂	? A	see over	IG 3
27	<i>R. tunneyi</i>	28/5	ME	R						SA		IG 3
28	<i>Equus caballus</i> ?	28/5	amphib									IG 3
29	<i>R. tunneyi</i>	29/5	ME	R		72			♂ M	A	-	IG 5
30	<i>R. tunneyi</i>	29/5	ME	R		65			F	A	-	IG 5

escaped!
droppings on road

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
25	enlarged teats, clean moist pouch head length 88.5, per 64.5 digital photos taken, hair taken
6	head length 75mm, per 61.5mm digital photos taken, hair taken
7	
8	droppings ID'd by Tricia - too big to be donkey.
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>INGUIS GAP</i>							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
21	<i>R. tunneyi</i>	29/5/04	LE	R		49			F	SA	-	IG5
32	<i>R. tunneyi</i>	29/5	ME	R		25			M	J	-	IG5
33	<i>R. tunneyi</i>	29/5	ME	R		71			F	A	-	IG5
34	<i>Z. agurui</i>	29/5	ME	R		49			F	A	<i>inguis</i>	IG4
35	<i>R. tunneyi</i>	29/5	ME	R		92			M	A	<i>over</i>	IG3
36	<i>R. tunneyi</i>	29/5	ME	R		64			M	A	<i>over</i>	IG3
37	<i>Isodon macrourus</i>	29/5	cage	R	<i>Murdoch hair</i> <i>Murdoch ear</i>	1015			M	A		IG3
38	<i>R. tunneyi</i>	29/5	ME	R		58			M	SA?		IG3
39	<i>R. tunneyi</i>	29/5	LE	R		58			M	A		IG3
40	<i>? Zygomys</i>	29/5	ME	<i>M</i> <i>54344</i>	<i>Murdoch</i> <i>WAM</i>	28	87.5	<i>damaged</i>	F	SA?		IG3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
35	testes not enlarged
36	testes not enlarged
37	<i>I. macrurus</i> ♂ head length 83.5 pes 67 hair collected 2 photos taken hair in Murdoch DVA ear in Murdoch DVA
8	
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Inglesham</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
41	<i>Egus asinus</i>	29/5/04	heard									near IG3
42	<i>2. argurus</i>	29/5	ME	R		-			-	-		IG1
43	<i>2. argurus</i>	29/5	ME	R		40			M	A	enlarged testes	IG1
44	<i>2. argurus</i>	29/5	ME	R		33			F	SA?		IG1
45	<i>2. argurus</i>	30/5	ME	R					F	A	? pregnant	IG1
46	<i>2. argurus</i>	30/5	ME	R		6			M	SA		IG3
47	<i>R. tunneyi</i>	30/5	ME	R		68			M	A		IG3
48	<i>I. macrourus</i>	30/5	C	R	hair	1340			F	A	over	IG3
49	<i>R. tunneyi</i>	30/5	LE	R	43g 87				F	SA?		IG3
50	<i>R. tunneyi</i>	30/5	ME	R		84			M	A	enlarged testes	IG3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
6	
7	
48	I. macrourus clean moist pouch some nipples & enlarged fm previous young.
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Inghs Gap</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
51	<i>D. hallucatus</i>	31/5/04	LE	R		435			F	A	—	M61
52	<i>P. brachyotis</i>	31/5	D	S								M63
53	"	31/5	D	S								M63
54	<i>P. brachyotis</i>	31/5	S	S								M63
55	<i>Z. angustus</i>	1/6/04	ME	R		29			F	A		M62
56	<i>D. hallucatus</i>	1/6	LE	R		330			M	A		M62
57	<i>Z. woodwardi</i>	1/6	LE	^M 54346	<i>Murdoch WAM</i>	117			F	A	over	M63
58	<i>D. hallucatus</i>	1/6	ME	R		390			F	A	over	M63
59	<i>Z. woodwardi</i>	1/6	ME	R		101			F	A	not breeding	M63
60	<i>D. hallucatus</i>	1/6	ME	R		285			M	A		M63

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
6	
57	Large adult female, vagina not perforated.
58	Pouch dirty, ? previous year, not in reproductive condition at present.
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Moor Her (Mt Matthew Gorge)</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
b1	<i>Z. angustus</i>	1/6/04	ME	^m 54845	<i>Murchison</i> WAM	35			M	A	not breeding	M631
b2	<i>D. hallucatus</i>	1/6	ME	R		300			F	A	not breeding	M631
b3	<i>D. hallucatus</i>	1/6	ME	R		505			M	A		M61
b4	<i>D. hallucatus</i>	1/6	LE	R		295			F	A	n.p.d.	M81
b5	<i>Z. woodwardi</i>	1/6	ME	R		142			M	A	-	M61
b6	<i>Z. angustus</i>	1/6	ME	R		60			M	A	over	M64
b7	<i>R. lunneyi</i>	1/6	ME	R		80			M	A	over	M64
b8	<i>R. lunneyi</i>	1/6	ME	R		22			M	J	-	M64
b9	<i>R. lunneyi</i>	1/6	ME	R		70			M	A	-	M64
b10	<i>Bos taurus</i>	1/6	pat	S							over	M65

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

ADD to A
R. lunneyi
M65
LE 85g
O Adult
+ no breed

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
66	no testes development
67	Scrotum somewhat developed, testes small, shrink after breeding?
8	
9	
70	The largest single cattle droppings I have ever seen!

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>MT Mathew Gorge (Mt Hart)</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
71	<i>Bos taurus</i>	2/6/04	track	S								Mar M64
72	<i>Z. woodwardi</i>	2/6	LE	R	Reptile	147			F	A	non-bred	M63
73	<i>D. hallucatus</i>	2/6	LE	R	Reptile	380			F	A		M63
74	<i>D. hallucatus</i>	2/6	ME	R		335			M	A		M63
75	<i>D. hallucatus</i>	2/6	LE	R		395			F	A	over	M61
76	<i>Z. woodwardi</i>	2/6	ME	R		59			F	J		M64
77	<i>R. tunneyi</i>	2/6	ME	R		76			M	A		M64
78	"	2/6	ME	R		29			F	J		M64
79	"	2/6	ME	R		42			F	J		M64
80	"	2/6	LE	R		64			M	A	no dev.	M64

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
6	
75	large park, clean, no current breeding
8	
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>MT HART</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
81	<i>Z. angustus</i>	2/6/04	LE	R		39			F	A	? pregnant	M62
82	<i>D. hallucatus</i>	2/6	ME	R		290			F	A	n.p.d.	M62
83	<i>D. hallucatus</i>	2/6	LE	R	Reaphne	340			M	A	-	M62
84	<i>R. tunneyi</i>	2/6	LE	R		83			F	A	? pregnant	M64
85	<i>R. tunneyi</i>	2/6	ME	R		95			F	A	-	M64
86	<i>R. tunneyi</i>	2/6	ME	R		86			F	A	? pregnant	M64
87	<i>P. brachyotis</i>	2/6	scats									M62
88	<i>M. robustus</i>	2/6	scats									M62
89	<i>M. robustus</i>	2/6	scats									M645
90	<i>Canis lupus</i>	2/6	scats									M64

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

fairly common
not collected
too wet

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)				Location <i>MT HART</i>								
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
91	<i>P. brachyotis</i> x4	2/6/04	S	S								M61
92	<i>P. dahl</i>	2/6/04	S	S								M61
93	<i>Petaurus brevicaeps</i>	2/6	S	S								M65
94	<i>D. hallucatus</i>	3/6	C	R		325			F	A	n.p.d.	M62
95	<i>D. hallucatus</i>	8/6	ME	R		325			M	A		M62
96	<i>Z. argurus</i>	3/6	ME	R		38			M	A	no dev.	M62
97	<i>R. tunneyi</i>	3/6	ME	R		76			F	A	non breed	M65
98	<i>R. tunneyi</i>	3/6	ME	R		87			F	A	non breed	M65
99	<i>R. tunneyi</i>	3/6	ME	R		70			F	A	non breed	M64
100	<i>R. tunneyi</i>	3/6	ME	R		27			F	SA	—	M64

with Jiri in
his camp -
photographed

recapture?

~~recapture~~

recapture

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>MT Hart</i>							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
101	<i>R. tunneyi</i>	3/6/04	ME	R		341			F	SA		M64
102	<i>R. tunneyi</i>	3/6	LE	R		63			M	A	no dev	M64
103	<i>D. hallucatus</i>	3/6	ME	R		340			M	A		M64
104	<i>D. hallucatus</i>	4/6	C	R								M62
105	<i>D. hallucatus</i>	4/6	C	R								M62
106	<i>D. hallucatus</i>	4/6	C	R		465			M	A		M65
107	<i>D. hallucatus</i>	4/6	LE	R		375			F	A	npd	M65
108	<i>R. tunneyi</i>	4/6	ME	R		145			M	A	none	M65
109	<i>R. tunneyi</i>	4/6	ME	R		115			M	A?	none	M65
110	<i>R. tunneyi</i>	4/6	LE	R		100			F	A?	none	M65

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>MT HALL</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
111	<i>R. tunneyi</i>	4/6/04	ME	R		85			F	SA?	nuc	M65
112	<i>R. tunneyi</i>	4/6	ME	R		70			M	SA?	nuc	M64
113	<i>R. tunneyi</i>	4/6	ME	R		30			F	J	nuc	M64
114	<i>R. tunneyi</i>	4/6	ME	R		85			M	A	ent. testes	M64
115	<i>R. tunneyi</i>	4/6	ME	R		85			F	A	? pregnant	M64
116	<i>Z. argurus</i>	4/6	ME	R		50			F	A	? pregnant	M64
117	<i>Z. argurus</i>	4/6	ME	R		70			M	A	ent. testes	M64
118	<i>R. tunneyi</i>	4/6	ME	R		92			M	A	ent. testes	M63
119	<i>D. hallucatus</i>	4/6	LE	R		375			M	A	-	M63
120	<i>D. hallucatus</i>	4/6	ME	R		-			-	A		

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

very wet
- released immediately

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <u>MT HART</u>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
221	<i>Z. woodwardi</i>	4/6/04	LE	R		137			M	A	no dev.	MG3
122	<i>Z. woodwardi</i>	4/6	C	R		150			M	A	no dev	MG3
123	<i>D. hallucatus</i>	4/6	ME	R		280			F	A	n/a	MG3
124	<i>Bos taurus</i>	6/6	trails droppings									MF1
125	<i>Bos taurus</i>	6/6	"									MF3
126	"	7/6										MF2
127	"	7/6										MF5
128	<i>Canis lupus</i>	7/6	scats									MF2
129	<i>Hydromys</i>	7/6	scats									creek bed
130	<i>Leggiana</i>	7/6	LE	M 54847	Murdoch WAM	12	65	46	F	A	-	MF5

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

photo + collected

MT HART

NOBATA FACUS

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
6	
7	
8	
9	
130	hindfoot 14 mm ear length 10 mm

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <u>MORGAN FALLS</u>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
131	<i>Z. argurus</i>	7/6/04	LE	M 54848	Murdock WAM	35	91mm	109.5mm	F	A	not bred	MF4
132	<i>R. tunayi</i>	7/6	ME	M 54849	"	58	126mm	109mm	F	A	not bred	MF5
133	<i>R. tunayi</i>	7/6	ME	M 54850	"	44	111mm	100mm	M	A	not taken	MF5
134	<i>Naemops robustus</i>	2/6	scats									MF all
135	<i>Petrogale brachyotis</i>	6/6	D/S	S					F			MF4
136	<i>Pseudocheirus dahl</i>	6/6	S	S								MF2
137	<i>P. brachyotis</i>	6/6	S	S								MF2
138	<i>Felis later</i>	6/6	S	S								MF4
139	<i>P. brachyotis</i>	6/6	S	S								MF4
140	<i>P. brachyotis</i>	7/6	D	S								MF2

3 seen together
3 seen

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
136	3 possums (family group? one smaller than other two) observed with binoculars ~50m feeding in Ficus and unknown tree. Across falls from us.
7	
8	
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Morgan Falls</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
141	<i>2. argurus</i>	8/6/04	LE	R		51			F	A	? breeding	MF2
142	<i>R. tunneyi</i>	8/6	ME	R		57			F	A	over	MF3
143	<i>Hydromys</i>	8/6	scats									MF
144	<i>Bos taurus</i> (3)	8/6	D	S								MF3
145	<i>2. argurus</i>	9/6	ME	R		11			M	J	-	MF4
146	<i>2. argurus</i>	9/6	ME	R		54			F	A	breeding condition	MF4
147	<i>R. tunneyi</i>	9/6	ME	M 54851	WMSOCH WAM	15	78	69	M	J	-	MF3
148	<i>2. argurus</i>	10/6	ME	R		38			M	A	-	MF4
149	<i>R. tunneyi</i>	10/6	LE	R		93			M	A	ear tester	MF5
150	<i>R. tunneyi</i>	10/6	ME	R		14			F	J	-	MF3

in middle of creek on rock

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
142	R. unnei - Adult, perforated to ^{to} is ^{is} vagina ^{vagina} but apparently not currently breeding.
3	
4	
5	
6	
7	
8	
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Morgan Falls</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
151	<i>P. dahlia</i> (3)	9/6	S	S								MF2
152	<i>P. brachyotis</i> (1)	9/6	S	S								MF2
153	<i>P. brachyotis</i>	9/6	S	S								MF4
154	<i>P. brachyotis</i>	9/6	S	S								MF4
155	<i>M. robustus</i>	9/6	S	S								MF3
6												
7												
8												
9												
0												

same 3 as before.

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlit (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- Summary of Spotlight Effort

Team NM + TW	Location Oscar Range
--------------	----------------------

Date 22 May 04	Time started 1820	Time finished 2025	Total time 2.05
Temperature warm	Wind light	Cloud light	Moon nil
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
2	20	-	
#1	40	-	
5	40	Euro	

Oscar
Ra

Date 23/5/04	Time started 1745	Time finished 2015	Total time 2.5
Temperature warm	Wind light	Cloud full cover	Moon none visible
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
4	100	Dingo: 2+ heard walking down track beside range.	

hrs
Oscar
Ra

Date 24/5/04	Time started 1835	Time finished 2005	Total time 1.30
Temperature warm	Wind none	Cloud none	Moon last Y4
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
4		Euro	
5		Euro x 2	

Oscar
Ra

Date 23 2/6/04	Time started 1900	Time finished 2130	Total time 2.5 hrs
Temperature warm	Wind 0	Cloud 100%	Moon covered
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	

MT
GLADYS

Date 3/6/04	Time started 1830	Time finished 2030	Total time 2 hrs
Temperature warm	Wind nil	Cloud 100%	Moon covered
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
line 2	80	nil	
of line 1	20	nil	

MT
GLADYS

Oscar Σ 6 hrs
Mt Gladys Σ 4.5 hrs

Kimberley Mammals -- Summary of Spotlight Effort

Team NMCK + TW

Location Cheddar Cliffs + Barker Gorge Cheddar Cliffs

Date 27/5/04	Time started 1905	Time finished 2230	Total time 3.5 hrs
Temperature warm	Wind still	Cloud 0	Moon 1/2
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
1	20%	Agile Wallaby	
2	5%	nil	
3	50%	Rock Wallaby, Possum	
Riverine Woodl	25%	nil	

Date 7 June 04	Time started 1800	Time finished 2015	Total time 2 1/4 hrs
Temperature warm	Wind —	Cloud —	Moon —
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
2	100	Petrogale ? burbridgei M52713	

Date 8 June 04	Time started 1800	Time finished 2030	Total time 2 1/2 hrs
Temperature warm	Wind —	Cloud —	Moon —
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
5	5	bats only	
3	95	bats only	

Date 9 June 04	Time started 1725	Time finished 2000	Total time 2 3/4
Temperature warm	Wind 0	Cloud 0	Moon 0
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
3	100%	sighting quietly 2 head torches at 2 locations along creek —	
		nil seen or heard except bats	

Date	Time started	Time finished	Total time
Temperature	Wind	Cloud	Moon
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	

Kimberley Mammals -- Traplines

37

Team <u>TV</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>Oscar Range</u>	Trapline name and/or code <u>1</u>
--	-----------------------------	---------------------------------------

Lat <u>17° 38' 19.2" S</u>	Date traps set <u>21/5/04</u>	Cages <u>10</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>124° 58' 15.6" E</u>	Date traps lifted <u>24/5/04</u>			

Topography and aspect <u>Northern Aspect, very gentle slope + plain</u>
--

Rock type, nature and extent of exposure <u>Small Limestone rise + alluvial flat</u>

Soil type, origin and depth <u>1 clayey apron + deep alluvial deposit</u>	Sample <u>YN</u>
--	------------------

VEGETATION (Muir system over page) <u>to 7m</u>				
Main trees	<u>Corymbia dampieri, occ. Adansonia gregorii to 8m</u>			
Main Shrubs	<u>Grevillea parallela + Dolichandrone heterophylla</u>			
Main grasses	<u>Triodia pascoana - Sorghum plumosum - Dicanthium fecundum</u>			
Others	<u>Hybanthus auranriacus, Stenodia grossa, Gomphrena spp.</u>			
Vines	<u>nil</u>			
Main grasses form & height	hummock <u>Triodia 0.5m</u>	Perennial tussock <u>Dicanthium 1m</u>	Perennial mats	Annual <u>Sorghum 1.5m</u>

FIRE	When was last fire	<u>201993 2003</u>	<u>201992 2002</u>	<u>< 3 yrs</u>	3-10 yrs	>10 yrs
------	--------------------	-------------------------------	-------------------------------	-------------------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) <u>Size of Triodia hummocks</u>
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) <u>(none)</u>
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

OTHER DISTURBERS (e.g cattle, pigs weeds) <u>Cattle</u>
--

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	
Photos <u>YN</u> ✓	Photo re-takes Y/N

Photos 9+10 } stick 1
Photos 11+12 }

✓


Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass <u>Tall Grass</u> Low Grass Herbs	<u>Open Mat Plants Hummock Grass</u> Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Also Caeminiid Land Snails feeding on algal mat

Additional Notes

Co-dominants: *Triodia pascoana* / *Sorghum plumosum* / *Dicanthium fecundum*
with scattered *Corymbia dampieri* to 7m



Other species noted:
Sesbania simplicifolia
Crotalaria linifolia
Goodenia odonnellii
Mimulus uedalii
Trichodesma zeylanica
Stemodia grossa
Striga curviflora
Gomphrena brachystylis
Corchorus sp.
Chamaesyce coghanii

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) ^{Tw} (TS/AT) (CP +)		Location Oscar		Trapline name and/or code 2	
Lat 17° 38' 25.7" S	Date traps set 21/5/04	Cages	Big Elliott	med Elliott	
Long 124° 56' 51.7"	Date traps lifted 25/5/04	10	11	18	
Topography and aspect Karst gully + ridges (Massive Devonian Reef)					
Rock type, nature and extent of exposure Limestone ranges, little soil					
Soil type, origin and depth Limestone skeletal + scree + litter					
Sample ☒ Y ☐ N					
VEGETATION (Muir system over page)					
Main trees <i>Ficus platypoda</i> <i>Celtis philippensis</i> <i>Corymbia dampieri</i> <i>Vitex glabrata</i> <i>Lophostemon</i>					
Main Shrubs <i>Flueggea virosa</i> <i>Dodonaea</i> <i>Plumbago</i> <i>grandiflorus</i>					
Main grasses <i>Tridax</i> ? <i>pascoana</i> <i>platyptera</i> <i>zeylanica</i>					
Others					
Vines <i>Jasminum didymum</i> , <i>Tinospora smilacina</i> , <i>Cayratia cardiophylla</i>					
Main grasses form & height	hummock ✓ 1.5m	Perennial tussock nil	Perennial mats nil	Annual nil	
FIRE		When was last fire	1993 2003	1992 2002	< 3 yrs
					3-10 yrs
					>10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) No visible fire scars on trees Hummock grass clumps in inner valley (massive)					
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) Rainforest species (<i>Celtis philippensis</i> , <i>Vitex glabrata</i> in Karst)					
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Massive limestone outcrops & creekline					
OTHER DISTURBERS (e.g. cattle, pigs, weeds) None					
HABITAT CHANGE Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage					
PTO for more notes space		Photos ☒ Y ☐ N		Photo re-takes Y/N	


Photos 3+4 stick 1

Lophostemon valley
Muir vegetation System
Karst outcrops
Corymbia valley

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% t
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns	Dense Ferns	Ferns	Open Ferns	Very Open Ferns
	Mosses, liverwort	Dense Mosses	Mosses	Open Mosses	Very Open Mosses

Additional Notes

Trapline 2



Kimberley Mammals -- Traplines

Team <u>TW</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>Oscar Range</u>	Trapline name and/or code <u>3</u>
--	-----------------------------	------------------------------------

Lat <u>17° 37' 44.9"</u>	Date traps set <u>21/5/04</u>	Cages <u>7</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>124° 56' 38.4"</u>	Date traps lifted <u>25/5/04</u>			

Topography and aspect

Creekline meandering across plain

Rock type, nature and extent of exposure

Soil type, origin and depth

Black Soil Plain

Sample Y/N

VEGETATION (Muir system over page)

Main trees Euc. teretica / Corymbia dampieri / Terminalia platyphylla

Main Shrubs Dodonaea lanceolata

Main grasses Dicanthium ^{fecundum} / Chrysopogon ^{fallax} / Heteropogon ^{contortus}

Others —

Vines —

Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual
		<u>1.5m Dicanthium</u>		

FIRE	When was last fire	<u>2003 1993</u>	<u>2002 1992</u>	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------------------	------------------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Presence of Heteropogon contortus

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

Cattle

HABITAT CHANGE

Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

Creek banks eroded + gullied + trampled (bare)

PTO for more notes space

Photos Y/N

Photo re-takes Y/N

Photos 7 + 8
Shack


Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland <u>Low Woodland A</u> Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass	Dense Mat Plants Dense Hummock Grass	Mat Plants Mid-Dense Hummock Grass	Open Mat Plants Hummock Grass	Very Open Mat Plants Open Hummock Grass
GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Tall Grass Dense Low Grass Dense Herbs	<u>Tall Grass</u> Low Grass Herbs	Open Tall Grass Open Low Grass Open Herbs	Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

co-dominants
Dicanthium + Chrysopogon fallax

TO MOUNT PERCY



In Creek

add Lophostemon grandiflorus to 10m
" Planchonia careya + Brachychiton diversifolius
some Euc. tectifica + Corymbia dampierii
Dodonaea lanceolata
Irotalaria linifolia
stemodia grossa
Ficus opposita

Kimberley Mammals -- Traplines

Team <u>TW</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>Oscar Range</u>	Trapline name and/or code <u>4</u>
--	-----------------------------	---------------------------------------

Lat <u>17° 38' 25.1" S</u>	Date traps set <u>21/5/04</u>	Cages <u>6</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>124° 56' 28.8" E</u>	Date traps lifted <u>25/5/04</u>			

Topography and aspect
Limestone Karst Outcrops + ridge / flat waxy valleys between

Rock type, nature and extent of exposure
Limestone ~ 50% + 50% flat-floored alluvial valley floors

Soil type, origin and depth
Deep fine textured clay-loam Sample ☒ YN ☒ X

VEGETATION (Muir system over page)

Main trees Adansonia + Terminalia + Bauhinia
gregarii hadleyana cunninghamii

Main Shrubs Erevillea parallela, Flueggea virosa, Dodonea platyptera

Main grasses Triodia pascoeana with Sorghum plumosum (common)

Others Gomphrena spp, Dichapera armata, Indigofera linifolia

Vines

Main grasses form & height	hummock 0.5 <u>Triodia p.</u>	Perennial tussock —	Perennial mats —	Annual <u>1.5m Sorghum plumosum</u>
----------------------------	----------------------------------	------------------------	---------------------	--

FIRE	When was last fire	<u>1993 (2003)</u>	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	--------------------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)
Boab bark flaked + lower branches ~~scorched~~ - tanned

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)
limestone ridges

OTHER DISTURBERS (e.g. cattle, pigs, weeds)
some cattle

HABITAT CHANGE Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

Annual Sorghum intruding into Triodia pascoeana (fire frequency)

PTO for more notes space

Photos Y/N ☒	Photo re-takes Y/N
--	--------------------


Photos 1 + 2
Sheet 1

Muir vegetation System

co-don Sorghum + Triodia

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE d 70-100%	MID-DENSE c 30-70%	SPARSE i 10-30%	VERY SPARSE r 2-10%
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland <u>Open Low Woodland A</u> Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass	Dense Mat Plants Dense Hummock Grass	Mat Plants Mid-Dense Hummock Grass	Open Mat Plants Hummock Grass	Very Open Mat Plants <u>Open Hummock Grass</u>
GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Tall Grass Dense Low Grass Dense Herbs	<u>Tall Grass</u> Low Grass Herbs	Open Tall Grass Open Low Grass Open Herbs	<u>Triodia pascuosa</u> Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes



Kimberley Mammals -- Traplines

Team <u>TW</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>Oscar Range</u>	Trapline name and/or code <u>5</u>
--	-----------------------------	------------------------------------

Lat <u>17° 38' 28.9" S</u>	Date traps set <u>21/5/04</u>	Cages <u>7</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>124° 55' 55.7" E</u>	Date traps lifted <u>25/5/04</u>			

Topography and aspect <u>Black Soil Plain</u>
--

Rock type, nature and extent of exposure
--

Soil type, origin and depth <u>Deep cracking clay E gilgai</u>	Sample <u>YN</u>
---	------------------

VEGETATION (Muir system over page)				
Main trees <u>Corymbia dampieri & Bauhinia cunninghamii</u>				
Main Shrubs <u>Hakea arborescens, Dolichandrone heterophylla</u>				
Main grasses <u>Cane Grass (Sorghum plumosum) to 2m with Mitchell Grass (Dicanthium</u>				
Others				
Vines				
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual
		<u>Dicanthium to 1.5m</u>		<u>Sorghum plumosum 2m</u>

to 1.5m (ecundum)

FIRE	When was last fire	<u>1903 2003</u>	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------------------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Already burnt in patches nearby

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

damp area

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

Cattle

HABITAT CHANGE

Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos YN

Photo re-takes Y/N


photos 5 + 6
stick 1

✓

Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes



Other plants noted:

Byblis liniflora
Mimulus vernalis
(mauve)

Cyperus sp.

Stemodia grossa
Goodenia odonellii
Ludwigia octovalvis
Sesbania simplicifolia to 2.5m
Crotalaria liniflora
Commelina ensifolia
Buchnera sp.
Flueggea virosa
Grevillea parallela
Crinum angustifolium (Lily)

Kimberley Mammals -- Traplines

CLIFFS

108 k W of Camp

Team <u>TW</u> (AB/TH) (NM/TH) (TS/AT) (CP +)	Location <u>EXEDDAR</u> <u>NAPIER RANGE</u>	Trapline name and/or code <u>1</u>
--	--	---------------------------------------

Lat <u>17° 15' 38.8"</u>	Date traps set <u>26/5/04</u>	Cages <u>8</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>124° 43' 06.7"</u>	Date traps lifted <u>30/5/04</u>			

Topography and aspect
Flats on S side of range

Rock type, nature and extent of exposure
nil

Soil type, origin and depth
Calcareous brown earth plain with gilgai holes

Sample Y/N ☒

VEGETATION (Muir system over page)

Main trees Bauhinia cunninghamii, Adansonia gregorii

Main Shrubs Sesbania simplicifolia, Hakea arborescens

Main grasses Dicanthium fecundum + Sorghum plumosum

Others

Vines nil

Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual
		<u>Dicanthium 0.5m</u>		<u>Sorghum 2m</u>

FIRE	When was last fire	<u>2003</u> <u>1993</u> ✓	<u>2002</u> <u>1992</u>	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------------------------------	----------------------------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

none

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

nil

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

Cattle

HABITAT CHANGE

Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N ☒

Photo re-takes Y/N

Photos 19 + 20
stick

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE d 70-100%	MID-DENSE c 30-70%	SPARSE i 10-30%	VERY SPARSE r 2-10%
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland <i>Bauhinia</i> <u>Low Woodland A</u> Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass	Dense Mat Plants Dense Hummock Grass	Mat Plants Mid-Dense Hummock Grass	Open Mat Plants Hummock Grass	Very Open Mat Plants Open Hummock Grass
GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Tall Grass Dense Low Grass Dense Herbs	<u>Tall Grass</u> <i>Curly Mitchell Grass</i> Low Grass Herbs	Open Tall Grass Open Low Grass Open Herbs	Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Ranges

Additional Notes

Plants

N
↑

C. dampieri

Sorghum
o *Ac suberosa*
o *Bauh*

Bauh o
o *Mitchell Grass*
o *Curly Mitchell Grass*
o *Sedgwickia*
o *Bauh*

bl. soil
gully with gilgai holes

o *E. tect*
o *Bauh* E

Track

Indigofera linifolia
Crotalaria linifolia
Grevillea parallela
Trichodesma zeylanica
Tinospora smilacina x 1
Hakea arborescens
Flueggea virosa
Neptunia sp.
Gomphrena flaccida
Chamaesyce sp.
Ptilotus calostachyus
Sida rohlenae
Chrysopogon fallax
Themeda triandra

Kimberley Mammals -- Traplines

37.
2.7 k W of Camp

Team <u>TW</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>CHEDDAR CLIFF</u> <u>NAPIER RA</u>	Trapline name and/or code <u>2</u>
--	---	---------------------------------------

Lat <u>17° 15' 28.5"</u>	Date traps set <u>26/5/04</u>	Cages <u>8</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>124° 42' 32.1"</u>	Date traps lifted <u>30/5/04</u>			

Topography and aspect

Flat plain on S side of range

Rock type, nature and extent of exposure

Sheet exposure in places (calcrete)

Soil type, origin and depth

Shallow Reddish-brown + black soil over limestone pavement

Sample Y/N ☒

VEGETATION (Muir system over page)

Main trees Corymba lamprocarpaMain Shrubs Hakea arborescensMain grasses Sorghum plumosum + Mitchell Grass (Dicanthium fecundum) - codominantOthers Grewia retusifolia, Indigofera linifolia, Portulaca sp.Vines nil

Main grasses form & height	hummock	Perennial tussock to 1.5m <u>Dicanthium fecundum</u>	Perennial mats	Annual <u>Sorghum plumosum</u> 2m
----------------------------	---------	---	----------------	--------------------------------------

FIRE	When was last fire <u>25th 1993</u> ☒	<u>1992</u> ☒	< 3 yrs	3-10 yrs	>10 yrs
------	--	---	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

none

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

nil

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

cattle

HABITAT CHANGE

Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N

Yes

Photo re-takes Y/N


21st + 22nd sticks

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% v
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland <u>Low Woodland A</u> Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass <u>Tall Grass</u> Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Plants noted



Crotalaria linifolia
Indigofera tinifolia
Chrysopogon fallax (etc)
Eragrostis sp.
Gomphrena
Spermacoce
Sida
Portulaca sp.
Cleome viscosa
Mitrasacme
Goodenia
Grewia retusifolia
Ficus opposita x 1
Themeda triandra
Hybanthus aurantiacus
Bauhinia cunninghami x 1
Adansonia gregorii x 1 (small)
Mitrasacme
Goodenia

Kimberley Mammals -- Traplines

4.0 W of Camp

Team <i>FW</i> (AB/TH) (NM/IR) (TS/AT) (CP +)	Location <i>CHADDA CLIFFS</i> <i>NAPIER Ra</i>	Trapline name and/or code <i>3</i>
--	---	---------------------------------------

Lat <i>17° 15' 19.0"</i>	Date traps set <i>26/5/04</i>	Cages <i>8</i>	Big Elliott <i>10</i>	med Elliott <i>18</i>
Long <i>124° 42' 47.5"</i>	Date traps lifted <i>30/5/04</i>			

Topography and aspect <i>S. Side + bp of Napier Range southern aspect</i>
--

Rock type, nature and extent of exposure <i>>70% rock exposure. very massive exposures</i>
--

Soil type, origin and depth <i>skeletal shallow in crevices etc</i>	Sample Y/N ☒
--	--

VEGETATION (Muir system over page)				
Main trees	<i>Celtis phillipensis, Gyrocarpus americanus, Vitex glabrata</i>			
Main Shrubs	<i>Plumbago zeylanica, Abutilon andrewsianum</i>			
Main grasses	<i>Triodia pascoeana</i>			
Others				
Vines	<i>Cayratia aff. cardiophylla, Adenia heterophylla</i>			
Main grasses form & height	hummock <i>Triodia 1.5m</i>	Perennial tussock	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	<i>>10 yrs</i>
------	--------------------	------	------	---------	----------	-------------------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Rainforest species in good condition

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Rock outcrops (massive)

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

some Passiflora foetida seen

HABITAT CHANGE

Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N ☒

Photo re-takes Y/N

*photo 13+14**stick**prob.*

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE d 70-100%	MID-DENSE c 30-70%	SPARSE i 10-30%	VERY SPARSE r 2-10%
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland <i>Celtis</i> Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass	Dense Mat Plants Dense Hummock Grass	Mat Plants Mid-Dense Hummock Grass	Open Mat Plants Hummock Grass	Very Open Mat Plants Open Hummock Grass
GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Tall Grass Dense Low Grass Dense Herbs	Tall Grass Low Grass Herbs	Open Tall Grass Open Low Grass Open Herbs	Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

S side Napier Range

Boab
rock
scree
Dipliptera

Gyrocarpus
Celtis
Plumbago
rilled Karst with Triodia

Celtis
Triodia
Vitex
Plumbago
Achras
Capparis jacobsonii
Hibiscus sp (big star calyx)
Cynanchum pedunculatum
Cleome viscosa
Tinospora smilacina
Jacquemontia paniculata
Strychnos lucida
Gyrocarpus americanus
Celtis philippensis
Bauhinia cunninghamii
Cayratia cardiophylla
Diospyros maritima
Adenia heterophylla
Sarcostemma viminalis
Brucea javanica
Flueggea virosa
Alstonia aff linearis
Melia azedarach
Opilia amantacea
Achras aspera

Species noted

Dipliptera armata
Terminalia hadleyana
Trichodesma zeylanica
Abutilon andrewsianum
Terminalia volucris
Eragrostis sp.
Plumbago zeylanica
Pennisetum basedowii
Grewia breviflora
Helicteres rhyocarpa
Trema ~~orientalis~~ tomentosa
Ficus mombasa

Kimberley Mammals -- Traplines

10.8 k W. of Camp

37.

Team <u>PS</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>QUEDAR CLIFFS</u> <u>NAPIERRA</u>	Trapline name and/or code <u>4</u>
--	--	---------------------------------------

Lat <u>17° 13' 48.1"</u>	Date traps set <u>26/5/04</u>	Cages <u>8</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>124° 38' 26.0"</u>	Date traps lifted <u>30/5/04</u>			

Topography and aspect
Flat plain

Rock type, nature and extent of exposure
—

Soil type, origin and depth
Black soil

Sample Y/N ☒

VEGETATION (Muir system over page)

Main trees Acacia suberosa

Main Shrubs Dolichandrone heterophylla, Sesbania simplicifolia

Main grasses Dicanthium fecundum

Others Bulbostylis barbara

Vines —

Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual
		<u>Dicanthium</u>		

FIRE	When was last fire	<u>2003</u> <u>4093</u>	<u>2002</u> <u>1992</u>	< 3 yrs	3-10 yrs	>10 yrs
-------------	--------------------	----------------------------	----------------------------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Acacia suberosa (Killed by fire, regen. from base)

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

none

OTHER DISTURBERS (e.g. cattle, pigs weeds)

(Cattle)

HABITAT CHANGE

Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N ☒


Photo re-takes Y/N

Photo 23+24
stick 1

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass <u>Tall Grass</u> Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes



lines of *Sesbania simplicifolia* to 2.5m with *Sorghum* on gilgai

Other plants

Crotalaria linifolia
Sorghum plumosum
Acacia suberosa
Buchnera
Ficus opposita x 1
Carissa lanceolata
Goodenia
Bulbostylis barbata
Mitrasacme
Themeda triandra
Ludwigia octovalvis
Triodia pascoeana (occ.)
Dolichandrone heterophylla (as mini shrub)

Kimberley Mammals -- Traplines

37.
6.8 km W of Camp in Barker Gap

Team <u>JW</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>Cheddar Cliff</u> <u>Wapier Range</u>	Trapline name and/or code <u>5</u>
--	--	---------------------------------------

Lat <u>17° 14' 49.3"</u>	Date traps set <u>26/5/04</u>	Cages <u>8</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>124° 40' 30.8"</u>	Date traps lifted <u>30/5/04</u>			

Topography and aspect up slopes into top of range
generally S aspect

Rock type, nature and extent of exposure 80% exposed limestone

Soil type, origin and depth skeletal s.s. , v. patchy
Sample Y/N ☒

VEGETATION (Muir system over page)

Main trees Terminalia volucris, T. hadleyana, Vitex glabrata

Main Shrubs Grewia reusifolia, Flueggea virosa

Main grasses Sorghum plumosum, Triodia pascoana, Heteropogon contortus

Others

Vines Operculina brownii

Main grasses form & height	hummock <u>Triodia</u>	Perennial tussock	Perennial mats	Annual <u>Sorghum 2m</u>
----------------------------	---------------------------	-------------------	----------------	-----------------------------

FIRE	When was last fire <u>2003 1993</u> ✓	<u>1992</u>	< 3 yrs	3-10 yrs	>10 yrs
-------------	---------------------------------------	-------------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)
Karst site degraded by fire ascending terrace
(Sorghum & Heteropogon replacing Triodia)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))
Rainforest species under cliffs in sinkholes
eg Vitex glabrata, Ficus virens, Diospyros
maritima

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)
Massive rocks, sinkholes in 'moonscape' plateau

OTHER DISTURBERS (e.g. cattle, pigs, weeds)
no, some weeds

HABITAT CHANGE Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage


loss of vine thicket, replaced by
savanna grassland

PTO for more notes space	Photos Y/N ☒	Photo re-takes Y/N
--------------------------	--	--------------------

Photos 15-18
Sheet 1

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland <u>Low Woodland A</u> Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass <u>Tall Grass</u> Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses



Plants noted

Abutilon andrewsianum
Trichoderma zeylanica
Ficus opposita
Albizia lebbek
Vitex glabrata
Heteropogon contortus
Celtis philipensis
Solanum
Diospyros maritima
Calceolaria ianabii

Alstonia linearis
Ficus virens
Commelina ensifolia
Cleome viscosa
Trema orientalis
Helicteres rhynchocarpa
Ficus platypoda
Terminalia hadleyana
 " *volucris*
 " *canescens*
Diplocyclos palmata
Flueggea virosa
 " " "

Grewia retusifolia
 " *bravifolia*
Cymbopogon procerus
Bauhinia cunninghamii
Operculina brownii
Dolichandrone heterophyll
Crotalaria linifolia
Gomphrena
Sida rohlenae
Eragrostis sp.

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location MT GLADYS	Trapline name and/or code 1a
--	-----------------------	---------------------------------

Lat 17° 21' 57.1"	Date traps set 3/5/04	Cages 4	Big Elliott 5	med Elliott 9
Long 125° 32' 01.5"	Date traps lifted 4/6/04			

Topography and aspect gentle slope

Rock type, nature and extent of exposure Granitoid Clay-loam Reddish c rocks in profile
--

Soil type, origin and depth several old flattened termite mounds skeletal soil, deep	Sample Y/N Y
--	-----------------

VEGETATION (Muir system over page)				
Main trees	Eucalyptus tectifica, Bauhinia cunninghamii			
Main Shrubs	Cochlospermum fraseri, Grevillea parallela			
Main grasses	Sorghum plumosum, Heteropogon contortus, Triodia bitextura			
Others	Indigofera linifolia			
Vines				
Main grasses form & height	hummock 0.5 Triodia m	Perennial tussock	Perennial mats	Annual Sorghum 1.5m

FIRE	When was last fire	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Fire scars & Heteropogon frequency

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

none

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

none

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

(Cattle)

HABITAT CHANGE

Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N


Photo re-takes Y/N

Photos 35 & 36
shot 1

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland <u>Low Woodland A</u> <u>Low Woodland B</u>	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants <u>Mid-Dense Hummock Grass</u> <u>Tall Grass</u> Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes



Plant list

Eucalyptus tectifica
 Bauhinia cunninghamii
 Solanum leopoldense
 Cymbopogon procerus
 Distichlis spicata
 Cochlospermum fraseri
 Indigofera linifolia
 Triodia bixetura
 Grevillea parallela
 Chamaesyce distans
 Goodenia Odonnellii
 Polycarpaea involucrata
 Terminalia canescens x 1
 Brachychiton diversifolius
 Cullen badocana
 Mitrasacme sp.
 Dolichandrone heterophylla (stunted)
 Corchorus pumilio
 Dodonaea physocarpa x 1
 Gomphrena cunninghamii
 Buchnera linearis

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location MT GLADYS	Trapline name and/or code Line 1 b
--	-----------------------	---------------------------------------

Lat 17° 21' 47.4"	Date traps set 31/5/04	Cages 4	Big Elliott 5	med Elliott 9
Long 125° 31' 41.9"	Date traps lifted 4/6/04			

Topography and aspect
gentle slope

Rock type, nature and extent of exposure
Granitoid Unit stony profile

Soil type, origin and depth
reddish gravel on surface
skeletal clay-loam, deep

Sample ☒ Y/N

VEGETATION (Muir system over page)

Main trees Eucalyptus tectifica, Corymbia dampieri, Brachychiton diversifolius

Main Shrubs Trichodesma zeylanicum, Grewia retusifolia

Main grasses Dicanthium feundum, Chrysopogon fallax, Sehima nervosum

Others

Vines

Main grasses form & height	hummock	Perennial tussock Sehima Dicanthium Chrysopogon	Perennial mats	Annual
----------------------------	---------	---	----------------	--------

FIRE	When was last fire	1993 ✓	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	--------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Fire scars on trees

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

none

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

none

OTHER DISTURBERS (e.g cattle, pigs weeds)

(Cattle)

HABITAT CHANGE Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N


Photo re-takes Y/N

Photos 37 + 38
Sheet 1

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE d 70-100%	MID-DENSE c 30-70%	SPARSE i 10-30%	VERY SPARSE r 2-10%
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland <u>Low Woodland A</u> Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass <u>Tall Grass</u> <u>Low Grass</u> Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes



Plant List

Eucalyptus tectifica
Corymbia dampieri x 2
Brachychiton diversifolius
Trichoderma zeylanicum
Pterocaulon sphacelatum
Bauhinia cunninghamii x 1
Buchnera linearis
Gomphrena sp.
Crotalaria linifolia
Indigofera linifolia
Chamaesyce distans
Indigofera hirsuta
Grewia reusifolia
Heteropogon contortus
Sida sp.

Kimberley Mammals -- Traplines

Team <u>TW</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>MT GLADYS</u>	Trapline name and/or code <u>Line 2</u>
--	------------------------------	--

Lat <u>17° 24' 06.9"</u>	Date traps set <u>31/5/04</u>	Cages <u>8</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>125° 35' 29.6</u>	Date traps lifted <u>4/6/04</u>			

Topography and aspect
Gently sloping Creekline, southerly aspect

Rock type, nature and extent of exposure
Dolomite scree exposed in creek & creek alluv. sand + deep alluvial soil

Soil type, origin and depth
alluvium, granitoid origin clay-loam

Sample Y N

VEGETATION (Muir system over page)

Main trees Eucalyptus camaldulensis, Terminalia platyphylla

Main Shrubs Acacia plectocarpa, Ludwigia spp.

Main grasses Heteropogon contortus, Setaria sp

Others

Vines

Main grasses form & height	hummock	Perennial tussock <u>Heteropogon 1m</u>	Perennial mats	Annual
----------------------------	---------	--	----------------	--------

FIRE	When was last fire	<u>2003</u> 1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	--------------------------------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Heteropogon contortus

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Acacia plectocarpa & A. neurocarpa on creek

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Permanent creek

OTHER DISTURBERS (e.g. cattle, pigs weeds)

Cattle, horses

* Stylosanthes hamata on track verge

HABITAT CHANGE

Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y N


Photo re-takes Y N

Photo 34 Shuck!

Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest <i>riverine</i>	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns	Dense Ferns	Ferns	Open Ferns	Very Open Ferns
	Mosses, liverwort	Dense Mosses	Mosses	Open Mosses	Very Open Mosses

Additional Notes



Plants noted

Terminalia platyphylla
Eucalyptus camaldulensis
Cymbidium canaliculatum (host)
Sesbania formosa
Lophostemon grandiflorus
Ficus coronulata
Timonius timon
Melaleuca argentea
Pandanus spiralis
Eucalyptus tectifica
Acacia suberosa
Acacia neurocarpa
Grewia retusifolia
Brachychiton diversifolius
Sorghum plumosum
Phyllanthus reticulatus
Bauhinia cunninghamii
Operculina brownii
Crotalaria linifolia
Indigofera linifolia
Livistona mariae var. *occidentalis*
Acacia plectocarpa
Ludwigia octovalvis
 " *perennis*
Sehima nervosum
Tephrosia dhaeosperma

Sida subspicata
Heteropogon contortus
Arundinella nepalensis
Setaria sp.

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location MT GLADYS	Trapline name and/or code Line 3
--	-----------------------	-------------------------------------

Lat 17° 24' 33.7"	Date traps set 31/5/04	Cages 8	Big Elliott 10	med Elliott 18
Long 125° 36' 02.7"	Date traps lifted 3/6/04			

Topography and aspect
? Dolomite or Granitoid Hill, northerley aspect steep slope & scree

Rock type, nature and extent of exposure
60% exposure overall or massive bare scree

Soil type, origin and depth
skeletal between scree

Sample Y/N

VEGETATION (Muir system over page)

Main trees	Eucalyptus tectifica, Bauhinia cunninghamii		
Main Shrubs	Cochlospermum fraseri		
Main grasses	Sesima nervosum, Sorghum plumosum		
Others	Indigofera linifolia		
Vines	Operculina brownii on rocks		
Main grasses form & height	hummock	Perennial tussock	Perennial mats
		Sesima 1m	Annual Sorghum 1.5m

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Abundant sorghum below scree

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Celtis & Vitex in scree/boulder face

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Massive rock outcrop + scree

OTHER DISTURBERS (e.g. cattle, pigs weeds)

Old donkey dung noted

HABITAT CHANGE

Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N

Photo re-takes Y/N

Photo 30 stick 1

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses


Additional Notes

Slopes

Euc. tectifica
Sehima nervosum
Buchnera linearis
Crotalaria linifolia
Chamaesyce distans
Cleome ~~viscosa~~
Indigofera linifolia

Plants in rock/summit

Bauhinia cunninghamii
Dolichandrone heterophylla
Flueggea virosa
Ficus platypoda
Vitex glabrata
Operculina brownii
Trema orientalis
Brachychiton viscidulus
Abutilon andrewsianum
Ehretia saligna
Trichodesma zeylanicum
Atalaya hemiglauc
Premna acuminata
Capparis jacobsonii
Cleome viscosa
Solanum leopoldense
Celtis philippensis
Cochlospermum fraseri
Terminalia canescens
Pennisetum basedowii
Cayratia trifolia
Eragrostis sp.
Senna glutinosa



Kimberley Mammals -- Traplines

Team <u>TW</u> (AB/TH) (NM/AR) (TS/AT) (CP +)	Location <u>MT GLADYS</u>	Trapline name and/or code <u>Line 4</u>
--	------------------------------	--

Lat <u>17° 23' 46.3"</u>	Date traps set <u>31/5/04</u>	Cages <u>8</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>125° 35' 32.1"</u>	Date traps lifted <u>4/6/04</u>			

Topography and aspect
Sandstone Gully + Gorge, Southern aspect

Rock type, nature and extent of exposure
King Leopold Sandstone, 60% (scree slopes etc)

Soil type, origin and depth
peaty skeletal on slopes, alluvial / loam in gully

+ gorge bottom
Sample ☒ Y/N

VEGETATION (Muir system over page)

Main trees Lophosiemon grandiflorus, Euc. ptychocarpa, E. tectifica

Main Shrubs Grevillea agrifolia, Cajanus cinereus, Ludwigia octovalvis

Main grasses Arundinella nepalensis, Setaria sp., Triodia sp.

Others Livistona mariae var. occidentalis (fan palm)

Vines

Main grasses form & height	hummock, 0.5 <u>Triodia m</u>	Perennial tussock <u>Arundinella 2m</u>	Perennial mats	Annual
----------------------------	----------------------------------	--	----------------	--------

FIRE	When was last fire	1993	1992	< 3 yrs	<u>3-10 yrs</u>	>10 yrs
-------------	--------------------	------	------	---------	-----------------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Fine scars + Heteropogon frequency on trees

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Riverine species

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Rocky gorge walls, permanent creek

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

Cattle (dung evidence)
* weeds - Euphorbia hirta
Stylosanthes hamata

HABITAT CHANGE Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space


Photos ☒ Y/N

Photo re-takes Y/N

Photo 34-36 stuck!

Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns	Dense Ferns	Ferns	Open Ferns	Very Open Ferns
	Mosses, liverwort	Dense Mosses	Mosses	Open Mosses	Very Open Mosses



Additional Notes

Plants noted

Livistona mariae var. *occidentalis*
Corymbia cadophora
Erythrophloeum chlorostachys
Grewia retusifolia
Canarium australianum
Ficus platypoda
Ficus hispida
Owenia vernicosa
Terminalia hadleyana
Coelosperrum fraseri
Brachychiton viscidulus
Eucalyptus tectifica
Melaleuca argentea
Trema ~~orientalis~~ tomentosa
Phyllanthus reticulatus
Trichodesma zeylanicum
Cajanus cinereus
Acacia neurocarpa
Buchanania obovata
Sida subspicata
Grevillea agrifolia
Dicliptera armata
Cymbopogon procerus
U...

Setaria sp.
Arundinella nepalensis
Chamaesyce sp.
Potamogeton sp. (aquatic)
Eriozaulon sp. (aquatic)
Eriachne glauca

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location MT GLADYS	Trapline name and/or code Live 5a
--	-----------------------	--------------------------------------

Lat 17° 24' 35.2"	Date traps set 31/5/04	Cages 4	Big Elliott 5	med Elliott 9
Long 125° 39' 07.1"	Date traps lifted 4/06/04			

Topography and aspect Gentle southern aspect, creek bank

Rock type, nature and extent of exposure Some stones in profile + surface rocks
--

Soil type, origin and depth Deep brown alluvial clay-loam	Sample Y/N
--	------------

VEGETATION (Muir system over page)				
Main trees	Eucalyptus tectifica, Bauhinia cunninghamii			
Main Shrubs	Crotalaria novae-hollandiae, Grewia retusifolia			
Main grasses	Dicanthium fecundum, Sehima nervosum, Cymbopogon procerus			
Others				
Vines	Cassytha filiformis			
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual
		Sehima 1m	2.5m	

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	-----------------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Creek - permanent

OTHER DISTURBERS (e.g cattle, pigs weeds)
Cattle
* Stylosanthes hamata (on verge)


HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	
Photos Y/N	Photo re-takes Y/N

Photo 33 Stick 1
Photo 46 Stick 1

Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	<u>Low Woodland A</u>	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1 0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	<u>Tall Grass</u>	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes



Plants noted

Planchonia careya
Lophostemon grandiflorus
Euc. tectifica
Euc. camaldulensis
Hakea suberea
Corymbia flavescens
Dicerna biarticulatum
Goodenia odonnellii
Pterocaulon sphacelatum
Trichodesma zeylanicum
Grewia rebusifolia
Portulaca bicolor
Flueggea virosa
Sida subspicata
Brachychiton diversifolius
Corchorus sp.
Jasminum didymum
Sehima nervosum
Cymbopogon procerus
Chamaesyce distans
Polycarpha sp.

Kimberley Mammals -- Traplines

Team <u>TS</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>MT GLADYS</u>	Trapline name and/or code <u>Line 5b</u>
--	---------------------------	---

Lat <u>17° 24' 40.3"</u>	Date traps set <u>31/5/04</u>	Cages <u>4</u>	Big Elliott <u>5</u>	med Elliott <u>9</u>
Long <u>125° 37' 57.9"</u>	Date traps lifted <u>4/6/04</u>			

Topography and aspect <u>Flat, seasonally swampy</u>

Rock type, nature and extent of exposure <u>Some stones below profile + on surface</u>

Soil type, origin and depth <u>Black heaving clay gilgai surface</u>	algal mat in gilgai holes	Sample <u>Y/N</u>
---	---------------------------	-------------------

VEGETATION (Muir system over page)				
Main trees	<u>Eucalyptus tectifica & Bauhinia cunninghamii</u>			
Main Shrubs	<u>Carissa lanceolata</u>			
Main grasses	<u>Eriachne glauca</u>			
Others	<u>Bonamia pinnosa (ground cover)</u>			
Vines				
Main grasses form & height	hummock	Perennial tussock <u>1m</u>	Perennial mats	Annual
		<u>Eriachne</u>		

FIRE	When was last fire	<u>2003</u> 1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	---------------------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) <u>nil</u>
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) <u>none</u>

OTHER DISTURBERS (e.g cattle, pigs weeds)
<u>(Cattle)</u>


HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	
Photos <u>Y/N</u>	Photo re-takes Y/N

Photo 32 Stick 1

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes



Plants noted

Eucalyptus tectifica
 Baobab cunninghamii
 Atalaya hemiglauc
 Grewia retusifolia
 Bonamia pinnosa
 Neplunia sp.
 Gomphrena sp.
 Indigofera linifolia
 Buchnera linearis
 Blumea integrifolia
 Abutilon andrewsianum
 Polycarpaea sp.
 Aeschynomene indica
 Ipomoea coptica
 Eriachne glauca

Kimberley Mammals -- Traplines

Team <u>TW</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>FOREST CREEK</u>	Trapline name and/or code <u>1a</u>
--	---------------------------------	--

Lat <u>14° 38' 57.9"</u>	Date traps set <u>6 June 04</u>	Cages <u>5</u>	Big Elliott <u>5</u>	med Elliott <u>10</u>
Long <u>126° 56' 39.5"</u>	Date traps lifted <u>10 June 04</u>			

Topography and aspect <u>NW aspect Sandstone scree slope & terraces</u>
--

Rock type, nature and extent of exposure <u>WARTON s/s 80% exposure</u>
--

Soil type, origin and depth <u>skeletal, sandy, shallow</u>	Sample <u>YN</u>
--	------------------

VEGETATION (Muir system over page)				
Main trees	<u>Corymbia greeniana, Terminalia hadleyana</u>			
Main Shrubs	<u>Petalostigma quadriloculare, Grevillea agrifolia</u>			
Main grasses	<u>Triodia microstachya</u>			
Others	<u>Livistona loriphylla (fan palm)</u>			
Vines	<u>Flagellaria indica</u>			
Main grasses form & height	hummock <u>Triodia lm</u>	Perennial tussock	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	<u>3-10 yrs</u>	>10 yrs
------	--------------------	------	------	---------	-----------------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) <u>size of Triodia hummocks</u>
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) <u>Acacia aff. translucens</u>
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) <u>sandstone outcrops</u>

OTHER DISTURBERS (e.g cattle, pigs weeds) <u>None</u>
--

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	
Photos <u>YN</u>	Photo re-takes Y/N

Photo 41 / stick 1
Photo 45
46

Muir vegetation System


LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	<u>Low Woodland A</u>	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	<u>Mid-Dense Hummock Grass</u>	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns	Dense Ferns	Ferns	Open Ferns	Very Open Ferns
	Mosses, liverwort	Dense Mosses	Mosses	Open Mosses	Very Open Mosses

Additional Notes

Plants noted

Owenia vernicosa
Petalostigma quadrilobulare
Triodia microstachya
Livistona loriphylla
Grevillea agrifolia
Acacia dunnii
Tephrosia sp.
Acacia aff. *transluens*
Sida sp. A
Phyllanthus aridus
Boronia lanuginosa
Lilicis aff. *superbus* (pink fls)
Lilbertia
Lilbertia
Zinnocarpus sp. A.
Sardenia megasperma
Tagellaria indica

Buchanania obovata
Corymbia greeniana
Bossiaea bossiaecoides
Erythrophleum chlorostachys
Vitex glabrata
Calytrix exstipulata



O Camp

↓
N

Kimberley Mammals -- Traplines

29

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location FOREST CREEK	Trapline name and/or code 1b
--	--------------------------	---------------------------------

Lat 14° 38' 54.1"	Date traps set 6 June 04	Cages 3	Big Elliott 5	med Elliott 8
Long 126° 56' 38.7"	Date traps lifted 10 June 04			

Topography and aspect	top + SE slope of sand dune perpendicular to river bend
-----------------------	---

Rock type, nature and extent of exposure	nil
--	-----

Soil type, origin and depth	deep fine yellowish sand	Sample ☒ Y ☐ N
-----------------------------	--------------------------	---

VEGETATION (Muir system over page)				
Main trees	Callitris intratropica (on dune crest), Pouteria arnhemica			
Main Shrubs	Calytrix exstipulata, Acacia tumida, Verticordia cunninghamii			
Main grasses	Triodia microstachya & Sorghum plumosum			
Others	Spermacoce sp & Polycarpaea longiflora			
Vines	Opilia amentacea (on dune crest)			
Main grasses form & height	hummock Triodia 1m	Perennial tussock	Perennial mats	Annual Sorghum 2m

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)	Size of Acacia tumida (5m)
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))	Unburnt Cypress on dune crest to 7m & tall Acacia tumida
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)	Drysdale River nearby

OTHER DISTURBERS (e.g. cattle, pigs weeds)	Cattle (dung, pads & dustwallow)
--	----------------------------------

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	Photos ☒ Y ☐ N
	Photo re-takes Y/N

Photo 42 stuck

Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Plants noted

Callitris intratropica
Calytrix exstipulata
Triodia microstachya
Jacksonia argentea
Hibbertia lepidota
Verticordia cunninghamii
Grevillea refracta
Grevillea heliosperma
Acacia tumida
Polycarpaea longifolia
Sorghum plumosum
Pouteria arnhemica
Pouteria sericea
Opilia amentacea
Corymbia polycarpa
Buchera linearis
Jasminum didymum

Livistona loriphylla
Premna acuminata
Exocarpos latifolius
Erythrophleum chlorostachys
Spermacoce sp. (dead)
Denhamia obscura
Syzygium eucalyptoides
Persoonia falcata
Canarium australianum
Triumfetta sp.

[see diagram 1A]

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location FOREST CREEK	Trapline name and/or code 2
--	--------------------------	--------------------------------

Lat 14° 38' 57.8"	Date traps set 6 June 04	Cages 8	Big Elliott 10	med Elliott 18
Long 126° 56' 42.9"	Date traps lifted 10 June 04			

Topography and aspect Western aspect Sandstone scree slope ± terrace

Rock type, nature and extent of exposure Warton Sandstone 85%
--

Soil type, origin and depth skeletal sandy shallow	Sample ☒ Y ☐ N
---	---

VEGETATION (Muir system over page)				
Main trees	Terminalia hadleyana to 6m, Owenia vernicosa			
Main Shrubs	Petalostigma quadrilobulare, Acacia dunnii			
Main grasses	Triodia microstachya, Triodia U leaf, resinous, serrated edge — upper slopes on			
Others	Livistona loriphylla (palm)			
Vines				
Main grasses form & height	hummock 1m Triodia	Perennial tussock	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc) size of Triodia hummocks
--

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) Acacia dunnii (slender to 6m)

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Sandstone pavements & boulders
--

OTHER DISTURBERS (e.g. cattle, pigs weeds) nil

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space	
Photos ☒ Y ☐ N	Photo re-takes Y/N

Photos 43 + 44 + 52 + 45
stack

Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	<u>Low Woodland A</u>	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	<u>Mid-Dense Hummock Grass</u>	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes


Plants noted

Cochlospermum fraseri
Petalostigma quadrilobulare
Sprevillea agrifolia
Sardenia megasperma
Sardenia resinosa
Boronia lanuginosa
Livistona loriphylla
Terminalia hadleyana
Buchanania obovata
Owenia vernicosa
Croton ~~argutus~~ arnhemicus
Planchonia careya
Jacksonia argentea
Ricinocarpus sp A
Pouteria arnhemica
Xanthostemon paradoxus
Acacia dunnii
Brachychiton aff. *tridentatus*

Distichostemon hispidulus

Triumfetta sp.

Triodia microstachya
Triodia sp.



camp

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location FOREST CREEK	Trapline name and/or code 3
--	--------------------------	--------------------------------

Lat 14° 39' 02.9"	Date traps set 6 June 04	Cages	Big Elliott	med Elliott
Long 126° 56' 33.2"	Date traps lifted 10 June 04	8	9	18
TRAPS SET: 8 June 04		+ 8	+ 10	+ 18

Topography and aspect NW aspect, gully into sandstone ranges

Rock type, nature and extent of exposure Warton Sandstone 40% exposure

Soil type, origin and depth deep alluvium (sandy)	Sample Y/N
--	------------

VEGETATION (Muir system over page) Xanthostemon eucalyptoides				
Main trees Syzygium nervosum, Carallia brachiata, Corymbia ptychocarpa				
Main Shrubs Exocarpos latifolius, Hibiscus sp.				
Main grasses Chloris sp (on creek edge)				
Others Pandanus spiralis & Livistona loriphylla				
Vines Flagellaria indica & Smilax australis				
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual
		Chloris 1m		

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Syzygiums + ferns [western side]

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Permanent Creek

OTHER DISTURBERS (e.g cattle, pigs weeds)

Cattle (pads + dung) damaging riverine understorey.
* Passiflora foetida (grazing on Chloris)

HABITAT CHANGE

Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N

Photo re-takes Y/N

Photo 48 + 49 stick 1

Muir vegetation System


LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Plants noted

Melaleuca argentea
Syzygium nervosum
 " *eucalyptoides*
Carallia brachiata
Buchanania obovata
Exocarpos latifolius
Corymbia ptychocarpa
Erythrophleum chlorostachys
Andanus spiralis
Livistona loriphylla
Smilax australis
Passiflora foetida
Corymbia polycarpa
Canarium australicum
Acacia plectocarpa
Grevillea pteridifolia
Melastoma affine
Outeria sericea
Cestrum aurantiacum

Stemodia lythrifolia
Hibiscus sp. (pink fl)
Dicranopteris linearis (fernbank)



Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location FOREST CREEK	Trapline name and/or code 4
--	--------------------------	--------------------------------

Lat 14° 38' 54.2"	Date traps set 6 JUNE 04	Cages 8	Big Elliott 10	med Elliott 18
Long 126° 56' 30.0"	Date traps lifted 8 June 04			

Topography and aspect Flat level banks of Drysdale River

Rock type, nature and extent of exposure
--

Soil type, origin and depth Deep alluvial sand	Sample Y/N
---	------------

VEGETATION (Muir system over page)				
Main trees	Melaleuca argentea (to 20m on 'island' at mouth of Forest Creek)			
Main Shrubs	Barringtonia acutangula, Phyllanthus reticulatus			
Main grasses	Sorghum plumosum, Eragrostis sp.			
Others				
Vines	* Passiflora foetida			
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual Sorghum 2m

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Drysdale River x Forest Creek (permanent)

OTHER DISTURBERS (e.g. cattle, pigs, weeds)
Flood has ripped most of fringing forest away - some regeneration * Passiflora foetida Cattle (pads + dung)

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
Once rich fringing paperbark grove now stripped to bare level banks + regenerating stumps + trees.	
PTO for more notes space	Photos Y/N
	Photo re-takes Y/N

photo 50 + 51 stick 1


Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Plants noted

Melaleuca argentea
Barringtonia acutangula
Eucalyptus camaldulensis
Antidesma ghaesembilla
Passiflora foetida
Scaevola orientalis (occ)
Selousia campestris
Xylocarpus reticulatus
Acacia tumida
 " *holosericea*
Tagellaria indica
Agrostis sp.
Sorghum plumosum



Kimberley Mammals -- Traplines

Team <u>TW</u> (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <u>FOREST CREEK</u>	Trapline name and/or code <u>5</u>
--	---------------------------------	---------------------------------------

Lat <u>14° 39' 01.2"</u>	Date traps set <u>6 JUNE 04</u>	Cages <u>8</u>	Big Elliott <u>10</u>	med Elliott <u>18</u>
Long <u>126° 56' 39.4"</u>	Date traps lifted <u>10 June 04</u>			

Topography and aspect <u>Western Aspect Scree Slope</u>
--

Rock type, nature and extent of exposure <u>Warton Sandstone scree</u>

Soil type, origin and depth <u>skeletal soil</u>	Sample <u>YN</u>
---	------------------

VEGETATION (Muir system over page) <u>Xanthostemon eucalyptoides</u>				
Main trees <u>Corymbia greeniana, Terminalia hadleyana, Owenia vernicosa</u>				
Main Shrubs <u>Petalostigma quadriloculare, Acacia dunnii</u>				
Main grasses <u>Triodia microstachya (fine leaf spinifex)</u>				
Others <u>Livistona loriphylla (ribbon leaf fan palm)</u>				
Vines				
Main grasses form & height	hummock 1m	Perennial tussock	Perennial mats	Annual
<u>Triodia</u>				

FIRE	When was last fire	1993	1992	< 3 yrs	<u>3-10 yrs</u>	>10 yrs
------	--------------------	------	------	---------	-----------------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Xanthostemon, Syzygium spp + Carallia

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Upper creekline (Forest Creek)

OTHER DISTURBERS (e.g cattle, pigs weeds)

HABITAT CHANGE

Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N

Photo re-takes Y/N

Photo 47 stick

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% v
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest <i>Creek</i> Low Forest A Low Forest B	Tall Woodland Woodland <i>sandstone</i> Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes


Plants noted (sandstone ⁱⁿ rocks)

Triodia microstachya (dom.)
Corymbia greeniana (camp end)
Verticordia cunninghamii
Grevillea cunninghamii
Buchanania obovata
Etalostigma quadrilobulare (dom.)
Leivostoma loriphylla
Hibbertia lepidota
Acacia dunnii
Apparis umbonata x1
Butleria arnhemica
Ardenia megasperma
Buchnera linearis
Coronia lanuginosa
Wenia vernicosa
Erminalia hadleyana

Acacia humifusa
Grevillea agrifolia
Planchonia careya
Cymbopogon procerus
Ficus platypoda x1
Corymbia polycarpa

Pandanus spiralis
Melastoma affine
Smilax australis
Syzygium nervosum
"Syzygium eucalyptoides"
Carallia brachiata
Xanthostemon eucalyptoides

W



Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/ TR) (TS/AT)												
Location <i>Oscar Range</i>												
Record Number	Foot notes	Date	*1	*2	*3	Weight	Head body	Tail length	Sex	Age	Rproduction	*4
1	<i>Pseudomys nanus</i>	22/5/04	ME	M52691	toe clip 22	34.5			♀	Adult	—	4
2	<i>Zyzomys argurus</i>	22/5/04	"	M52690	"	32.0			♂	"	scrotal	2
3	<i>Pseudomys nanus</i>	"	"	M52688	"	16.4				S/A		5
4	" "	"	"	M52687	"	21.2				S/A		5
5	<i>Rattus tunneyi</i>	"	"	M52689	"	30.5				S/A		5
6	<i>Euro</i>	21/5/04		seen								4
7	"	"		scat								2
8	<i>Macropus antelopinus</i>	22/5/04		seen						Adult		0.5 k w d Camp in grassland
9	<i>Hipposideros alic</i>	22/5/04		recorded + seen								Foraging B/S/A grassland between lines 3 + 4
0	<i>Pseudomys nanus</i>	23/5/04		released						A		4 (valley floor grassland)

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Cattle

21/5/04 scat
+ tracks

1, 2, 3, 4, 5

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)				Location <i>Oscar Range</i>								
Record Number	Foot notes	Date	*1	*2	*3	Weight	Head body	Tail length	Sex	Age	Rproduction	*4
	Species		Source	Record type	Other samples & code numbers							Trapline
11	<i>Rattus tunneyi</i>	23/5/04	ME	M25694	toe for DNA X2	31gm			♂	S/A		5
12	"	"	LE	M25693	X2	78gm			♀	A		5
13	<i>Pseudomys delicatulus</i>	"	ME	M25692	X2	8.5gm			♂	A		5
14	<i>Tachyglossus aculeatus</i>	"	feces	—	—	—						4
15	Possum feces		"									4
16	<i>Pseudomys nanus</i>	24/5/04	"	released	ear tissue 01 X2	35g			♂	A	scutal test	4
17	<i>Rattus tunneyi</i>	"	"	"	tail tissue 02 X2	88g			♂	S/A		5
18	<i>Pseudomys nanus</i>	"	"	"	hair 03 X2	21g			♀	Young adult		5
19	<i>Zyzomys argurus</i>	25/5/04	ME	M52695	tissue 04 X2	22.5g			♂	S/A		2
20	<i>Pseudomys nanus</i>	"	"	released	—	36g			—	Adult		5

scratches + shot high up in valley on range on top of limestone ridge

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

TW

Team (AB/TH) (NM/TH) (TS/AT)					Location <i>Oscar Ranges</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
21	<i>Rattus tunneyi</i>	25/5/04	LE	released	tissue 05	48g			♂			5
22	"	"	ME	released	tissue 06	41g			♀			5
23	<i>Taphozous georgianus</i>	24/5/04	seen	seen	—	—			—			gorge near 2
24	Euro x 2	"	"	"	—	—			x			near line 4
25	Dingo 2+	23/5/04	heard									4
26	<i>Chaerophorobensis</i>	25/5/04	"									Barker Gap
27	<i>Saicolaimus flaviventris</i>	25/5/04	"									"
28	<i>Macropus agilis</i>	26/5/04	seen									1
29	Euro	"	faeces	(shit)								3
30	Rock Wallaby	"	"	"								3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

BARKER GAP,
NAPIER RANGE
"
"

4

Tw/Jiri + Marie Lockman

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Barker Gorge / Cheddar Cliff</i>							
Record Number	Foot notes Species	Date	*1 Source	*2 Record type	*3 Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	*4 Trapline
31	<i>Echidna</i>	26/5/04		faeces	—							3
32	<i>Euro</i>	27/5/04		"								5
33	<i>Rock Wallaby</i>	"		"								5
34	<i>Sandy Wallaby</i>	"	D	seen								2
35	<i>Petrogale brachyotis</i>	"	S	"	7.9 k	West of Barker Gap camp						cf 3, but bare sides of range
36	" "	"	S	"	"	"	"	"	"	"		cf. 3.
37	? Possum partial jaw	"			outside	cave at Cheddar Cliffs — may be recent.						cf 5
38	? Possums	"	S	seen at distance in	rock fig on	rim of range		7.7 k	W of camp			
39	"	"	S	"	"	"	"	"	"	8.2 km W of camp		
40	Possum, Rock Wallaby + ?	27/5/04	faeces	collected under	rock fig on	"	"	"	"	7.7 + 8.2 km W of camp		

} faeces in plastic bag in dry-pants box.

- *1 SOURCE Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)
- *2 RECORD TYPE WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR
- *3 OTHER SAMPLES Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number
- *4 TRAP LINE Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

5

TW, Jiri + Marie Lockman

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Cheddar Cliffs, Napier Range</i>							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
41	<i>Rattus tunneyi</i>	28/5/04	LE	escaped	—	65						1
42	<i>Pseudomys nanus</i>	"	ME	M52699	CK04 x2 tail	45			♂	A	scrutal	1
43	<i>Varanus acanthurus</i>	"	"	released								2
44	<i>Pseudomys nanus</i>	"	ME	M52698	CK03 x2 tail	33			♀	A	not preg	4
45	Euro faeces	"		faeces								5
46	<i>Zyzomys argurus</i>	"	LE	M52697	CK02 x2 tail	46			♀	A		3
47	Euro	"		faeces								3
48	Possum skull ? <i>Petrogale daltii</i>	"		M52696	CK01 x2 dry skin							3
49	<i>Echidna</i> shit	"		faeces								3
50	<i>Rattus tunneyi</i>	29/5/04	LE	M526700	CK05 x2	67			♂	A	scrutal	1

specimen in with McDowell

skeleton & dry skin & pa perhaps 2-3 months dead. In exposed site under tree outside cave

- *1 SOURCE Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)
- *2 RECORD TYPE WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR
- *3 OTHER SAMPLES Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number
- *4 TRAP LINE Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)				Location CHEDDAR CLIFFS / NAPIER R								
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
S1	Macropod carcass ? Rock Wallaby	29/5/04	carcass	calcaneum taken CK1								~5 k W of camp Outside a cave at foot of range
S2	Macrodera gigas	"	heard	heard	flying into back of cave complex							5.5 k W of Camp in Barker Gap
S3	Pseudomys delicatulus	30/5/04	ME	MS2701		11		left E Jiri Lockman			no photo	4
S4	Pseudomys nanus	"	ME	released		25	"	"	"	"	"	4
S5	Rattus tunneyi	"	ME	MS2702	tissue (toe) CK07 x2	47						4
S6	Nail-tailed Wallaby X2	"	seen	seen	—	—						North end of Wombarella Gap
S7	Cattle	29/5/04	scat tracks									1,2,5
S8												
S9												
S0												

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

6

specimen 2
Malt
McDonnell

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/LE) (TS/AT)					Location MOUNT GLADYS							
Record Number	Foot notes	Date	*1	*2	*3	Weight	Head body	Tail length	Sex	Age	Rproduction	*4
61	Pseudomys nanus	1/6/04	LE	M52704	MG02 x 2	24g						1b
62	Rattus villosissimus	"	ME	M52703	MG01 x 2	105g			♂	A	Scrotal	1b
63	Pseudomys nanus	"	ME	M52705	MG03 x 2	21.5g			♀			3
64	Agile Wallaby	"	seen	—								Gibb River Rd near Inghir
65	Pseudomys nanus	2/6/04	ME	M52707	MG05 x 2	27.2g			♂	A	Scrotal	1b
66	Euro	"										old shirt in cleft in
67	Echedne	"										"
68	Rattus villosissimus	"	ME	M52706	MG04 x 2	90gm			♂	A	Scrotal	4
69	Trichasaurus vulpecula	Feb 1987	seen									eastern foot of Mt Ord
70	Rattus villosissimus	3/6/04	ME	M52708	MG06 x 2	98gm			♂	A	Scrotal	1b

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Trap

line 3

"

in bloodwood trees + ~10 k from our Mt Gladys campsite

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location MT Gladys							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
71	Pseudomys nanus	3/6/04	ME	M52709	M607 x 2	19.1			♂			3
72	Taphozous georgianus	"	5 seen									3
73	Possum farras	2/6/04	collected						~ 2-3	days old		4
74	Rock wallaby farras	2/6/04	"						~ 2-3	days old.		4
75	Pseudomys nanus	4/6/04	ME	M52710	M608 toe tissue x 2	34			♀	A		1a
76	Rattus? villosusimus	4/6/04	ME	M52711	M609 toe tissue x 2	25.5				Juv		1b
77	Sac. flaviventris	3/6/04	heard									Mt Gladys Camp
78	Chaerophyllus farras	4/6/04	heard									Silent Grove
79	bandicoot diggery	31/5/04	diggery		In heavy							4
80	Cattle	3/6/04	scat tracks		riverine grassland at mouth of gorge.							1, 2, 4, 5

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

in my prep kit

cleft in rock exposed site under 3 Ficus platypoda

under 1 Ficus platypoda, exposed site

Mt Gladys Camp

Silent Grove camp/ranger station

9

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/ TR) (TS/AT)				Location <i>Forest Creek, DRNP</i>								
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
81	<i>Zyzomys argurus</i>	7/6/04	ME	M52712	FC01 x2 toe tissue	40.5g			♀	A		2
82	Euro	6/6/04	Seen									2
83	Euro	7/6/04	Scat									1
84	Euro	"	Scat									3
85	Cattle	"	Scat + tracks									4
86	"	"	"									3
87	"	"	"									1
88	<i>Petrogale? burbridgei</i>	"	spotlight + gun	M52713	FC02 x2 ear clip	1.2 kg	31.5	35	♂		Scrotal	2: further up scrotal slope
89	<i>Zyzomys argurus</i>	8/6/04	ME	M52714	FC03 x2 toe	26.5				S/A		5
90	" "	"	ME	M52715	FC04 x2 toe	41.5			♀	A		2

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Hind Foot = 10.3 cm
 EAR = 4.9 cm
 Photos 1-8 Slide 2

Kimberley Mammals -- RECORDS

TW

Team (AB/TH) (NM/TH) (TS/AT)				Location FOREST CREEK, D.R.N.P.								
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
91	<i>Zyzomys argurus</i>	9/6/04	ME	released	FC05 x2 tail	36 gm						5
92	<i>Zyzomys argurus</i>	9/6/04	ME	released	FC06 x2 tail	48 gm				A		2
93	<i>Miniopterus schreibersii</i>	"	Mist net	MS2716	—	11			♂	A		3
94	" "	"	"	MS2717	—	10.7			♂	A		3
95	<i>Nyctophilus walkeri</i>	"	"	MS2718	—	5.4			♂	A		3
96	" "	"	"	MS2719	—	4.4			♀	A		3
97	<i>Myotis molucellarum</i>	"	"	MS2720	—	9.6			♂	A		3
98	<i>Nyctophilus 2 bifax</i>	"	"	MS2721	—	8.6			♀	A		3
99	<i>Zyzomys argurus</i>	10/16/04	ME	MS2722	FC07 x2 toe	4.4			♂	A		5
100	<i>Pseudomys sp</i>	"	ME	MS2723	FC08 x2 toe	11			?	?		3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlit (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- Summary of Spotlight Effort

Team	INS/AT	Location	Geikie
------	--------	----------	--------

Date	21/5/04	Time started	6.50	Time finished	7.10	Total time	20 min
Temperature	Warm	Wind	—	Cloud	—	Moon	—
Trap line (name or similar habitat)	Riparian	% total time	100	Mammals seen (Species and number)			
				Agile Wallaby x 4 (Barking Owl 1)			

- Light deer

Date	22/05/04	Time started	6.40	Time finished		Total time	8.00
Temperature	Warm + Humid	Wind		Cloud	8/8	Moon	Nil
Trap line (name or similar habitat)		% total time		Mammals seen (Species and number)			
"Raft Walk" Limestone				Euro x 1 Limestone			
Riparian				Agile x 3 Riparian			
				Hippoboscids ater (?) several			
7.30 Rain - shelter				heard			
7.40 Resume				H. ater - caught in GWR abutment Block			
				Rain overnight			

Date		Time started		Time finished		Total time	
Temperature		Wind		Cloud		Moon	
Trap line (name or similar habitat)		% total time		Mammals seen (Species and number)			

Date		Time started		Time finished		Total time	
Temperature		Wind		Cloud		Moon	
Trap line (name or similar habitat)		% total time		Mammals seen (Species and number)			

Date		Time started		Time finished		Total time	
Temperature		Wind		Cloud		Moon	
Trap line (name or similar habitat)		% total time		Mammals seen (Species and number)			

Kimberley Mammals -- Summary of Spotlight Effort

Team TS / AT	Location McSherry Gap
---------------------	------------------------------

Date 26/5/04	Time started 1830	Time finished 1900	Total time 30 min
Temperature Warm	Wind gentle	Cloud Nil	Moon 1/8
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
16 Crack at Camp	80%	Nil - Battery Died.	

Date 27/5/04.	Time started 1830	Time finished 2100	Total time 2 1/2 hrs
Temperature Warm	Wind gentle - moderate	Cloud Nil	Moon 1/4
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
Fairfield Leopold Rd	100	Nocturnal Wallaby x 2	
West towards		Cat x 6 - shot 2	
Ward		Wil Bats	
10.5 km		Immature Night Heron	
Vehicle		Bush Thick knee x 4	
		Boobah x 2	

[illegible][illegible][illegible]

Kimberley Mammals -- Summary of Spotlight Effort

Team	TS/AT		Location	SILENT CROVE	
Date	31/05/04		Time started	1945	
Temperature	mild		Wind	Light	
			Cloud	100%	
			Moon	3/4 obscured by cloud	
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)			
SILENT CROVE ENTRANCE	100%	Onychomys leucogaster x1			
Road from		Dingo x3 Spotted Nighthawk			
Compound south to		Cat x1 Outlet Nightjar			
Swish Xing at 12.7 km.		Barn Owl x1 Bush Thickknee			
		Baboon x6 Cattle			
		Tawny Frogmouth x1			

Date 2/26/04	Time started 1020	Time finished 2110	Total time 1:30
Temperature 69°f	Wind W.1	Cloud W.1	Moon Full
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
Silt Grove Composite		Mr. agilis x 4.	
2, Bell Creek Gorge	100%		
Corroch ~ 12 Km		Barking Owl	
		Beech	
		Barn Owl	
		Spotted Nightjar - perched on branch	

[illegible][illegible][illegible]

Kimberley Mammals -- Summary of Spotlight Effort

Team	TS NT.
Location	Solea falls.

Date <u>7/6/04</u>	Time started <u>6:40</u>	Time finished <u>8:10</u>	Total time <u>1.5 hr.</u>
Temperature <u>mild</u>	Wind <u>S light</u>	Cloud <u>NIL</u>	Moon <u>NIL</u>
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
<u>B and 4</u>	<u>100%</u>	<u>NIL mammals</u>	
<u>river portion</u>		<u>1 Frog</u>	
		<u>2 Geckos</u>	
		<u>1 small Croc</u>	
		<u>Few Bats - recording</u>	
		<u>Few spiders!</u>	

Date 8/6	Time started 1845	Time finished 2015	Total time 1:30
Temperature mild	Wind SW	Cloud Nil	Moon Nil
Trap line (name or similar habitat)	% total time	Mammals seen (Species and number)	
Comp 5	40	Hydromys Chrysogaster x 1.	
Line 5			
Soleo Falls 5	50	"Smallest" animal at top of Soleo Falls near	
Comp.	10	River Snail Feeding Station	
		1x Childrens Python	

[illegible][illegible][illegible]

Kimberley Mammals -- Traps

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Gcirkie		Trapline name and/or code Line 1 Over ridge		
Lat 18 06 23	Date traps set 21/5/04	Cages	Big Elliott	med Elliott		
Long 125 41 25	Date traps lifted 25/5/04	8	11	19		
Topography and aspect Heavily dissected S face of limestone range						
Rock type, nature and extent of exposure Limestone Large boulders on heavily isolated/eroded bed rock >90%						
Soil type, origin and depth Skeletal stony clay loam						
						Sample Y/N Y
VEGETATION (Muir system over page) LBr (mixed) / SBr mixed - SD mixed / Hr patches						
Main trees Various PTO						
Main Shrubs "						
Main grasses Tnodia - 2 spp						
Others See over						
Vines Parryflowia Tinospora Sarcostema See over						
Main grasses form & height	hummock ✓	Perennial tussock	Perennial mats	Annual		
FIRE	When was last fire	2003	2002	< 3 yrs	3-10 yrs	>10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Mature Hummock grass patches between bare rock. Veg 1p ind Celtis Sarcostema Gleditsia - patchy charcoal occasionally seen. Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) See above and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Large areas bare rock						
OTHER DISTURBERS (e.g. cattle, pigs, weeds) Some (1-2) parryflowia etc Basically pristine						
HABITAT CHANGE		Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage				
NA.						
PTO for more notes space		Photos Y/N Y		Photo re-takes Y/N		

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Coc fruz
 Cle tom
 Cle flo
 1 Santalum lance
 6 Ficus (ham) MS 116
 1 Dodonaea (feath)
 Cynanchum floumbordon
 1 Tri zey
 Ter sphadleyana
 Tin mi
 1 Tri spp ANS 16
 Mubia maplers
 1 aj cinerous P21 MS
 Baucon
 Sar aus
 Sol las
 1 ap climb lasyanka
 Jas del
 Roskellul
 Hypoestes
 Grewia ber.
 Phytret
 Atahem
 Par Joe
 Lop gra (rare)
 1 Setaria dielsii 1612
 1 Pteris capitata Philobus P21
 1 Ipo eri
 Abu and
 Cuc mel
 ? Lenchus sp ANS 1599
 1 Dichopora
 1 Hyb aur
 1 Ipo ni 1614
 1 poly 399
 1 Trumfetta sp
 1 Corchorus albidus 1610
 1 Tridax purgens 1607
 1 Croton tom

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location GETKIE	Trapline name and/or code 2 Ridge edge		
Lat 18°06'31" - 37	Date traps set 21/5/04	Cages	Big Elliott	med Elliott	
Long 125°41'21"-04	Date traps lifted 25/5/04	8	11	19	
Topography and aspect Flat with occasional limestone mss. Line runs E/W along ecotone of range foot and alluvial flat behind levee					
Rock type, nature and extent of exposure Limestone - bedrock flats to gently slopes but adjacent to range. 30/70% exposure varied.					
Soil type, origin and depth skeletal stony clay loam to deep clay loam Colluvial / alluvial - former more stony					
Sample Y/N Y					
VEGETATION (Muir system over page) LBr (rare < 0.1%) / TL (TL-Jr) or HI					
Main trees Bauhinia lophostemon					
Main Shrubs					
Main grasses Tridax pungens + Tridax sp					
Others					
Vines - rare					
Main grasses form & height	hummock ☒	Perennial tussock	Perennial mats	Annual	
FIRE					
When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Massive hummocks < 1.5m tall - sparse herb cover generally (low flammability) + rare charring					
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) N/A					
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Sparse herbs in most areas stony range to N & road / riparian veg to south					
OTHER DISTURBERS (e.g. cattle, pigs weeds) Pigs Weeds / Pamphora not dominant - Melochia in herb fields. Pigs tracks widespread but not major factor					
HABITAT CHANGE					
Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage ?					
PTO for more notes space			Photos Y/N Y	Photo re-takes Y/N N	

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

pro dia

Additional Notes

* Baucon Buchnera asperata
 * Euc mic Cro med
 Lop gia Cro ret
 Pas fac Tri zey
 Malv ame Nbu and
 Cardiasperme Hib pan
 1 Rostellul Hib spp
 2 Melochia art Cucumis melo
 Phymad. Ste gros
 Port fil Cymbop
 1 Ptil. cap Phypalus
 Ptd. - gae Eup hir
 Hyp enn Opilina aeg
 Poa ann Sida sp
 Gre pyr. Comelinc ens
 Ind lineae Stylos. ham
 Ind may
 Ind collut.

Luffa
 Tradia ~~pro dia~~ 1605
 Erag of tenenull
 Boerh. stenk
 Bid bipen
 Cardiosperme
 Ipomeno P 21
 Didiptia
 Hybaur
 Philotes Josifamix 1601

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location <i>Geikie</i>	Trapline name and/or code <i>3 (Creek in Range)</i>		
Lat <i>18 06 31 - 16</i>	Date traps set <i>21-5-04</i>	Cages	Big Elliott	med Elliott	
Long <i>125 41 36 - 27</i>	Date traps lifted <i>25-5-04</i>	<i>8</i>	<i>11</i>	<i>19</i>	
Topography and aspect <i>Creek line occupying 1/3 - 1/2 of gorge</i>					
Rock type, nature and extent of exposure <i>Limestone cliffs with steep lateral valley occasional - filled c Boulders 50%</i>					
Soil type, origin and depth <i>Heavy clay in valley bottom</i>					
Sample Y/N <i>Y</i>					
VEGETATION (Muir system over page) <i>LAc / G7c - patches & Jc - r patches</i>					
Main trees <i>Emicrotheca Tplatyphylla Barringtonia (Nuclea M leucadendron) Lophostemon</i>					
Main Shrubs <i>Phyllanthus reticulatus</i>					
Main grasses <i>? Tussock - sterile Sedge < 0.5m</i>					
Others <i>Trichodesma Melochia Phyllanthus mad.</i>					
Vines <i>Paniflora - common but not dominant.</i>					
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual	
		<i>✓</i>			
FIRE	When was last fire	1993	1992	< 3 yrs	> 10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) <i>Fire may come down lateral gorges rarely. - contain Celtis etc</i>					
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) <i>see above & over</i>					
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) <i>Gorge walls Flood prone valley floor with bare ground & herbs in most areas</i>					
OTHER DISTURBERS (e.g. cattle, pigs weeds) <i>Pigs. Not major prob Paniflora Melochia</i>					
HABITAT CHANGE		Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage <i>N/A</i>			
PTO for more notes space		Photos Y/N <i>Y</i>		Photo re-takes Y/N <i>N</i>	

Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1 0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

* Euc mie
 Nauclea
 Mel au.
 Barringtonia
 Ter platy
 Lop gie
 Tur goy
 * Aitahens
 Alternanthera
 Amaranthus
 Aechyranthes
 Commens
 Solanum sp
 Malvastrum
 Passiflora
 Cardiospermum
 Phyl med
 Ph. l. color
 Rostellularia
 Eup hirta
 Melochia
 Santalum
 Abou and
 Grewia over → side gorge
 Celtis — " "
 Cro blue
 Watth ind
 Grass 1604
 1605
 Sedge 1606
 Ludwigia per 1602
 Amman auric 1603

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Geikie	Trapline name and/or code Line 4 leve bank		
Lat 18 06 48 - 38	Date traps set 21/5/04	Cages 8	Big Elliott 11	med Elliott 19	
Long 125 41 - 12 - 28	Date traps lifted 25/5/04				
Topography and aspect Levebank between River & Line 3 / Range runs EW					
Rock type, nature and extent of exposure NIL					
Soil type, origin and depth deep heavy alluvial clay loam					
Sample Y/N X					
VEGETATION (Muir system over page) LBr / SA + SB r / GLL - r & Jr					
Main trees <i>Banksia</i> (<i>E microtheca</i>)					
Main Shrubs <i>Santalum</i> <i>Crotalaria</i> <i>Cunning</i>					
Main grasses <i>Urochloa</i> <i>poligera</i> & <i>pubescens</i>					
Others See over					
Vines <i>Panyflora</i>					
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual ✓	
FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) low fuel except panyflora Mature <i>Santalum</i> & oth shrubs little charcoal <i>Banksia</i> without recent epicormic etc or					
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) See over					
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) low fuel zones to N of Line 2) & S (riparian) - low fuel in site					
OTHER DISTURBERS (e.g. cattle, pigs weeds) <i>Panyflora</i> Gall Dominant throughout Pigs oth weeds eg Gallon's Curse					
HABITAT CHANGE		Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage			
PTO for more notes space		Photos Y/N X		Photo re-takes Y/N N	

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

* Santalum
 * Bauhinia
 * Phyllanth
 Phyl. mad.
 Grewia brev.
 'Rope'
 * Paspale
 Celtis
 * Ach. asp.
 * Bidens
 Commens
 Cro. nov.
 * Cro. conn.
 * Nerve
 Hib. merawk
 151600 Hib. penten.
 Malv.
 Nalt. ind.
 Abu and

* Nt. a hem
 * Calatropus
 Euphorbia
 Ceanoth.
 Ceanoth.
 Enneap. coe.
 Euc. mic.
 Euc. can.
 Hyb. enn.
 Reichen. of holes.
 Selanc. apic.
 T. modic.
 Portul. fil.
 Sporob. abl. P21

Uropob P21
 Propel P21
 Cro. ret.
 Ipo. enu.
 Hah. arbor.
 Opil. aeg.
 Eudorolus als.
 Aca. farnes.
 "Marsdenia"

Kimberley Mammals -- Traps

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Gerbie	Trapline name and/or code 5 - Riparian.
--	--------------------	--

Lat 18 06 43 (Wend)	Date traps set 21/5/04	Cages 8	Big Elliott 4	med Elliott 14
Long 125 41 22 (Wend)	Date traps lifted 25/5/04			

Topography and aspect	Alluvial Terrace on N bank of Fitzroy
-----------------------	---------------------------------------

Rock type, nature and extent of exposure	NIL
--	-----

Soil type, origin and depth	Mundial Clay/Loam	Sample Y/N	Y
-----------------------------	-------------------	------------	---

VEGETATION (Muir system over page) ML to MC / Vines & patches of SD				
Main trees E Microtheca + he over				
Main Shrubs See over				
Main grasses — Urochloa pelagica & D pob rarer				
Others Herbs Tuckersma				
Vines Pandanora				
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Massive Pandanora mantles in tall trees Deep litter layers

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Riparian trees eg Fraxinus, F. coronata, Nauclea

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

River to South. low fuel to N (see Line 2 & 4)

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

Pandanora - Dominates 80/90% over throughout; Pigs. many areas of "rooted" soil & tracks - ground cover & climber canopy to canopy

HABITAT CHANGE

Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N X

Photo re-takes Y/N

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

* Das Joe
* Euc mic
* Fracemose
F. coronata
Nauale
Barrington
Mel arg
Mel lev.
Atahem
* Tri des
Wal ind
Mal ame
"Marsdeni"

Euphorbia
* Solanum locani
Ab. andrewsii
Achyranthes
M. 1605
Heliosp. ind.
Mib meracensis
Calotropis
Cen echinatus
Cardios perma

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location McSherry Gap	Trapline name and/or code 1A Camp Hillside 1B Camp
--	--------------------------	--

Loop

Lat 17 43 19"	Date traps set 25/5/04	Cages	Big Elliott	med Elliott
Long 125 06 42	Date traps lifted 29/5/04	4	5	9

Topography and aspect	Lower slope of small elongate Hill
-----------------------	------------------------------------

Rock type, nature and extent of exposure	Phyllite bedrock, low exposure slope Mulch on ground surface $\leq 80\%$
--	---

Soil type, origin and depth	Skeletal loam derived locally from Phyllite rock	Sample Y/N	yes
-----------------------------	---	------------	-----


VEGETATION (Muir system over page) LBr / HL				
Main trees Eucalyptus with Eucalyptus Boab at foot of hill				
Main Shrubs				
Main grasses Coily Sprinter ?T. Litexture				
Others				
Vines				
Main grasses form & height	hummock ☒	Perennial tussock	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)
Trochilid flowering but small. Eucalyptus grown out.
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))
NIL
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)
NIL besides creek to West

OTHER DISTURBERS (e.g. cattle, pigs weeds)
Cattle minor - mainly bare weeds minor PTO

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
	?
PTO for more notes space	Photos Y/N ☒ Photo re-takes Y/N



Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% t
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Euc 399

Euc tec - base of hill

Bocob -

Sorghum stipoides - densest at base of hill - but flowering here

Colpaea

Dicliptera

Helolus pusiformis

Crotalaria montana

Buchnera

Crotalaria medicaginea

Trichodesmos

Cochlospermum parvifolium

Terminalia catappa

Cardinalis flaccida

Goodenia

Travenola

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <i>McSherry Gap</i>	Trapline name and/or code <i>1B Camp Creek</i>
--	---------------------------------	---

Lat <i>17 43 19</i>	Date traps set <i>25/5/04</i>	Cages <i>4</i>	Big Elliott <i>6</i>	med Elliott <i>10</i>
Long <i>125 06 42</i>	Date traps lifted <i>29/5/04</i>			

Topography and aspect	<i>Creek incised into alluvial plain</i>
-----------------------	--

Rock type, nature and extent of exposure	<i>NA</i>
--	-----------

Soil type, origin and depth	<i>Alluvial clay loam with sandy patches deep</i>	Sample Y/N	<i>Y</i>
-----------------------------	---	------------	----------

VEGETATION (Muir system over page) <i>LAC-i / SAC-r / GTd</i>				
Main trees	<i>Eucalyptus Terminalia platyphylla Lophoceros Basmia</i>			
Main Shrubs	<i>Phyllanthus reticulatus & sapling Terminalia platyphylla</i>			
Main grasses	<i>Heteropogon Schima "Blady" ANS 1689</i>			
Others				
Vines	<i>Panicum</i>			
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	<u>3-10 yrs</u>	>10 yrs
------	--------------------	------	------	---------	-----------------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)	<i>Dense tree canopy - old shrubs - creek bed</i>
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))	<i>Riparian sp - see list over</i>
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)	<i>Creek bed tree canopy</i>

OTHER DISTURBERS (e.g. cattle, pigs, weeds)	<i>Cattle some pigs some weeds - mostly panicum - not major problem</i>
---	---

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
<i>?</i>	
PTO for more notes space	Photos Y/N <i>Y</i> Photo re-takes Y/N

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% t
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Ludwigia peruviana
Amm. 
Ipomoea  *erio.*
Hibiscus mevarianus
H. pentaphylla
Trichodesma
Malaya
Phytolobus
Eubalia aurea
Vigna
Pariflora
Nelsonia
Boab.
Crot. medica
Crot. retusa
Crot. montana
Chloris
Melochia
Sida acuta

"Blady" 1639
Tas. mol.
Ehretia
Hydrophylla ang.
Basilica
Grew. retusa
Premna
Ficus coronata
F. opposita
Sedge - Geelba 1606
Hakea arborescens
Rhynchos
Erythrina

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (GP +)		Location <i>McSherry Gap</i>		Trapline name and/or code <i>2 old Spempr.</i>		
Lat <i>17 34 08</i>	Date traps set <i>25/5/04</i>	Cages	Big Elliott	med Elliott		
Long <i>125 06 52</i>	Date traps lifted <i>29/5/04</i>	<i>8</i>	<i>11</i>	<i>19</i>		
Topography and aspect <i>gullies</i> <i>gently undulating plain incised by</i>						
Rock type, nature and extent of exposure <i>Granodiorite < 5% exposed</i> <i>Surface stone mulch</i>						
Soil type, origin and depth <i>probably colluvial with high clay sand</i> <i>& stone content</i>						
Sample Y/N ☒						
VEGETATION (Muir system over page) <i>LBr / Hctotti or GTe-r or GLi</i>						
Main trees <i>Eucalyptus Hakea arborescens</i>						
Main Shrubs <i>/</i>						
Main grasses <i>Triodia NW 1624 or Heteropogon Sorghum styriac. or Xerochloa</i>						
Others <i>/</i>						
Vines <i>/</i>						
Main grasses form & height	hummock ☒	Perennial tussock ☒	Perennial mats	Annual ☒		
FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	
> 10 yrs						
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) <i>Large old triodic hummocks - some scarring</i>						
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) <i>NA</i>						
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) <i>NA</i>						
OTHER DISTURBERS (e.g. cattle, pigs weeds) <i>MINOR - appears to have been severely</i> <i>scalded/eroded in past - now regenerating.</i>						
HABITAT CHANGE		Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage <i>? Gullying & loss of A' horizon</i>				
PTO for more notes space		Photos Y/N ☒		Photo re-takes Y/N		

NW 1623

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Phytolacca exaltata
M. capitatus
Banksia
Boob
Panicum decomf.
Chrysopogon fal.
Xerodhloa
Dicelyptera
Coomed
Bochnera
Vigna
Rhynchosia
Trichadenia

Wue Enodine
Cymbopogon sp.
Euphorbia sp. *steub.*
Stemodia ~~sp.~~ *rupestris*
Ehretia
Grassia retusifolia *und* *Boob.*
Bra div
Crotalaria montana
Phyl. reticulata - *und* *Boob*

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location McSherry Gap	Trapline name and/or code 3 - Reef Limestone
--	--------------------------	---

Lat 17 33 49	Date traps set 28/5/04	Cages 8	Big Elliott 11	med Elliott 19
Long 125 06 14	Date traps lifted 29/5/04			

Topography and aspect

Rock type, nature and extent of exposure

80%
surface boulder Limestone bedrock with some
leaves & soil mulch.

Soil type, origin and depth

Skeletal in situ

Sample Y/N ☒

VEGETATION (Muir system over page)

LB r / HL

Main trees

Cochlospermum Boob.

Main Shrubs

Various

Main grasses

Tridax sp. sterile (= NM 1624?)

Others

Many more

Vines

minor

Main grasses
form & height

hummock ☒

Perennial tussock

Perennial mats

Annual

☒

FIRE

When was last fire

1993

1992

< 3 yrs

3-10 yrs ☒

>10 yrs

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)

Age of Tridax

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

Ficus in valleys on rocky sites
Adenium

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

Rock - including bedrock on creek / river
in valley bottoms

1/2 page

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

HABITAT CHANGE

Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage

PTO for more notes space

Photos Y/N

☒

Photo re-takes Y/N

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% t
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Grevillea pyramidalis
Senna venusta
Diactyone
Trichoderm
Portul. filiformis
Conchocarp of *Sida*
Trumflettia sp.
Terminalia hadleyana
Euphorbia sp. herb.
Hibiscus 11/1619
Philolus fruticosus
P. gardneri
P. 11/1637
P. capitatus
Nerolana
Buckner
Ficus hampdenii pod.
F. oppositifolia

Stemodia viscosa
Weddellia 11/1638
Solanum lasiopetalum
Tephrosia rosea
Parataphrosia
Parafloerfeldia
Sarcostema
Tournefortia did.
Timospora
Phyll. reticulata
Adenia
Gom. canescens
Dodonaea polyzyga
Agrostis
Lycopodium

Lindernia 1632
Sedge 1633
Lophostemum
Opil. aeg.
Cynanchum
Santalin
Racewood
Sporobolus 1635

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location <i>McSherry Gap</i>		Trapline name and/or code <i>4 Blackrod Paddock</i>		
Lat <i>17 34 01 +20</i>	Date traps set <i>25/5/04</i>	Cages	Big Elliott	med Elliott		
Long <i>125 05 47-54</i>	Date traps lifted <i>29/5/04</i>	<i>8</i>	<i>11</i>	<i>19</i>		
Topography and aspect <i>Flat</i>						
Rock type, nature and extent of exposure <i>NA</i>						
Soil type, origin and depth <i>Deep cracking Black soil</i>					Sample Y/N <i>Y</i>	
VEGETATION (Muir system over page) <i>Jr / GTa - e</i>						
Main trees <i>(Bauhinia)</i>						
Main Shrubs						
Main grasses <i>Schuma (Sorghum hybrid in places)</i>						
Others <i>Sesbania simpliciflora</i>						
Vines						
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual		
		☒	☐			
FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	
					>10 yrs	
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) <i>?</i>						
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) <i>NIL</i>						
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) <i>/</i>						
OTHER DISTURBERS (e.g. cattle, pigs, weeds)						
<i>Cattle few weeds</i>						
HABITAT CHANGE		Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage <i>?</i>				
PTO for more notes space		Photos Y/N <i>Y</i>		Photo re-takes <i>Y/N</i>		

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Song-stipoid
 Rivinida perenne
 Panner decomp
 Weddellia 1630 asperna
 Sporob
 Buchner
 Rhynchosia
 Comelina
 Ipomoea
 Carina
 Eragrostis 1631

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location <i>McSherry Gap</i>	Trapline name and/or code <i>5 McSherry Gap n.p.m.</i>		
Lat <i>17 33 49</i>	Date traps set <i>25/5/04</i>	Cages <i>8</i>	Big Elliott <i>11</i>	med Elliott <i>19</i>	
Long <i>125 05 58</i>	Date traps lifted <i>29/5/04</i>				
Topography and aspect <i>Incised creek & alluvial flat occupy gap in limestone range</i>					
Rock type, nature and extent of exposure <i>NA</i>					
Soil type, origin and depth <i>deep alluvial clay loam</i>					
Sample Y/N <i>Y</i>					
VEGETATION (Muir system over page) <i>LAC-LAr / SL-Sr / GTC-L & Jr</i>					
Main trees <i>Euc cam Term. platyph. Euc tect. Boob. Bauhinia lophosperm.</i>					
Main Shrubs <i>Various</i>					
Main grasses <i>Heteropogon Schum. Blady AWS 1639</i>					
Others <i>Various herbs</i>					
Vines <i>Panylore.</i>					
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual	
		☒			
FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs
					>10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) <i>Malve shrubs little scorch.</i>					
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) <i>Multiloes.</i>					
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) <i>Cliffs - Creek bed. Road</i>					
OTHER DISTURBERS (e.g. cattle, pigs weeds) <i>Cattle. Weeds Panylore.</i>					
HABITAT CHANGE		Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage			
PTO for more notes space		Photos Y/N <i>Y</i>		Photo re-takes Y/N	

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Hibiscus merauk
H. pentaphyllus
Cro. medicag.
Cro. refusa
Cro. montana
Cro. crisp.
Chrysopogon
Eragrostis 1631
Vigna
Passiflora
Achyranthes
Ipomoea
Malvastrum
Rope
Alternanthera pungens
Whitechloa
Dicliptera

Blady 115 1639
Commelina
Phyl. reticulata
Ind. colutea
Grewia refusa
Tan. did.
J. mol
Premna
Sylo hamata
Ficus coronata
F. opposita
Eup. hirta
R. wince
Tinospora
Sida acuta
Nelsonia

Calotropis
Planchonia
ammanica

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Silent Grove	Trapline name and/or code 1 - Bell Creek riparian
--	--------------------------	--

Lat 17 31 13.8	Date traps set 31 May 04	Cages 8	Big Elliott 11	med Elliott 19
Long 125 13 53	Date traps lifted 4 June			

Topography and aspect	River levee and edge of flood plain
-----------------------	-------------------------------------

Rock type, nature and extent of exposure	NA
--	----

Soil type, origin and depth	Sandy loam -- more sandy on the levee
	Sample Y/N Y

VEGETATION (Muir system over page) LAL-r / Tr rGTc-i				
Main trees	E camaldulensis & Terminalia platyphylla (M. leucodendron on creek bank)			
Main Shrubs	Bucanania repens & Nauclea plicata			
Main grasses	"Blady" = MNS 1639. Het cont Schima			
Others	See list			
Vines	See list			
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)	Charring & shrub recovery
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))	N/A
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)	Creek to North

OTHER DISTURBERS (e.g. cattle, pigs, weeds)	Cattle Pigs not major Some weeds - not major
---	---

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
	7
PTO for more notes space	Photos Y/N Y Photo re-takes Y/N

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

levee
 Blady
 Buchananii
 Planchone
 Hyptis
 Helicostachya
 Acacia pleurocarpa
 Whitechloe
 Freycinet
 Crot retusa
 Side acute
 Mel. Acrodonia
 Pandanus spiralis
 Euphorbia
 Ne helioser.
 Cro montana
 Cro juncea

Hibiscus - ans
 Grex retusifolia
 Bolbroch.
 Cram ANS
 Flood plain
 Zornia
 Rhyssalis
 Grex retusifolia
 Helicostachya
 Cro montana
 Hyptis
 Eucalypt
 McSherry Grass ANS
 Sorghum
 IPO

Ticus opposita
 Planchone
 Bonamia

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Silent Gorge	Trapline name and/or code Gardner's Gap Line 2		
Lat 17 02' 28"	Date traps set 31/5/04	Cages 8	Big Elliott 11	med Elliott 19	
Long 125 13' 33"	Date traps lifted 4/6/04				
Topography and aspect Gorge with permanent creek flowing North					
Rock type, nature and extent of exposure KL Sandstone 90% Extensive bed rock Outcrop on ridges & scree on slopes - boulders stream creek bed.					
Soil type, origin and depth Humus rich loam - skeletal & deep skeletal below boulders & on scree Sample Y/N Y					
VEGETATION (Muir system over page) LAL-r / S L-r / GTe-r + Jr					
Main trees E miniata and E app - white bark - remnant of fire hole - in creek.					
Main Shrubs Mixed see list					
Main grasses NMS 1639 NMS 1643 See list.					
Others See list } no dominants					
Vines See list }					
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual	
FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Some evidence of fire killed trees etc. but those there now are mature probably not frequent near mouth at N end					
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) Rainforest elements Adiantum Canarium See list					
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Rock faces of gorge & probably permanent, grace grass & creek					
OTHER DISTURBERS (e.g. cattle, pigs, weeds) Pigs - deep rooting near entrance see 30m up side slope some weeds - not dominant					
HABITAT CHANGE		Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage ?			
PTO for more notes space		Photos Y/N X		Photo re-takes Y/N	

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Silent Grove	Trapline name and/or code Minuata Line 3
--	--------------------------	---

Lat 17°03'07"S	Date traps set 31/5/04	Cages 8	Big Elliott 11	med Elliott 14
Long 125 14 11	Date traps lifted 4/6/04			

Topography and aspect	Colluvial slope at foot of sandstone ridge - runs E/W
-----------------------	---

Rock type, nature and extent of exposure	High leopard sandstone < 5% surface matrix of stones & gravel associated & drainage lines.
--	---

Soil type, origin and depth	Sandy loam deep	Sample Y/N Y
-----------------------------	-----------------	--------------

VEGETATION (Muir system over page) LA c-i / GT c-i				
Main trees E Minuata				
Main Shrubs Buchanan's Ironwood & see list.				
Main grasses Sorghum stipoides				
Others Protea - see list				
Vines see list				
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual ✓

FIRE	When was last fire	1993 2003	1992	< 3 yrs	3-10 yrs	> 10 yrs
------	--------------------	-----------	------	---------	----------	----------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)
 Buchanan's & Ironwood etc all kept to < 1m
 by frequent fire Range reports for last year

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))

NIL

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)

NIL.

OTHER DISTURBERS (e.g. cattle, pigs, weeds)

Pigs. + ? cattle - not major.

HABITAT CHANGE

Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage

?

PTO for more notes space

Photos Y/N

Y,

Photo re-takes Y/N

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Silent Grove		Trapline name and/or code Line 4 Rock Ridge		
Lat 17 03 07	Date traps set 31/5/04	Cages	Big Elliott	med Elliott		
Long 125 14 11	Date traps lifted 3/6/04 73 days	8	11	19		
Topography and aspect Sandstone ridge runs EW.						
Rock type, nature and extent of exposure King Leopold ss. 95% Bedrock with boulder scatter - tending to scree on lower slope						
Soil type, origin and depth Skeletal Rich Loam. dark brown.						
Sample Y/N Y						
VEGETATION (Muir system over page) LB, / SA, / GTr + Hr						
Main trees Mixed						
Main Shrubs Mixed						
Main grasses Sorghum stipoides early sprouts						
Others See list						
Vines NA						
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual ✓		
FIRE	When was last fire	1983 2003	1992	< 3 yrs	> 10 yrs	
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) Ranger reported fire last year.						
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) Acacias.						
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Rock. would make it patchy						
OTHER DISTURBERS (e.g. cattle, pigs, weeds) NA						
HABITAT CHANGE Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage ?						
PTO for more notes space		Photos Y/N Y		Photo re-takes Y/N		

Thommo.

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location <i>Silent Grove</i>	Trapline name and/or code <i>Camp line</i>
--	---------------------------------	---

Lat	Date traps set <i>3/6</i>	Cages	Big Elliott	med Elliott
Long	Date traps lifted <i>4/6</i>	<i>10</i>	<i>10</i>	<i>0</i>

Topography and aspect	<i>Creek margin and adjacent flat.</i>
-----------------------	--

Rock type, nature and extent of exposure	<i>NA</i>
--	-----------

Soil type, origin and depth	<i>Petty loam near creek red sandy loam further out</i>	Sample Y/N <i>N</i>
-----------------------------	---	---------------------

VEGETATION (Muir system over page) <i>LAc-i/GTd-c near creek to LBr-GTd-i away from creek</i>				
Main trees	<i>Melaleucadend. near creek. Eucalyptus away.</i>			
Main Shrubs	<i>Few</i>			
Main grasses	<i>Whiteclover near creek. Heteropogon away</i>			
Others	<i>Few</i>			
Vines	<i>Few</i>			
Main grasses form & height	hummock	Perennial tussock ☒	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	-----------------	-----------------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)	<i>Range advice 2003</i>
--	--------------------------

Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))	<i>NA</i>
--	-----------

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)	<i>NA</i>
--	-----------

OTHER DISTURBERS (e.g. cattle, pigs, weeds)	<i>Pigs, weeds, Hyptis.</i>
---	-----------------------------

HABITAT CHANGE	Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage
PTO for more notes space 9	
Photos Y/N <i>N</i>	Photo re-takes Y/N

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Solea Falls	Trapline name and/or code Line 1. Creek West
--	-------------------------	---

Lat 14.40.43	Date traps set 6/6/04	Cages 8	Big Elliott 10	med Elliott 19
Long 126 59 29 Loop	Date traps lifted			

Topography and aspect	Creek line meandering through sandstone boulder country
-----------------------	---

Rock type, nature and extent of exposure	Sandstone 70% Boulders on bedrock
--	-----------------------------------

Soil type, origin and depth	Sand to sandy loam 2 m deep
Sample Y/N	Y

VEGETATION (Muir system over page) (A) Rainforest Sc/GTc (B) Luvetonia LBe-r 1GTc-1 (C) Savanna LBr/GTc				
Main trees	(A) See list (B) Luvetonia (C) Euc sp & Corymba			
Main Shrubs	See list variable			
Main grasses	Sorghum sp. sp.			
Others	Tirodus "Hologonifera"			
Vines	See list			
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual ☒

FIRE	When was last fire	1993	1992-03	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	---------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)
Burnt 2003 - Neaua seedlings < 40 cm etc ? 2001/2 - erect stems charred on wood by 2003 fire.
Patches men fire
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))
Pouter Adenium Ficus etc in a patch on creek.

and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)
Blocky sandstone protects pockets.

OTHER DISTURBERS (e.g. cattle, pigs weeds)
Cattle pads & dung old. No weeds Donkeys - footprints old.

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage ?
PTO for more notes space	Photos Y/N Y Photo re-takes Y/N N

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Solea Falls		Trapline name and/or code Line 2. Brads Sandstone North		
Lat 126 59 25 - 29	Date traps set 6/6/04	Cages	Big Elliott	med Elliott		
Long 14 40 35.4 - 43	Date traps lifted	8	10	19.		
Topography and aspect Sandstone boulder on bedrock broken but flat						
Rock type, nature and extent of exposure See above						
Soil type, origin and depth Skeletal sand.						
						Sample Y/N y
VEGETATION (Muir system over page) LBr/GTc-i						
Main trees <i>Corymbia Owenii</i>						
Main Shrubs <i>Various see list</i>						
Main grasses <i>Sorghum stipoides</i>						
Others <i>Tridax "stolonifera"</i>						
Vines						
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual		
FIRE	When was last fire	1993	1992 03	< 3 yrs	3-10 yrs	>10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) 2003 - mixed patches - over hot. 2001/2 - 2003 fire charred still-end stems killed by an earlier fire Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) N/A and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Bloody sandstone → patches shelter						
OTHER DISTURBERS (e.g. cattle, pigs, weeds)						
Minimal						
HABITAT CHANGE		Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage				
		?				
PTO for more notes space		Photos Y/N y		Photo re-takes Y/N		

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Solea Falls.		Trapline name and/or code Line 3 Sandstone South.		
Lat 14 40 43		Date traps set 6/6/04		Cages 8	Big Elliott 10	med Elliott 19
Long 126 59 29 Loop.		Date traps lifted				
Topography and aspect Blochy sandstone boulders on bedrock						
Rock type, nature and extent of exposure Sandstone 90%						
Soil type, origin and depth Skeletal sand						Sample Y/N Y
VEGETATION (Muir system over page)						
Main trees						
Main Shrubs						
Main grasses						
Others						
Vines						
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual ✓		
FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) See Line 2 - Solea Falls						
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) See Line 2						
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) See line 2						
OTHER DISTURBERS (e.g. cattle, pigs, weeds) See line 2						
HABITAT CHANGE		Record indications of change. E.g. recollections: Dead cypress but no regeneration: rainforest shrinkage ?				
PTO for more notes space			Photos Y/N Y,		Photo re-takes Y/N	

Kimberley Mammals -- Traplines

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location Solea Falls		Trapline name and/or code line 4 River cliffs		
Lat 1440 - 51.5	Date traps set 6/6/04	Cages	Big Elliott	med Elliott		
Long 126 59 - 38	Date traps lifted	8	11	19		
Topography and aspect Cliff line of massive boulders and bedrock along E bank of overflow channel						
Rock type, nature and extent of exposure Sandstone $\approx$ 100%						
Soil type, origin and depth Skeletal loamy sand in crevices						
Sample Y/N ☒						
VEGETATION (Muir system over page) SA _r /H _r						
Main trees ☒						
Main Shrubs Ficus small Rosteric Stychnos Xanth. paradox "coffee"						
Main grasses Phod.						
Others						
Vines Flagellum						
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual		
		☒				
FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
Comment (e.g. indicators used to assess age, longer term effects, intensity etc) No fire signs						
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias)) No fuel						
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc) Rock & river bed.						
OTHER DISTURBERS (e.g. cattle, pigs weeds) NA						
HABITAT CHANGE		Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage NA				
PTO for more notes space		Photos Y/N ☒		Photo re-takes Y/N		


Muir vegetation System

LIFE FORM/HEIGHT CLASS		CANOPY COVER			
		DENSE 70-100% ^d	MID-DENSE 30-70% ^c	SPARSE 10-30% ⁱ	VERY SPARSE 2-10% ^r
T	Trees >30m	Dense Tall Forest	Tall Forest	Tall Woodland	Open Tall Woodland
M	Trees 15-30m	Dense Forest	Forest	Woodland	Open Woodland
LA	Trees 5-15m	Dense Low Forest A	Low Forest A	Low Woodland A	Open Low Woodland A
LB	Trees <5m	Dense Low Forest B	Low Forest B	Low Woodland B	Open Low Woodland B
KT	Mallee tree form	Dense Tree Mallee	Tree Mallee	Open Tree Mallee	Very Open Tree Mallee
KS	Mallee shrub form	Dense Shrub Mallee	Shrub Mallee	Open Shrub Mallee	Very Open Shrub Mallee
S	Shrubs >2m	Dense Thicket	Thicket	Scrub	Open Scrub
SA	Shrubs 1.5-2.0m	Dense Heath A	Heath A	Low Scrub A	Open Low Scrub A
SB	Shrubs 1.0-1.5m	Dense Heath B	Heath B	Low Scrub B	Open Low Scrub B
SC	Shrubs 0.5-1.0m	Dense Low Heath C	Low Heath C	Dwarf Scrub C	Open Dwarf Scrub C
SD	Shrubs 0.0-0.5m	Dense Low Heath D	Low Heath D	Dwarf Scrub D	Open Dwarf Scrub D
P	Mat plants	Dense Mat Plants	Mat Plants	Open Mat Plants	Very Open Mat Plants
H	Hummock Grass	Dense Hummock Grass	Mid-Dense Hummock Grass	Hummock Grass	Open Hummock Grass
GT	Bunch grass >0.5m	Dense Tall Grass	Tall Grass	Open Tall Grass	Very Open Tall Grass
GL	Bunch grass <0.5m	Dense Low Grass	Low Grass	Open Low Grass	Very Open Low Grass
J	Herbaceous spp.	Dense Herbs	Herbs	Open Herbs	Very Open Herbs
VT	Sedges >0.5m	Dense Tall Sedges	Tall Sedges	Open Tall Sedges	Very Open Tall Sedges
VL	Sedges <0.5m	Dense Low Sedges	Low Sedges	Open Low Sedges	Very Open Low Sedges
X	Ferns	Dense Ferns	Ferns	Open Ferns	Very Open Ferns
	Mosses, liverwort	Dense Mosses	Mosses	Open Mosses	Very Open Mosses

Additional Notes

Stychnos
Smooth fig
F. opposita
Pouteric ser.
Xanthost. paradoxus
Flagellum
Vitex
"Coffee"
Junc. triodic
Calytrix brownii
Gardenia 'B'

Kimberley Mammals -- Traplines



37

Team (AB/TH) (NM/JR) (TS/AT) (CP +)	Location Solea Falls	Trapline name and/or code line 5 'Island'
--	-------------------------	--

Lat 14.40.36	Date traps set 6/6/04	Cages 8	Big Elliott 11	med Elliott 19
Long 126.59.38	Date traps lifted			

Reduced
4xc 2mgh
3xLE
6xME

Topography and aspect	"Island" of sandstone between channel & side channel
-----------------------	--

New line
Falls 5a

Rock type, nature and extent of exposure	Sandstone 60%+ Blocks on bedrock
--	----------------------------------

4xc 2mgh
8xLE
13xME

Soil type, origin and depth	Skeletal Sand
Sample Y/N Y	

VEGETATION (Muir system over page) S _c -r / H _c				
Main trees				
Main Shrubs Ac. kelleri Gandemia 'B', Calytrix brownii A. donnellii				
Main grasses Proelia ANS r Proelia ANS				
Others				
Vines N. denhamii & T. mosopora				
Main grasses form & height	hummock	Perennial tussock	Perennial mats	Annual

FIRE	When was last fire	1993	1992	< 3 yrs	3-10 yrs	>10 yrs
------	--------------------	------	------	---------	----------	---------

Comment (e.g. indicators used to assess age, longer term effects, intensity etc)
Appears long unburnt. Old grasses. Mature - thick - no charring
Notes on presence and status of fire sensitive spp (e.g. Cypress, rainforest species, obligate seeders (including most Acacias))
See list
and features offering fire shelter (e.g. rock formations, creeks, vegetation type etc)
River r off shoot make this a fire-safe island

OTHER DISTURBERS (e.g. cattle, pigs weeds)
NA

HABITAT CHANGE	Record indications of change. E.g recollections: Dead cypress but no regeneration: rainforest shrinkage
	NIL?
PTO for more notes space	Photos Y/N Y
	Photo re-takes Y/N

Muir vegetation System

LIFE FORM/HEIGHT CLASS	CANOPY COVER			
	DENSE 70-100% d	MID-DENSE 30-70% c	SPARSE 10-30% i	VERY SPARSE 2-10% r
T Trees >30m M Trees 15-30m LA Trees 5-15m LB Trees <5m	Dense Tall Forest Dense Forest Dense Low Forest A Dense Low Forest B	Tall Forest Forest Low Forest A Low Forest B	Tall Woodland Woodland Low Woodland A Low Woodland B	Open Tall Woodland Open Woodland Open Low Woodland A Open Low Woodland B
KT Mallee tree form KS Mallee shrub form	Dense Tree Mallee Dense Shrub Mallee	Tree Mallee Shrub Mallee	Open Tree Mallee Open Shrub Mallee	Very Open Tree Mallee Very Open Shrub Mallee
S Shrubs >2m SA Shrubs 1.5-2.0m SB Shrubs 1.0-1.5m SC Shrubs 0.5-1.0m SD Shrubs 0.0-0.5m	Dense Thicket Dense Heath A Dense Heath B Dense Low Heath C Dense Low Heath D	Thicket Heath A Heath B Low Heath C Low Heath D	Scrub Low Scrub A Low Scrub B Dwarf Scrub C Dwarf Scrub D	Open Scrub Open Low Scrub A Open Low Scrub B Open Dwarf Scrub C Open Dwarf Scrub D
P Mat plants H Hummock Grass GT Bunch grass >0.5m GL Bunch grass <0.5m J Herbaceous spp.	Dense Mat Plants Dense Hummock Grass Dense Tall Grass Dense Low Grass Dense Herbs	Mat Plants Mid-Dense Hummock Grass Tall Grass Low Grass Herbs	Open Mat Plants Hummock Grass Open Tall Grass Open Low Grass Open Herbs	Very Open Mat Plants Open Hummock Grass Very Open Tall Grass Very Open Low Grass Very Open Herbs
VT Sedges >0.5m VL Sedges <0.5m	Dense Tall Sedges Dense Low Sedges	Tall Sedges Low Sedges	Open Tall Sedges Open Low Sedges	Very Open Tall Sedges Very Open Low Sedges
X Ferns Mosses, liverwort	Dense Ferns Dense Mosses	Ferns Mosses	Open Ferns Open Mosses	Very Open Ferns Very Open Mosses

Additional Notes

Trodia reungus NNS
Trodia sp. NNS
Grevillea agnifolia
Gardenia B - fruit in purple box.
Ac. kelleni NNS 1651
Ac. aff. translocens NNS. 1652
Tachyoma argent NNS 1653
Hebe NNS 1650.
Calytrix brownii
 exspp.
Verticordia
Isaemum NNS -
Sorghum stipoid
Neacume
Eragrostis sp.
Vitex
Santalum

Peisocoma
Boronia NNS.
Neacume
Ficus opposita
Tinospora
Owenia venicosa
Buchanania
Xanthost. paradoxus

Kimberley Mammals -- RECORDS

- Lines
1. Ridge
 2. Ridge edge
 3. ridge creek
 - 4 riparian edge
 - 5 riparian river.

Team (AB/TH) (NM/JR) (TS/AT)				Location <i>Gerbie Gorge</i>								
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
1	<i>Euro</i>	21/5/04	F	M54671	/	/	/	/	/	Ad	/	Ridge. (1)
2	<i>Mac. agiles</i>	21/5/04	SD	S	/	/	/	/	♂♀	Ad ju	✓	Ridge edge (2)
3	"	"	"	"	/	/	/	/	"	"	✓	ridge creek 3
4	"	"	"	"	/	/	/	/	"	"	/	riparian edge 4
5	"	"	"	"	/	/	/	/	"	"	✓	riparian 5
06	<i>Euro.</i>	"	"	"	/	/	/	/	"	"	—	Ridge 1
07	<i>Pig.</i>	20/5/04	D	S	/	/	/	/	?	Ad.	—	4 2 3 5
08	<i>Nailtail</i>	20/5/04	D	S	/	/	/	/	/	Ad	—	4
09	<i>Ghost Bat</i>	21/5/04	O Pro	Head	/	/	/	/	/	/	—	5
10	<i>Planigale</i>	22/5	SE	54676	/	chewed out of bag	/	/	♂	Ad	—	1

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	Agile Wallabies very common throughout riparian area many seen.
3	
4	
5	
6	
7	Much pig rooting in 5 (river) some in 4 & 3 & 2
8	
9	Ghost bats squeaky twitters heard in trees over camp throughout night but not continuous
0	

151
75

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>Gerike</i>							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
11	<i>Z. argurus</i>	22/5	SE	R	M54672 MU WAM EP	34	-	-	♀	ad	NLNP	3
12	"	"	"	R	M54673 MU WAM EP	37	-	-	♂	ad	-	3
13	<i>Mus</i>	"	"	M54674	MU	13.5	76	75	♂	ad	-	3
14	<i>Z. argurus</i>	"	SE	M54675	MU WAM EP	39	108	96	♂	ad	Notes	1
15	Blank		SE	M54676					♂	ad		1
16	<i>Rattus rattus</i>	"	SE	M54677	MU WAM EP	125			♂	ad		5
17	<i>Tachyglossus aculeatus</i>	"	O	Scot								1
18	<i>P. nanus</i>	23/5	SE	M54678	WAM Tissue MU Tissue	28	93	96	♀	Ad	Recently coupled	2
19	<i>P. nanus</i>	"	SE									2
20	<i>Z. argurus</i>		SE	R		35			♀	Ad	NLNP	1

~~35~~
no tail.

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlit (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	Testes partly descended. Hindfoot 21 Ear 20 Lighter colour on dorsal surfaces than those seen in North Kimber 2003 and East Kimberley
5	
6	
7	
8	No ectoparasites on animal CALM Hair Sample Reproduction: - blood on entrance to vent - vent - nipples just showing
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location <i>GEIKIE GORGE NP</i>							
Record Number	Foot notes Species	Date	*1 Source	*2 Record type	*3 Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	*4 Trapline
21	<i>Zy. argurus</i>	23/5/04	ME	R		28			♂		NLNP	1
22	<i>Zy. argurus</i>	"	ME	R		30			♀		NLNP Nipples recessed	2
23	<i>Zy. argurus</i>	"	ME	R		21			♂	SAD		3
24	<i>Mus</i>		ME	R		-			-			4
25	<i>Mus</i>		ME	R		-			-			4
26	<i>Mus</i>	23/5/04	LE	R		-			-			4
27	<i>Tachyglossus aculeatus</i>	23/5/04	CO	Scat								2
28	<i>Zy. argurus</i>	24/5/04	ME	R		31			escape			1
29	<i>Zy. argurus</i>	24/5/04	ME	R		25			♂	Juv		1
30	<i>Zy. argurus</i>	24/5/04	ME	photo		76						1

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

"Red"

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location GIEKIE GORGE NP							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
31	Zy argurus	24/5/04	LE	photo								2
32	Mus	"	ME	R		—			—		—	5
33	Mus	"	ME	R		—			—		—	5
34	Mus	24/5/04	ME	R		—			—		—	5
35	Zy argurus	24/5/04	ME	R		43			♀	Ad	MLUP	3
36	Felis catus	24/5/04	O	Scat with feathers in it.								4
37	Zy argurus	25/5/04	ME	WAM 54679	WAM mu No EP	20g			♀	SAd		4
38	Mus		ME	R								5
39	Zy argurus		ME	R		32g			♀	Ad	NLNP NipReg	1
40	Mus		ME	R								4

no tail

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Check Spelling of McSherry
 Old Airport Station Maps show
 McSherry Creek
 McSherry Yard/Bore.

1a Camp Hill
 1b Camp Creek
 2 Spinifex/Sorghum
 3. Limestone Range
 4. Blacksoil
 5. McSherry Riparian

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)				Location McSHERRY GAP								
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
1	Zy. argurus	27/05/04	ME	WAM 54680	WAM MU EP 54680 v	28.5	95	108	♂		Small Testes	3
2	Euro		D									3
3	Zy. a		ME	WAM 54681 R	MU No EP	25			♀	J	Imperbrate	3
4	Zy a		ME	R		56g			♀	Ad	Preg	3 1
5	Zy a		ME	R		28			♂		Small Testes	3 3
6	Zy a		LE	R		24			♀		Imperbrate	3 3
7	R. funneyi		LE	WAM 54676	WAM MU No EP v	15.5	87	70	♂	J		5
8	Felis catu		D TS									Main Road
9	Sus scrofa	27/05/04	O	Scat.								1b
10	Rattus funneyi	28/05/04	ME	WAM 154682	WAM MU No EP v	52	121	116	♂	SAd		5

No tail

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
6	
7	
8	Tracks sighted on road at McSherry Cap Aur observed black/dark tabby on road ~ 0930
9	Dung observed in creek bed
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)				Location MCHERRY GAP								
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	Trapline
11	Euro x2	28/5/04	D									3
12	Zy. argurus	"	ME	R		23			♂	SAd		3
13	Zy. n	"	LE	R		26			♀	Ad	NLNP	3
14	Zy. n	"	ME	Recapture Pregnant female - see record #4								3
15	Zy. n.	"	ME	R		25g			♀	escape		3
16	Zy. n	"	ME	R		17g			♀	SAd		3
17	Zy. n.	"	ME	R		35g			♂	Fig Tester		3
18	Rattus rattus	"	ME	escaped								4
19	P. nanus	29/5/04	ME	M54683	WAM 54683 mu No EP Hair	18	80	83	♂	SAd		2
20	Zy argurus	29/5/04	ME	R	EP-hair in vial	14g			escaped			3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlit (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)				Location. McSHERRY GAP								
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
21	Zy. argurus	20/5/04	LE	R		25			♀	Ad	NLNP	3
22	Zya.	"	ME	R		32			♂	Ad		3
23	Zy. a	"	ME	R		23			♀	Ad		3
24	Zy. a	"	ME	R		29			♂	Ad	Ing Testes	3
25	Zy. a	"	ME	escaped		escaped			♂	Ad		3
26	Zy. a	"	ME	R		29			♀	SAd		3
27	Zy. a	"	ME	escaped		escaped			-	-		3
28	Zy. a	"	ME	R		44			♀	Preg		3
29	Zy. a	"	ME	R		30			♀			3
30	Zy. a	20/5/04	ME	R		31			♀	NLNP		3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

1. Camp.
2. Grasses
3. Limestone Hill
4. Black soil
5. McSherry

Team (AB/TH) (NM/JR) (TS/AT)				Location <i>McSherry Gap</i>								
Record Number	Foot notes Species	Date	*1 Source	*2 Record type	*3 Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Reproduction	*4 Trapline
31	<i>Dog</i>	29/5/04	O									5
32	<i>P. nanus</i>	30/5/04	ME	R		30g			♂	Ad		2
33	<i>Zy. ar.</i>		LE	R		30g			♀	Ad	Preg	3
34	<i>Zy. ar.</i>		ME	R		39g			♀	Ad	Preg	3
35	<i>R. tunneyi</i>		ME	R		21g			♀	Juv		4
36	<i>R. tunneyi</i>		LE	R		-			♀	Ad	Preg	4
37	<i>R. tunneyi</i>		LE	m54685	WAM# TIS Hair MU No EP (✓)	78	137	108	♀	Ad		4
38	<i>P. delicatulus</i>		ME	m54684	WAM# TIS Hair MU No EP (✓)	16g	72	82	♀	Ad	Preg	4
39												
40												

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

- 2 Bellch Flats
2. Gardiner Gorge
3. Minista
4. Quarteite Hill
5. Tectifica Grassland

Team (AB/TH) (NM/JR) (TS/AT)				Location SILENT GROVE								
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
21	R. tunneyi	1 June 04	ME	R		75			♂	Ad	1	1
22	Return		ME	escape								1
23	R. tunneyi		ME	MS4640	MU WAM N:LEP (V)	68	133	123	♂	Ad		1
24	R. tunneyi		LE	R		90			♂	Ad		1
25	R. tunneyi		ME	R		90			♂	Ad		1
26	R. tunneyi		ME	R		85			♂	Ad		1
27	R. tunneyi		ME	escape								1
28	P. nanus		ME	MS4691 (V)	MU WAM N:LEP	22	93	90	♂	Ad		1
29	D. hallucatus		ME	R	MS4686 MU N:LEP	475			♀		No Pouch level	2
30	Zy. argurus		ME	MS4687 (V)	MS4687 MU EP	28	90	111	♀	Ad		2

Trapline 4
- pulled on 3/06
LE + Cages set
at Ranger Quarters
for Bandicoot

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)					Location SILENT GROVE							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
11	Zy. argurus	1/06/04	ME	R		23			♀	SAd		2
12	Zy. argurus		ME	R		32			escape			2
13	D. hallucator		ME	R	M 54688 ME EP	275			♀	SAd	hit Pouch Deer	2
14	Zy. argurus		ME	escape						Ad		2
15	Zy. a		ME	R		52			♀	Ad	WNA	2
16	R. funeyi		ME	R		80			♂	Ad	Large Testes	2
17	Sus.		D.	Black male in creekline								2
18	R. funeyi	1/10/04	ME	R		90			♂	Ad		3
19	R. funeyi		ME	R		22			♂	Juv		3
20	R. funeyi		ME	R		80			♂	SAd		3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location <u>SILENT GROVE</u>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	HB Total Length	Tail length	Sex	Age	Reproduction	Trapline
31	<i>Onychomys leucogaster</i>	2/10/04	S									
32	<i>Rattus</i> v 2	2/06/04	ME	M54692	WAM Hair MU NILEP (D)	105	147	124	♀	Ad	NLNP	2
33	<i>Rattus</i> v ?	2/06/04		M54693	WAM Hair MU NILEP (D)	110	161	141	♀	Ad	NLNP	3
34	<i>Rattus</i> v ?	2/06/04	ME	M54694	WAM Hair MU NILEP (D)	135	164	156	♂	Ad	Large Testes	1
35	<i>R. tunneyi</i>		ME	R		22g			♀	Juv		1
36	<i>R. tunneyi</i>		ME	R		39			♀	Juv		1
37	<i>R. tunneyi</i>		ME	R		85			♂	Ig Testes		1
38	<i>R. tunneyi</i>		LE	R		44			♂			1
39	<i>R. tunneyi</i>		ME	R		22			♀	S.Ad		1
40	<i>R. tunneyi</i>		ME	R		60			♂		Ig Test	1

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)
M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR
Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	Spotlighted 31/05/04 on Silent Grove Entrance Rd.
32	Looked different in trap. Rattus tunneyi or villosissimus??
33	Looked different in trap R. tunneyi Rattus tunneyi or villosissimus??
34	Longer guard hairs ~ 20mm. Possibly Rattus villosissimus??
5	
6	
7	
8	
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location SILENT CIRCLE							
Record Number	Foot notes	Date	*1	*2	*3	Weight	Total Length	Tail length	Sex	Age	Reproduction	*4
21	R. tunneyi	1/06/04	ME	R		80			♀	Ad	Preg	3
22	R. tunneyi		ME	R		23			♂	SAd		3
23	D. hallucator		C	R	LAM	855			♂	Ad		3
24	R. tunneyi		ME	R		42			♀	SAd	Not Pert	3
25	R. tunneyi		ME	R		120			♂	Ad	Large Testes	3
26	R. tunneyi		ME	R		68			♀	Ad	Reg Nipples	3
27	P. nanus		ME	R		32			♂		Testes Big	5
28	R. tunneyi	1/06/04	ME	R		61			♀	Ad		5
29	Dingo	31/05/04	D.	O								
30	Cat	31/05/04	S									

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)
 M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR
 Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	Very large Quoll.
4	
5	
6	
7	
8	
9	Observed from Bell Gorge Gap park 21/02/05.
0	Spotlighted on Silent Grove Entrance Rd 31/05/04

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location SILENT GROVE							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	HB Total Length	Tail length	Sex	Age	Rproduction	Trapline
41	R. tunneyi	2/06/04	ME	R		90			♀	Ad	Preg	1
42	R. tunneyi	2/06/04	ME	R		45			♀	JAd		1
43	Z. argurus	"	ME	R		48			♀	Ad	N-LNP	2
44	Z. argurus	"	LE	R		45			♀	Ad	N-LP	2
45	Z. argurus	"	ME	escape								2
46	Zy. argurus	"	ME	escape					♀	Ad		2
47	R. tunneyi	"	ME	R		65			♂		Testis	3
48	R. tunneyi	"	LE	R		80			♂	Ad	Large Testes	3
49	R. tunneyi	"	ME	R		100			♀	Ad	Lactating	3
50	R. tunneyi	2/06/04	ME	R		70			♂		Testes Is	3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location SILENT GROVE										
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
51	R. tunneyi	2/06/04	ME	R		105			♀		Nip Regress	3
2	R. tunneyi	"	ME	R		24			♂		Imm	3
3	R. tunneyi	"	ME	escaped								3
4	R. tunneyi	"	LE	R		95			♀		Lactating	3
5	Rattus tunneyi	"	ME	R		49					escaped	3
6	Rattus tunneyi	"	LE	R		70			♀		WLNP	4
7	Zy. argurus	"	ME	R		24			♂			4
8	Rattus tunneyi	"	ME	R		75			♂		Ig Testes	4
9	D. hallucator	"	ME	R		310			♀		No Pouch Develop	4
60	R. tunneyi	2/06/04	ME	R		95			♂			5

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
6	
7	
8	
9	widjingari wid jin gar ri
0	wallawa

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location SILENT GROVE							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	HB Total Length	Tail length	Sex	Age	Rproduction	Trapline
61	R. tunneyi	2/6/04	LE	R		43			♀	Imm		5
62	R. tunneyi		ME	R		60			♀	Imm		5
63	R. tunneyi		ME	R		75			♂		Taste Ig	5
64	R. tunneyi		LE	R		70			♂		Taste Ig	5
65	R. tunneyi		LE	R		85			♂		Taste Ig	5
66	R. tunneyi	2/6/04	ME	R		44			♀	Imm		5
7	M. antilopeus	1/6/04	O									
8	D. harrisi???	3/6/04	Cage	m54695 ①	wam mu * Ecto Hair	850			♂	Ad		3
9	M. agilis	2/6/04	S									Bell Gorge Campsite
70	P. bradyotis	3/6/04	D									3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	
6	
7	Observed by AT. Lane & S. dy. 1400 hrs. adjacent to Sike & Grove "Barrow-whul"
8	Large male. Yellow markings on base of liver samples for DNA. throat Large, healthy male, fat on body & tail. (london chest) orange fur
9	Several individuals seen in vicinity of Bell Creek Conspires w/ spotlighting mostly on very burnt patches.
70	TS sighted in Gardners Gap. One hopping up gorge wall.

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location SILENT GROVE							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
71	Rattus tunneyi	3/06/04	ME			70				escape		1
72	" "	"	ME			70			♀	Ad	NLNP	1
73	" "	"	LE			50			♂		Ting	1
74	" "	"	ME			70			♂		Ting	1
75	" "	"	ME			85			♀	Pres NL		1
76	" "	"	ME			53			♂	SAd		1
77	" "	"	LE	escaped								1
78	" "	"	ME			85			♂		Ting	1
79	" "	"	ME			53		escape				1
80	" "	"	LE			85			♂		Ting	1

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)
M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR
Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location SILENT GROVE							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
81	Rattus tunneyi	1	LE			95			♂		Tiny	1
82	" "	1	ME			65			♂		Tiny	1
83	" "	1	ME			20			♂	Juv		1
84	Dasyurus hallucatus	3/6/04	ME			290			♀		UPD	2
85	Zyromys argurus		ME			53			♀		Nipples Enlarged	2
86	Zyromys argurus		ME			53			♀		Preg	2
87	Rattus tunneyi	3/6/04	ME			100			♂	Ad	Tiny	3
88	" "		ME			115			♀	Ad	Preg	3
89	" "		ME			70			♀	Ad	NLWP	3
90	" "	3/6/04	ME		couple							3

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlit (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location <i>Silent Grove</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
91	<i>Rattus funebris</i>	3/10/04	LE			70			♀		NLWP	3
92	" "		ME			45			♂	sAd		?
93	" "		ME			100			♀		Preg	3
94	" "		LE			escape			♂		T. enlarged	3
95	" "		ME			90			♀		NLWP	3
96	" "		ME			75			♀		bigler not enlarged	3
97	" "		ME			66g			♂		T. Ins	3
98	" "		ME			95g			♀		NLWP	3
99	" "		ME			20			♂	juv		3
100	<i>Zyzomys argurus</i>	3/10/04	LE			55			♀		NLWP	4

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location SILENT GROVE							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
101	Rattus funebris	3/06/04	ME			55			♀		Perf.	4
102	Zyromys eremicus !!	3/06/04	SE			58			♀		NLNP	5
103	Rattus funebris		ME			90			♂		Tiny	5
104	" "		ME			65			♀		Unperf.	5
105	" "		LE			75			♂		Tiny	5
106	" "		SE			25			escape	Juv		5
107	" "		LE			85			♂		Tiny	5
108	" "		ME			35			♀	Juv		5
109	" "		ME			47			♀		Perf NLNP	5
110	" "		ME			65			♂			

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location SILENT CROVE							
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
111	Rattus funnigi	4/6/04	x5		Species only identified. Not weighed or sexed				14 Adult + 1 Juvenile			1
112	Zyomys oratus	4/6/04	X5									2
113	Rattus funnigi	4/6/04	x8									3
114	Rattus v. ??	4/6/04	x1		Only ID species							3
115	Rattus funnigi	4/6/04	x5		No other Data Taken.							5
116	Rattus funnigi	4/6/04	x2									6
117	D. hallucatus	4/6/04	x1		WAM mid Tissues.							6,
8												
9												
0												

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line: Use name or code number for trap line. If not on trapline, describe habitat in Notes

Line 4 lifted on slab.

Line 6 = 10 LE
12 days

Kimberley Mammals - NOTES to go with Records

11	Littig Traps - Line 2 Rattus funebris x 15 (14 Adult + 1 Juv)
2	Zygom argurus x 5
3	
4	
5	
6	
7	
8	
9	
0	

Kimberley Mammals -- RECORDS

- 1 Sandstone West (T)
- 2 Sandstone North (B)
- 3 Sandstone South (T)
- 4 River Cliffs
- 5 Soles Island

Team (AB/TH) (NM/JR) (TS/AT) (CP +)		Location SOLEA FALLS										
Record Number	Foot notes		*1	*2	*3							*4
	Species	Date	Source	Record type	Other samples & code numbers	Weight	HB Total Length	Tail length	Sex	Age	Reproduction	Trapline
1	Zy. argurus	7/06/04	ME	M54220	WAM 54220 MU 54220 EP Hair (V)	59	HB 121	127	♀	Ad	Preg.	2
2	SWINK	7/06/04	ME	R151819								5
3	Petrogale brachyotis	6/06/04	D									River Cliffs
4	Petrogale brachyotis	7/06/04	D									Soles Falls -
5	Macropus robustus	7/06/04	Scat									1,2,3,4,5
6	Pseudantechinus ningbingi	7/06/04	D									1
7	Bos	6/06/04	Scat									2
8	Equus	7/06/04		Heard at Sunrise								
9	Petrogale brachyotis		Scat									1,2,3,4,5
10	Felis		O									2 4

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES


*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes



base of

Kimberley Mammals - NOTES to go with Records

1	
2	
3	
4	
5	Scats everywhere.
6	
7	
8	
9	
10	Prints in sand.

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location SOLEA FALLS							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
21	Canis	7/6/04	O	Tracker								1, 3, 4, 5
12	Tachydossus aculeatus	7/6/04	O	Tracker	Scat on Line 5							4
13	Zy.a	8/6/04	ME	Tissue R	WAM 54697 MUM NL EP Hair	29			♂	SAd		2
14	Zy.a	8/6/04	ME	Tissue R	WAM 54697 MUM NL EP	22			♀	Perf.	NLWP	2
15	Hydromys chrysogaster	8/6/04	S									Solea Falls
16	Zy.a	9/6/04	ME	R		31g			♂	Ad		4
17	Zy.a	9/6/04	LE	R	escaped							3
18	Zy.a	9/6/04	ME	R		39			♀	Ad	Preg.	1
19	Zy.a	10/6/04	ME	R		—			♂	Ad	Testes Drooped	6 Solea Falls
20	Zy.a	10/6/04	ME	R		—			♀	Ad	NLWP	2

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as code number

Trap line: Use name or code number for trap line. If not on trapline, describe habitat in Notes

21

Dingo

munt camp 10/6/04

Kimberley Mammals - NOTES to go with Records

1	Prints in sand - common
2	Tracks in sand at base of river cliffs
3	
4	
5	Spotlighted at Slea Falls ~ 200hrs. "Smallish" animal at waters edge. Retreated into ledge in flat rock. Feet lay next to feeding station with cage trap.
6	
7	
8	
9	
0	

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT)				Location <u>INCIDENTAL / ANACDOTAL SIGHTINGS</u>								
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Head body	Tail length	Sex	Age	Rproduction	Trapline
1	Brush Tail Possum											Geikie Gorge
2	Brush Tail Possum		between WAMP + Gibb River Rd									
3	Short eared Rock-wallaby											Windjana Gorge NP
4	Nail Tail Wallaby	27 May	DIS									Fairfield Leopold Rd
5	Tachyglossus											Geikie Gorge NP
6	D. hallucator											Silent Grove
7	Bandicoot											Silent Grove
8	D. hallucator											Tony's Shed
9	Suscotia											Silent Grove Camp site
10	Petaurus brevipes											Silent Grove

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE, ME): Sighted in daylight (D); Spotlighted (S); Found remains (F); Other (O - describe in notes)

WAM-# or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use WAM # or team leader's initials + record number on the data sheet as code number

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	Mike Greenwood had seen BTP a "few years ago" (Ranger LGNP)
2	Mitch Hunter (Ranger WGNP) observed on road "few years ago"
3	Mitch Hunter (Ranger WGNP) observed in Gorge & on north side of reef.
4	at. base of Bahamin tree on Fairfield Leopold Rd. 1230 hrs. TS. disturbed while collecting Mistletoe.
5	Mitch Hunter observed at LGNP.
6	Quolls living in Ranger Quarters at Silent Grove.
7	Ranger Quarters at Silent Grove.
8	Red O'Donnell work shed at Silent Grove - rearranging furniture.
9	Shot by Sergeant Ray Dave in 2003 at Silent Grove. (From his lounge on the verandah.
10	Silent Grove. Jack Dale saw regularly around silent Grove, verandah. ^{marked} Ranger Dave said that a tourist had observed in Mango tree in campground

Uranganba

Kimberley Mammals -- RECORDS

Team (AB/TH) (NM/JR) (TS/AT) (CP +)					Location <i>INCIDENTAL / ANACDOTAL SIGHTINGS</i>							
	Foot notes		*1	*2	*3							*4
Record Number	Species	Date	Source	Record type	Other samples & code numbers	Weight	Total Length	Tail length	Sex	Age	Reproduction	Trapline
11	Bush Tail Possum											Silent Grove
12	Rock Wallaby											Silent Grove
13	Scaly Throated Possum											Silent Grove
14	Rock Ringtail Possum											Silent Grove
15	Edibana											Silent Grove
16	Euro											Silent Grove
17	<i>Onychogalea unguifera</i>	4 June 04	S									Theda Stn
18	<i>Macropus agilis</i>	5 June 04	D									Theda Stn
19	<i>Macropus antilopinus</i>	5 June 04	D									Theda Stn
20												

*1 SOURCE

*2 RECORD TYPE

*3 OTHER SAMPLES

*4 TRAP LINE

Trap type (C, LE ME): Sighted in daylight (D); Spotlit (S); Found remains (F); Other (O - describe in notes)

M-Number or Discarded (D); Sight record (S); Released (R); Photographed (P) or combine e.g. PR

Genetic samples (G); Ectop-parasites (EP); Other (O - describe in notes). Use team leader's initials + record number on the data sheet as **code number**

Trap line. Use name or code number for trap line. If not on trapline, describe habitat in Notes

Kimberley Mammals - NOTES to go with Records

1	Jack Dale ^{very} positive - the shown picture that this was around Silent Grove, Found on creek lines. "Wumbart" "Wum-bart" Very positive
2	Jack Dale - common on hills around "whole country"
3	Jack Dale - "proband" country but not very positive ~ some doubt
4	Jack Dale - not positive ~ very doubtful
5	Common on whole country Widgiemari good eating - lots of fat. Wid gee nah dee
6	"Wulumba"
17	Kalumburn Rd ~ 5km S of Thecla Str. In car headlights ~ 1730
18	Kalumburn Rd South of Thecla Station. 4 individuals 0-18km from Homestead.
9	On Thecla Str - Airstrip ~ 1000hrs
0	