

Field trial to compare baiting efficacy of *Eradicat*® and *Curiosity*® baits

D. Algar, N. Hamilton, M. Onus and S. Hilmer (Science Division, Woodvale)
S. Comer, C. Tiller, L. Bell and J. Pinder (South Coast Region, Albany)
E. Adams and S. Butler (South Coast Region, Esperance)



Plate 1. Feral cat walking across a monitoring plot at Cape Arid National Park with a predated Southern Brown Bandicoot

INTRODUCTION

Baiting is recognised as the most effective method for controlling feral cats (Short *et al.* 1997; EA. 1999; Algar *et al.* 2002; Algar and Burrows 2004; Algar *et al.* 2007; Algar and Brazell 2008) when there is no risk posed to non-target species. The preferred feral cat bait medium (Algar *et al.* 2007) is similar to a chipolata sausage in appearance — it is approximately 20 g wet-weight, dried to 15 g, blanched and then frozen. The bait is composed of 70 % kangaroo meat mince, 20 % chicken fat and 10 % digest and flavour enhancers that are highly attractive to feral cats (Patent No. AU 781829) (see detailed description in Algar and Burrows 2004).

There are two poison bait products intended for the management of feral cat populations in Australia. When the above bait medium (pH 5-6) is dosed with sodium monofluoroacetate (compound 1080), the bait product is known as *Eradicat*®. When the above bait medium is buffered with sodium bicarbonate to pH neutral-alkaline and dosed with para-aminopropiophenone (PAPP) it is known as *Curiosity*®. *Eradicat* and *Curiosity* are registered trademarks of the Western Australian and Commonwealth governments respectively.

A collaborative project between the Department of Sustainability, Environment, Water, Population and Communities (DSEWPaC), the Department of Sustainability and Environment (DSE) and the Department of Environment and Conservation (DEC) has been developing the *Curiosity*® bait product. The project involves bringing together the buffered feral cat bait medium and an encapsulated pellet known as the 'Hard Shell Delivery Vehicle' (HSDV), which contains the toxicant PAPP. The use of the acid-soluble HSDV, in a buffered bait, ensures that the toxin does not disperse throughout the bait but releases in the cat's stomach where it quickly overwhelms the cat's physiological processes (Johnston *et al.* in press). This method of delivering the toxicant also plays a key role in reducing the potential exposure of non-target species. When feeding, feral cats simply shear food items into manageable portions and swallow those portions whole. Thus, they will reliably swallow a pellet that is implanted into a bait. Conversely, most wildlife species process food items more thoroughly in the mouth. This means most animals other than cats tend to reject the HSDV as they eat whereas it is reliably consumed by feral cats (Marks *et al.* 2006; Hetherington *et al.* 2007; Forster 2009;

Johnston unpub. data). Direct injection of PAPP toxin into the bait (i.e. without the pellet delivery device) is not appropriate because it would significantly increase the amount of toxin required and hence significantly increase the risk of non-target poisoning. The pellet delivery device contains about 78 mg of PAPP toxin in pellet form (Johnston *et al.* in press).

A number of cafeteria pen trials have been conducted to test for differences in acceptability of the two bait mediums. These pen trials were conducted at the Perth Cat Haven that provided an opportunity to work with essentially semi-feral cats rather than domestic cats in catteries. Cats in the Haven were housed in individual cages. The cats were offered a choice of the two non-toxic bait mediums. The baits were randomly placed, approximately 20 cm apart. Bait preference was assessed by the medium first selected and consumed by an individual. The baits were offered at the normal time of feeding. All available cats were offered the bait mediums and those which showed interest, initially sniffed each bait type and then selected their choice thus; the experimental design offered a bait choice. Baits were only offered once to any individual cat to avoid any learned behaviour that may have confounded the trial and also to simulate toxic bait delivery in the field. Those cats that did not consume a bait were generally shy and remained in their sleeping boxes. Stress of recent capture and their new surroundings most likely accounted for their behaviour. A number of individuals consumed more than one bait type and the order of preference was also recorded.

A total of 43 cats consumed at least one bait. Analysis of cats' preferences for the two bait mediums, indicated a significant difference in their choice for bait mediums ($\text{Chi}^2 = 31.8$, 1df, $P < 0.001$) with 40 of the cats consuming *Eradicat*® first. The *Eradicat*® bait was the most preferred while the *Curiosity*® bait was the least preferred. However, in 40% of the occasions when the *Eradicat*® bait was consumed first, cats then also chose to eat the *Curiosity*® bait. To test whether this difference is real or an artefact common to cafeteria trials, a trial is required under normal field conditions where bait consumption is assessed in the absence of choice.

The objective of this trial was to compare the efficacy of *Eradicat*® and *Curiosity*® baits in the field to see whether there was any significant difference in baiting efficacy between the two bait types during an operational baiting campaign.

METHODOLOGY

Study area

The trial was conducted in Cape Arid National Park (CANP) and in the adjoining Nuytsland Nature Reserve (NNR). This broad area is located on the south coast of Western Australia (see Figure 1) at 33° 47' 21"S, 123° 24' 47"E (CANP centroid) and 33° 45' 0"S, 123° 41' 24"E (NNR centroid). The area of conservation estate and baiting cells are described below and provided in Table 1.




Figure 1. Location of Cape Arid National Park and Nuytsland Nature Reserve

CANP Park experiences a typical Mediterranean climate, with pronounced winter rainfall and frequently long dry summers with years of high summer rainfall associated with trough movement and thunderstorm activity. Average annual rainfall for CANP (Bureau of Meteorology Station 009879) is 596 mm.

Vegetation of the study area is largely *Eucalyptus incrassata* and *E. tetragona* mallee-heath with large patches of proteaceous shrublands. Water courses are dominated by *E.*

occidentalis woodlands, and smaller incursions of *E. redunca* mallee scrub occur in the vicinity of Thomas River.

Table 1. Location and size of baited cells

| National Park/Nature Reserve | Area (km ²) | <i>Eradicat</i> ® baiting cell (km ²) | <i>Curiosity</i> ® baiting cell (km ²) | % National Park/Nature Reserve baited |
|------------------------------|-------------------------|---|--|---------------------------------------|
| CANP | 2,781 | 973 | 259 | 44 |
| NNR | 6,079 | 227 | - | 4 |
| Total | 8,860 | 1,200 | 259 | 16 |

This location was selected for the following reasons:-

- This location has not been baited for feral cats in the past and surveys have indicted an abundant feral cat population (Comer and Tiller unpub. data);
- There are no non-target species at risk from the proposed baiting program. A complete mammal, reptile and bird species list present in CANP/NNR is provided in Appendix 1;
- The program will assist in the research and recovery efforts for the Critically Endangered Western Ground Parrot (*Pezoporus [wallicus] flaviventris*). The South Coast Threatened Birds Recovery Team identified feral cat predation as likely to be the primary key threatening process for the survival of the species (Comer *et al.* 2010);
- Financial support for the overarching program has been provided by DEC's Nature Conservation Service Special Projects funds, DSEWPaC, South Coast NRM and the State NRM funds directed through the department. DEC has also provided considerable 'in-kind' support through Regional, District, and Science Division resources.

CANP/NNR was divided into two study areas. The larger area (1,200 km²) was baited with *Eradicat*® baits as financial resources were limited and these baits are currently less expensive and easier to manufacture in large volumes. The smaller area (259 km²) was baited with *Curiosity*® baits. The two sites were selected to provide an area of sufficient size to allow enough cats to be trapped (see below) and their activity to be monitored pre- and post-baiting. The trapping locations and monitoring transects are sufficiently distant from the other baiting application, such that mortality of animals can only be ascribed to the one bait type. The study areas are shown in Figure 2.




Figure 2. CANP and adjacent NNR baiting cells, *Curiosity*[®] versus *Eradicator*[®] bait trial

Cat trapping and radio-collaring

Feral cats were trapped several weeks prior to the baiting program, at locations around the track network, in both study areas (see Figure 3 for trap locations). The trapping technique involved the use of padded leg-hold traps Victor ‘Soft Catch’[®] traps No. 3 (Woodstream Corp., Lititz, Pa.; U.S.A.) with a mixture of cat faeces/urine and a olfactory lure (Cat-astrophic, Outfoxed, Victoria) as the attractant. Trap sets were parallel to the track along the verge at 0.5 km intervals. Open-ended trap sets were employed with two traps positioned lengthwise (adjoining springs touching) and vegetation/sticks used as a barrier along the trap sides. The dates of commissioning and decommissioning traps are provided in Table 2.




Figure 3. Location of feral cat trap sites in CANP during 2011

Table 2. Dates of commissioning and decommissioning traps

| Trap No. | Commissioned | Decommissioned | No. trap nights |
|------------------|---------------------|-----------------------|------------------------|
| TT 1-40 | 15/2 | 24/2 | 360 |
| TT 41-42 | 16/2 | 24/2 | 16 |
| TT 43-48 | 20/2 | 24/2 | 24 |
| P 49-53, P 79-86 | 20/2 | 24/2, *P 83 | 51 |
| P 54-66, P 72-78 | 21/2 | 24/2 | 60 |
| P 67-71 | 22/2 | 24/2 | 10 |
| TR 1-19 | 15/2 | 21/2 | 114 |
| GAB 1-12 | 15/2 | 21/2 | 72 |
| GAB 13-31 | 16/2 | 21/2 | 95 |
| GAB 32-34 | 20/2 | 21/2 | 3 |
| PC 1-17 | 22/2 | 24/2, *PC 16 | 33 |
| TOTAL | | | 838 |

*both traps retrieved 21/2 because of non-target activity

Trapped cats were sedated with an intramuscular injection 4 mg/kg Zoletil 100[®] (Virbac, Milperra; Australia). All animals captured were sexed and weighed and coat colour recorded; a broad estimation of age (as either kitten, juvenile or adult) was registered using weight as a proxy for age. A VHF radio-telemetry collar with mortality signal (Sirtrack Ltd, New Zealand) was fitted to trapped cats. Cats were released at the site of capture.

Baiting program

To optimise baiting efficacy, it is essential that the baiting campaign was conducted prior to the onset of late autumn/winter rainfall, which long-term weather records suggested began in April/May (Bureau of Meteorology). A dedicated baiting aircraft deployed the baits at previously designated bait drop points. The baiting aircraft flew at a nominal speed of 130 kt and 500 ft (Above Ground Level) and a GPS point is recorded on the flight plan each time bait leaves the aircraft. The bombardier releases a bag of 50 baits into each 1 km map grid, along flight transects 1 km apart (see Figures 4a and b), to achieve an application rate of 50 baits km⁻². The ground spread of 50 baits is approximately 250 x 150 m (D. Algar, unpub. data). Under this regime, a total of 60,000 *Eradicat*[®] baits and 12,950 *Curiosity*[®] baits were deployed.




Figure 4a. Bait drop locations in the *Eradicat*[®] cell




Figure 4b. Bait drop locations in the *Curiosity*® cell

Surveys of cat activity

Two independent methods were used to monitor baiting efficacy. Baiting efficacy was firstly determined from the percentage of radio-collared cats found dead following the baiting program. The second method involved surveys of cat activity at sand plots to derive indices of activity. The difference in indices pre- and post-baiting was then used as a measure of baiting efficacy.

A track survey transect was established along the Thomas River track in the *Curiosity* baited site and along the Pasley track in the *Eradicat* baited zone. The Thomas River transect was 10.0 km in length and in the larger *Eradicat* baited zone, Pasley transect was 15 km long. These two transects provided a broad coverage of the entire area and an efficient and representative sampling of the population using the surrounding habitat. As multiple indexing assessments were to be made through time on the same area, then the same locations were used (Engeman *et al.* 2002).

The Thomas River and Pasley transects comprised 20 and 30 permanently marked sand pads respectively located at 0.5 km intervals (see Figure 5). Each sand pad was constructed from a 1 m patch of sand that covered the width of the road/track; either end of the sand pad was blocked by vegetation that forced animals to walk across the pad. Two types of plots, passive and active plots, were employed to monitor animal

presence/absence. Passive plots have no attracting lure and detect animals during the normal course of their movements. These plots often generate sample sizes that are too low to adequately monitor population changes (Fleming *et al.* 2001). The active plots contained a lure to attract animals to the plot and thereby increase the likelihood of detecting animals particularly at low density.




Figure 5. Location of sand plot monitoring transects in CANP

To limit potential short comings of using either plot method, a combination of both plot types was employed. The passive plots were located at the 0.5 km sand pads and the active plots were placed at the 1.0 km sand pads. At the active plots an audio lure (Felid Attracting Phonic, Westcare Industries, Western Australia) was used to attract cats to the sand plots during the two survey periods. The audio lures were removed outside the survey periods.

Each plot was observed for the presence or absence of tracks, as it is not possible to determine the number of intrusions by individual animals onto the plot. Each day, the plots were swept to clear evidence of previous activity. Cat activity at the sand plots was recorded over five nights during two survey periods; these were not consecutive nights because of interruption by rain.

Calculation of indices and analyses

Because individuals typically cannot be identified on the basis of track characteristics, it is customary to ignore the number of detections and simply record whether an animal was detected at the station (Ray and Zielinski 2008). The presence/absence data are more robust to statistical analysis than the total number of detections recorded at a station or multiple-station sample units. Thus in this case, sand plot stations have an index of usage expressed as the mean number of positive plots per night. The 'Plot Activity Index' (PAI) is formed by calculating an overall mean from the daily means (Engeman *et al.* 1998; Engeman 2005). The VARCOMP procedure within the SAS statistical software package produced the variance component estimates.

The efficacy of individual baiting programs for both feral cats and foxes was then assessed by comparing these indices immediately prior to and following individual baiting programs. Data were analysed for significant differences using a 'z'-test (Elzinga *et al.* 2001).

RESULTS

Cat trapping and radio-collaring

Twenty-one cats were trapped comprising 12 male and nine females (Table 3). Eleven of these animals were trapped within the *Curiosity*® baiting cell and ten within the *Eradicat*® baiting cell. The location of cat captures is provided in Figure 6. Bodyweight (mean \pm s.e.) for males was 4.5 ± 0.2 kg and 3.0 ± 0.1 kg for females. Nineteen radio-collars were available; ten cats were collared in the *Curiosity*® baiting cell, the eleventh cat died in a trap, cause unknown. Nine cats were collared in the *Eradicat*® baiting cell, the tenth cat was released without a collar following processing. All cats appeared to be in excellent body condition and searches for ectoparasites proved negative.




Figure 6. Location of cat captures

Table 3. Capture records of trapped cats

| Date | Sample No | Trap No | Sex (M/F) | Weight (kg) | Coat colour | Age | Radio-collar frequency (Mg Htz) |
|-----------|-----------|---------|-----------|-------------|-------------|-----------|---------------------------------|
| 16/2/2011 | CANP 1 | TT 13 | M | 4.6 | Tabby | Adult | 151.5470 |
| 16/2/2011 | CANP 2 | TT 16 | F | 3.2 | Tabby | Adult | 151.3480 |
| 16/2/2011 | CANP 3 | TR 18 | M | 3.9 | Tabby | Adult | 151.0482 |
| 16/2/2011 | CANP 4 | G 03 | F | 2.6 | Tabby | Sub-adult | 151.5292 |
| 17/2/2011 | CANP 5 | TT 28 | F | 3.3 | Tabby | Adult | 151.0097 |
| 17/2/2011 | CANP 6 | TR 01 | M | 4.8 | Tabby | Adult | 151.4887 |
| 17/2/2011 | CANP 7 | G 17 | M | 5.2 | Tabby | Adult | 151.4666 |
| 18/2/2011 | CANP 8 | TR 01 | M | 3.6 | Tabby | Adult | 151.0281 |
| 18/2/2011 | CANP 9 | TR 18 | F | 2.8 | Tabby | Adult | 151.5087 |
| 18/2/2011 | CANP10 | G 27 | M | 4.5 | Tabby | Adult | 151.4480 |
| 20/2/2011 | CANP11 | G 29 | M | 5.0 | Tabby | Adult | 151.3893 |
| 20/2/2011 | CANP12 | TR 17 | F | 2.8 | Tabby | Adult | 151.3084 |
| 20/2/2011 | CANP13 | TR 18 | M | 6.0 | Tabby | Adult | Died in trap |
| 21/2/2011 | CANP14 | TT 14 | M | 4.5 | Tabby | Adult | 151.3272 |
| 21/2/2011 | CANP15 | P 01 | M | 4.2 | Tabby | Adult | 151.0662 |
| 21/2/2011 | CANP16 | G 29 | M | 4.5 | Black | Adult | 151.0880 |
| 23/2/2011 | CANP17 | TT 19 | F | 2.9 | Tabby | Adult | 151.4093 |
| 23/2/2011 | CANP18 | PC 15 | F | 2.9 | Tabby | Adult | 151.4268 |
| 23/2/2011 | CANP19 | PC 10 | F | 3.4 | Tabby | Adult | 151.3673 |
| 24/2/2011 | CANP20 | PC 04 | M | 2.6 | Tabby | Sub-adult | 151.5885 |
| 24/2/2011 | CANP21 | P 12 | F | 2.9 | Tabby | Adult | No collar |

Baits, baiting program and impact

The production of *Eradicat*[®] baits was completed in November 2010 at the DEC Harvey bait factory and consisted of 60,000 toxic baits for CANP. A further 13,000 *Curiosity*[®] baits were produced in Victoria by Scientec Research Pty Ltd for the trial comparison between the effectiveness of each bait type. All toxic baits contained Rhodamine B to facilitate the non-target uptake trials.

Eradicat[®] baits and *Curiosity*[®] baits were sweated on 18 March. The two flights required to deliver all *Curiosity*[®] baits were conducted on 20 March. Prior to baiting, the plane was experiencing problems with the GPS program (i.e. not identifying exclusion zones for the cell). On 21 March, rain and low cloud was experienced at CANP during the first bait flight. This reached the Esperance airstrip and delayed the second flight by several hours, in total, only three flights were completed this day. The final two flights to complete delivery of *Eradicat*[®] baits to CANP were conducted on 22 March. Rainfall (2.0

mm) was recorded on the 20 March and 0.2 mm on 21 March, no rainfall was recorded over the following five days.

The status of radio-collared cats was assessed three weeks after the baiting program and is presented in Table 4. Two of the ten radio-collared cats in the *Curiosity*[®] bait cell and two of the eight found radio-collared cats in the *Eradicat*[®] bait cell are presumed to have died from baiting.

Table 4. The status of radio-collared cats following the aerial baiting program

| Bait cell | Sample No. | Status |
|-------------------------------|--------------------|--|
| <i>Curiosity</i> [®] | CANP 3 | Alive |
| “ | CANP 4 | Dead |
| “ | CANP 6 | Alive |
| “ | CANP 7 | Alive |
| “ | CANP 8 | Alive |
| “ | CANP 9 | Alive |
| “ | CANP10 | Alive |
| “ | CANP11 | Alive |
| “ | CANP12 | Alive |
| “ | CANP16 | *Not found initially, later found dead |
| | | |
| <i>Eradicat</i> [®] | CANP 1 | Alive |
| “ | CANP 2 | Alive |
| “ | CANP 5 | Alive |
| “ | CANP14 | Dead |
| “ | CANP15 | Alive |
| “ | CANP17 | Dead |
| “ | CANP18 | Alive |
| “ | CANP19 | Alive |
| “ | CANP20 | **Not found |
| “ | CANP21 (no collar) | Unknown |

*Mortality date to be checked to confirm death due to baiting

**Radio-collar presumed to have malfunctioned

The activity indices for cats, foxes and varanids before and after baiting are presented in Table 5. The only major impact of the baiting program was on cat activity indices in the *Eradicat*[®] baiting cell where a significant reduction in activity was recorded ($z = 3.391$, $P < 0.001$). Interestingly, cat activity indices increased in the *Curiosity*[®] baiting cell following baiting although this was not significant ($z = -1.152$, $P = 0.125$).

Table 5. Activity indices (mean \pm s.e.)

| Species | Site | PAI (pooled across active and passive plots) | | |
|---------|-------------------------------|--|-------------------|-----------------------|
| | | Pre-bait | Post-bait | Significance |
| Cat | <i>Curiosity</i> [®] | 0.080 \pm 0.027 | 0.140 \pm 0.045 | $z = -1.152, P=0.125$ |
| | <i>Eradicat</i> [®] | 0.180 \pm 0.043 | 0.027 \pm 0.013 | $z = 3.391, P<0.001$ |
| Fox | <i>Curiosity</i> [®] | 0.010 \pm 0.010 | 0.010 \pm 0.010 | $z = 0.000, P=0.500$ |
| | <i>Eradicat</i> [®] | 0.007 \pm 0.007 | 0.000 \pm 0.000 | $z = 0.999, P=0.159$ |
| Varanid | <i>Curiosity</i> [®] | 0.020 \pm 0.020 | 0.010 \pm 0.010 | $z = 0.447, P=0.372$ |
| | <i>Eradicat</i> [®] | 0.053 \pm 0.020 | 0.007 \pm 0.006 | $z = 0.554, P=0.290$ |

DISCUSSION

Results from this trial are inconclusive with regard to differences in field acceptability of *Eradicat*[®] and *Curiosity*[®] baits. Mortality of radio-collared cats in both baiting cells was low at 25% and 20% in *Eradicat*[®] and *Curiosity*[®] zones respectively. Despite this, cat activity indices indicated that baiting had a significant impact in the *Eradicat*[®] cell with a decline in cat activity following baiting. Conversely, although not significant, cat activity increased in the *Curiosity*[®] cell post-baiting. The relocation of the sand plot survey transect in the *Curiosity*[®] bait cell because of fire, see below, was not ideal because of its proximity to the unbaited boundary but limited access prevented placement elsewhere. As a consequence, cats at either end of this transect probably had less opportunity to encounter baits, particularly in the short-term following deployment.

To be able to make valid statistical comparisons of cat activity index scores, it is better to have data from a number of transects across the site, rather than a single continuous circuit. In this way, potential variability in activity across the site is accounted for. When using multiple transects to generate an activity index it is necessary to separate the transects by sufficient distance such that the probability of a single animal being recorded on more than one transect in any single survey period is minimized and therefore the transects are independent sampling units. Unfortunately, a fire within the study area several weeks prior to the commencement of this trial (see Figure 2) with subsequent re-alignment of the baiting cells and the lack of track access across the entire study area precluded the use of multiple transect use. As such, comparison of activity scores pre- and post-baiting within and between baiting cells should be made with caution.

Poor bait uptake by cats in either cell, in comparison with previous trials, was most likely a result of reduced bait attractiveness/palatability. Both feral cat bait types require preparation by thawing and sweating prior to deployment in the field. To have sufficient baits prepared to suit the timing of flights necessitates that many baits are prepared the day prior to delivery. However, poor weather conditions (i.e. wind > 25 kts, rain and low cloud) or issues with the plane or equipment can delay bait delivery which may require storage of prepared baits for a number of days. When this occurs, sweated baits are stored in the Western Shield Bait Truck, which contains a sealed refrigeration unit on the

tray. Over a period of a number of days, the lack of air movement in this refrigeration unit can cause mould to develop on the surface of the baits and/or rancidity to commence. In this trial, the longest period between sweating and bait delivery for some baits was four days (the final flights of *Eradicat*[®] to CANP).

The problem with deploying baits that are developing mould and/or rancidity is exacerbated when baits reach the ground. The dense vegetation and cool and damp conditions produce a micro-climate that increases the rate of bait decay. Decaying baits are not attractive/palatable to cats and bait uptake is significantly reduced. Bait longevity in the field is also compromised and thus quality bait availability over time is condensed.

To improve baiting efficacy in the more temperate regions it is recommended that: -

- An efficient artificial method to sweat baits in the field is developed. Reliance on environmental conditions to sweat baits is likely to result in poor quality baits being distributed;
- A test for bait stability/longevity is undertaken in all future trials to gain some measure of bait availability over time;
- Trials to assess baiting efficiency during late summer are conducted. Despite the prey resource likely to be more abundant during these warmer months, bait integrity and longevity will be improved and therefore potentially bait uptake.

ACKNOWLEDGEMENTS

We would like to thank Esperance District support staff: - Klaus Tiedemann; Rob Jose; Nigel Johnstone, Dave Thornberg, and Steve Owen and conservation employees Ben Kremlins and Tracey Klinger. Finally, we gratefully acknowledge the financial support and encouragement from Julie Quinn (Department of Sustainability, Environment, Water, Population and Communities) and financial support from the Department of Environment and Conservation's Special Nature Conservation Program and South Coast NRM Inc.

REFERENCES

- Algar, D., Angus, G.J., Williams, M.R. and Mellican A.E. (2007). Influence of bait type, weather and prey abundance on bait uptake by feral cats (*Felis catus*) on Peron Peninsula, Western Australia. *Conservation Science Western Australia* **6(1)**, 109-149.
- Algar, D. and Brazell, R.I. (2008). A bait-suspension device for the control of feral cats. *Wildlife Research* **35**, 471-476.
- Algar, D., Burbidge, A.A. and Angus, G.J. (2002). Cat Eradication on the Montebello Islands. In 'Turning the Tide: the eradication of invasive species'. (Eds. C.R. Veitch and M.N. Clout) pp 14-18. Invasive Species Specialist Group of the World Conservation Union (IUCN, Auckland.)
- Algar, D. and Burrows, N.D. (2004). A review of Western Shield: feral cat control research. *Conservation Science Western Australia* **5(2)**, 131-163.
- Buckmaster, A.J. (2009). Risk assessment of the likelihood of consumption of the Curiosity® cat bait and the encapsulated toxicant by all Australian terrestrial vertebrate animals: A desktop analysis. Report to the Victorian Government, Department of Sustainability and Environment, Arthur Rylah Institute for Environmental Research.
- Calver, M.C., King, D.R., Bradley, J.S., Gardner, J.L., and Martin G.R. (1989a). Assessment of the potential target specificity of 1080 predator baiting in Western Australia. *Australian Wildlife Research* **16**, 625-638.
- Comer, S., Burbidge, A.H., Tiller, C., Berryman, A., and Utber, D (2010). Heeding Kylorings warning: south coast species under threat. *Landscape* **26**, 48-53.
- Elzinga, C.L., Salzer, D.W., Willoughby, J.W. and Gibbs, J.P. (2001). *Monitoring Plant and Animal Populations*. Blackwell Science Inc. USA.

- Engeman, R.M., Allen, L. and Zerbe, G.O. (1998). Variance estimate for the Allen activity index. *Wildlife Research* **25**, 643–648.
- Engeman, R.M. (2005). Indexing principles and a widely applicable paradigm for indexing animal populations. *Wildlife Research* **32**, 203–210.
- Engeman, R.M., Pipas, M.J., Gruver, K.S., Bourassa, J. and Allen, L. (2002). Plot placement when using a passive tracking index to simultaneously monitor multiple species of animals. *Wildlife Research* **29**, 85-90.
- EA (Environment Australia) (1999). *Threat Abatement Plan for Predation by Feral Cats*. Environment Australia, Biodiversity Group, Canberra.
- Fleming, P.J.S., Corbett, L., Harden, B. and Thomson, P. (2001). *Managing the Impacts of Dingoes and Other Wild Dogs*. Bureau of Rural Sciences, Australian Government Publishing Service, Canberra.
- Forster, G. (2009). Non-target species uptake of feral cat baits containing Rhodamine B. Unpublished B.Sc. (Hons) thesis. Department of Agricultural Sciences, Latrobe University, Bundoora.
- Hetherington, C.A., Algar, D., Mills, H. and Bencini, R. (2007). Increasing the target-specificity of *Eradicat*[®] for feral cat (*Felis catus*) control by encapsulating a toxicant. *Wildlife Research* **34**, 467–471.
- Johnston, M.; Algar, D.; O'Donoghue, M. and Morris, J. (in press). Field efficacy of the Curiosity feral cat bait on three Australian islands. In: Veitch, C.R.; Clout, M.N. and Towns, D.R. (eds.). *Island invasives: Eradication and management*, IUCN, (International Union for Conservation of Nature), Gland, Switzerland.
- King, D.R. (1990). 1080 and Australian fauna. Agriculture Protection Board Technical Series No. 8.

- Marks, C.A., Johnston, M.J., Fisher, P.M., Pontin, K. and Shaw, M.J. (2006). Differential particle size: promoting target-specific baiting of feral cats. *Journal of Wildlife Management* **70**, 1119-1124.
- McIlroy, J. C. (1984). The sensitivity of Australian animals to 1080 poison. VII. Native and introduced birds. *Australian Wildlife Research* **11**, 373-85.
- McIlroy, J. C., King, D. R., and Oliver, A. J. (1985). The sensitivity of Australian animals to 1080 poison. VIII. Amphibians and reptiles. *Australian Wildlife Research* **12**, 113-18.
- Ray, J.C. and Zeilinski, W.J. (2008). Track stations. In: *Noninvasive Survey Methods for Carnivores* (eds R. A. Long, P. MacKay, W. J. Zielinski and J. C. Ray) pp 75-109. Island Press, Washington.
- Short, J., Turner, B., Risbey, D.A. and Carnamah, R. (1997). Control of feral cats for nature conservation. II. Population reduction by poisoning. *Wildlife Research* **24**, 703–714.

Appendix 1

Definitions

Sensitivity to 1080

Approximate Lethal Dose₅₀ data (LD₅₀) where LD₅₀ is the amount of toxin theoretically required to kill 50% of test animals are standardized to mg pure 1080 kg⁻¹, have been taken from Anon. (2002)^A, Twigg *et al.* (2003)^B and Martin *et al.* (2002)^C. Approximate Lethal Dose (ALD) the dose which causes 10% of deaths are provided, in parenthesis, where known from the above references. ALD₅₀ data are greater than the ALD by a factor of less than or equal to 1.5 in approximately 80% of species. LD₅₀ and ALD data are taken from the most recent source and referenced to the above authors by superscript, rather than from the original work. Where data for different populations differ, they are presented as a range, if unknown, they are left blank. Only data from Western Australian populations have been cited.

Sensitivity to PAPP

Sensitivity of Australian vertebrates included in this analysis was obtained from available literature or from personal comments from past and present researchers investigating this toxicant (eg. IA CRC). Additionally Savarie *et al.* (1983) undertook PAPP studies on several North American species that also exist in Australia. Data for these species is also shown. There is large variation in sensitivity to PAPP both intra and inter genus. As a result no extrapolation of sensitivity levels was made between species that had been tested and those that had not, except to note that similar species had been tested.

References

- Algar, D. (2006). A summary of research undertaken to identify non-target risks in the use of the feral cat bait *Eradicat*[®] and encapsulation of the toxin. Unpublished Report, Department of Environment and Conservation.
- Anon. (2002). 1080 Summary Information. Miscellaneous Publication No. 011/2002, ISSN1326-4168. Government of Western Australia.

- Hetherington, C.A., Algar, D., Mills, H. and Bencini, R. (2007). Increasing the target-specificity of *Eradicat*[®] for feral cat (*Felis catus*) control by encapsulating a toxicant. *Wildlife Research* **34**, 467–471.
- Marks, C.A., Johnston, M.J., Fisher, P.M., Pontin, K. and Shaw, M.J. (2006). Differential particle size: promoting target-specific baiting of feral cats. *Journal of Wildlife Management* **70**, 1119-1124.
- Martin, G.R., Twigg, L.E., Marlow, N.J., Kirkpatrick, W.E., King, D.R. and Gaikhorst, G. (2002). The acceptability of three types of predator baits to captive non-target animals. *Wildlife Research* **29**, 489-502.
- Savarie, P.J., Pan, H.P., Hayes, D.J., Roberts, J.D., Dasch, G.J., Felton, R., and Schafer, Jr. E.W. (1983). Comparative acute oral toxicity of para-aminopropiophenone (PAPP) in mammals and birds. *Bulletin of Environmental Contamination and Toxicology* **30**, 122-126.
- Twigg, L.E., Martin, G.R., Eastman, A.F., King, D.R. and Kirkpatrick, W.E. (2003). Sensitivity of some Australian animals to sodium fluoroacetate (1080): additional species and populations, and some ecological considerations. *Australian Journal of Zoology* **51**, 515-31.

MAMMALS

| Common Name | Scientific name | Size (g) | 1080 Sensitivity (mg/kg) | PAPP Sensitivity (mg/kg) | Potential for bait consumption | Potential for pellet consumption | Reason for risk assessment / Risk mitigation |
|--|--|----------|---|--------------------------|--------------------------------|----------------------------------|---|
| Western Grey Kangaroo | <i>Macropus fuliginosus</i> | 54000 | | Unknown | No | No | Herbivorous |
| Gould's Wattled Bat | <i>Chalinolobus gouldii</i> | 16 | 18.8 ^A (14.1) ^C | Unknown | No | No | Insectivorous |
| Southern Brown Bandicoot subsp fusciventer | <i>Isoodon obesulus fusciventer</i> | 1000 | | 6 CRC | Yes | No | Baits consumed in pen trials <i>WA Non-target bait acceptance study (Algar 2006)</i> . 100% pellet rejection <i>Hetherington et al. (2007)</i> |
| Bush Rat | <i>Rattus fuscipes</i> | 225 | 17-43 ^B (27.6) ^C | 696 CRC | Yes | Limited | Baits consumed in RB trials. 14% animals had low RB exposure on encapsulated RB trials. <i>Marks et al 2006</i> . 1 animal tested Nil pellets consumed. <i>WA ball bearing data-1</i> |
| Grey-bellied Dunnart subsp griseoventer | <i>Sminthopsis griseoventer griseoventer</i> | 25 | 4.2 ^B , (2.82) ^B | Unknown | Yes | No | Too small |
| Honey-possum | <i>Tarsipes rostratus</i> | 10 | | Unknown | No | No | Specialist feeder on pollen and nectar |

BIRDS

| Common Name | Scientific name | Length (cm) | 1080 Sensitivity (mg/kg) | PAPP Sensitivity (mg/kg) | Potential for bait consumption | Potential for pellet consumption | Reason for risk assessment / Risk mitigation |
|------------------|---------------------------|-------------|--------------------------|--------------------------|--------------------------------|----------------------------------|--|
| Inland Thornbill | <i>Acanthiza apicalis</i> | 11.5 | | unknown | No | No | Too small |

| | | | | | | | |
|--------------------------------------|--|------|-------------------|---------|----------|----------|--|
| Yellow-rumped Thornbill | <i>Acanthiza chrysorrhoa</i> | 13 | | unknown | No | No | Too small |
| Rufous Fieldwren Subsp campestris | <i>Calamanthus campestris</i> <i>campestris</i> | 13 | | unknown | No | No | Too small |
| Shy Heathwren | <i>Hylacola cautus</i> | 14 | | unknown | No | No | Too small |
| Redthroat | <i>Pyrrholaemus brunneus</i> | 11.5 | | unknown | No | No | Too small |
| White-browed Scrubwren | <i>Sericornis frontalis</i> | 13 | | unknown | No | No | Too small |
| Weebill | <i>Smicrornis brevirostris</i> | 9 | | unknown | No | No | Too small |
| Brown Goshawk | <i>Accipiter fasciatus</i> | 50 | | unknown | No | No | Live prey only |
| Wedge-tailed Eagle | <i>Aquila audax</i> | 101 | 9.1 ^A | unknown | No | No | Live prey and carrion, unlikely to recognise bait as food (Algar et al. 2007) only |
| Swamp Harrier | <i>Circus approximans</i> | 61 | | unknown | No | No | Live prey only |
| Spotted Harrier | <i>Circus assimilis</i> | 61 | | unknown | No | No | Live prey only |
| Black-shouldered Kite | <i>Elanus axillaris</i> | 38 | | unknown | No | No | Live prey only |
| Little Eagle | <i>Hieraaetus morphnoides</i> | 55 | | unknown | No | No | Live prey only |
| Square-tailed Kite | <i>Lophoictinia isura</i> | 56 | | unknown | No | No | Live prey only |
| White-bellied Sea-eagle | <i>Haliaeetus leucogaster</i> | 90 | | unknown | No | No | Live prey only |
| Osprey | <i>Pandion haliaetus</i> | 65 | | unknown | No | No | Live prey only |
| Australian Owlet-nightjar | <i>Aegotheles cristatus</i> | 24 | | Unknown | No | No | insectivorous |
| Sacred Kingfisher | <i>Todiramphus sanctus</i> | 23 | | Unknown | No | No | Live prey only |
| Chestnut Teal | <i>Anas castanea</i> | 48 | | Unknown | No | No | Water filter feeder |
| Grey Teal | <i>Anas gracilis</i> | 67 | | Unknown | No | No | Water filter feeder |
| Australasian Shoveler | <i>Anas rhynchotis</i> | 53 | | Unknown | No | No | Water filter feeder |
| Pacific Black Duck | <i>Anas superciliosa</i> | 60 | 11.8 ^A | Unknown | No | No | Water filter feeder |
| Musk Duck | <i>Biziura lobata</i> | 72 | | Unknown | No | No | Water filter feeder |
| Cape Barren Goose | <i>Cereopsis novaehollandiae</i> | 90 | | Unknown | No | No | Water filter feeder |
| Australian Wood/Maned Duck | <i>Chenonetta jubata</i> | 30 | 11.8 ^A | Unknown | No | No | Water filter feeder |
| Black Swan | <i>Cygnus atratus</i> | 140 | | Unknown | Unlikely | Unlikely | Water filter feeder. Don't bait near watercourses / lakes |

| | | | | | | |
|---------------------|------------------------------------|-----|---------|----------|----------|--|
| Pink-eared Duck | <i>Malacorhynchus membranaceus</i> | 45 | Unknown | No | No | Unlikely to recognise bait as a food source |
| Australian Shelduck | <i>Tadorna tadornoides</i> | 74 | Unknown | No | No | Water filter feeder / grain |
| Australasian Darter | <i>Anhinga novaehollandiae</i> | 94 | Unknown | Unlikely | Unlikely | Feeds on fish and crustaceans. Unlikely to recognise bait as food. Don't bait near waterways |
| Fork-tailed Swift | <i>Apus pacificus</i> | 19 | Unknown | No | No | Insectivorous |
| Great Egret | <i>Ardea alba</i> | 100 | Unknown | Possible | Possible | May recognise the bait as a food source |
| White-necked Heron | <i>Ardea pacifica</i> | 106 | Unknown | Possible | Possible | May recognise the bait as a food source |
| White-faced Heron | <i>Egretta novaehollandiae</i> | 70 | Unknown | Possible | Possible | May recognise the bait as a food source |
| Nankeen Night Heron | <i>Nycticorax caledonicus</i> | 64 | Unknown | Possible | Possible | May recognise the bait as a food source |
| Dusky Woodswallow | <i>Artamus cyanopterus</i> | 18 | Unknown | No | No | Insect / nectar feeders |
| Pied Butcherbird | <i>Cracticus nigrogularis</i> | 36 | Unknown | Possible | Possible | Carrion eaters |
| Australian Magpie | <i>Cracticus tibicen</i> | 44 | Unknown | Possible | Possible | Carrion eaters |
| Grey Butcherbird | <i>Cracticus torquatus</i> | 30 | Unknown | Possible | Possible | Carrion eaters |
| Grey Currawong | <i>Strepera versicolor</i> | 50 | Unknown | Possible | Possible | Carrion eaters |

| | | | | | | | |
|-------------------------------|------------------------------------|------|--------------------|---------|----------|----------|---|
| Short-billed Black-cockatoo | <i>Calyptorhynchus latirostris</i> | 60 | | Unknown | Unlikely | Unlikely | Feeds on seeds fruit and occasional invertebrates. Unlikely to recognise bait as food source |
| Black-faced Cuckoo-shrike | <i>Coracina novaehollandiae</i> | 36 | | Unknown | Unlikely | unlikely | Unlikely to recognise bait as a food source |
| White-winged Triller | <i>Lalage sueurii</i> | 18.5 | | Unknown | Unlikely | unlikely | Too small. Unlikely to recognise bait as a food source |
| Spotted Nightjar | <i>Eurostopodus argus</i> | 33 | | Unknown | No | No | Aerial insectivore. |
| Emu | <i>Dromaius novaehollandiae</i> | 200 | 102.0 ^A | Unknown | Possible | Possible | Eats some carrion. May see bait as food source (Algar et al. 2007), PAPP tolerance (Johnston unpub. data) |
| Red-capped Plover | <i>Charadrius ruficapillus</i> | 16 | | Unknown | No | No | Too small |
| Hooded Dotterel/Hooded Plover | <i>Thinornis rubricollis</i> | 23 | | Unknown | Possible | Possible | Includes invertebrates and small animals in diet. May recognise bait as food |
| Red-kneed Dotterel | <i>Erythrogonyx cinctus</i> | 19 | | Unknown | No | No | Too small |
| Banded Lapwing | <i>Vanellus tricolor</i> | 28 | | Unknown | Possible | Possible | Includes invertebrates and small animals in diet. May recognise bait as food |
| Rufous Treecreeper | <i>Climacteris rufa</i> | 17.5 | | Unknown | No | No | Insectivore. Too small |
| Crested Pigeon | <i>Ocyphaps lophotes</i> | 34 | 23.5 ^A | Unknown | No | No | Granivore |
| Common Bronzewing | <i>Phaps chalcoptera</i> | 36 | 37.6 ^A | Unknown | No | No | Granivore |

| | | | | | | | |
|---------------------------|------------------------------------|-----|-------------------|----------------|----------|----------|--|
| Brush Bronzewing | <i>Phaps elegans</i> | 31 | | Unknown | No | No | Granivore |
| Little Crow | <i>Corvus bennetti</i> | 48 | 12.8 ^A | Unknown | Yes | Unlikely | Omnivorous carrion eater. May reject pellet if similar to <i>C. coronoides</i> |
| Australian Raven | <i>Corvus coronoides</i> | 52 | | 129 CRC *** | Yes | Unlikely | Pellet rejected 37/40 Pellet consumed 3/40. (<i>Johnston unpub. data</i>) |
| Fan-tailed Cuckoo | <i>Cacomantis flabelliformis</i> | 27 | | Unknown | Possible | Possible | Predominant insectivore but may take small vertebrates. |
| Pallid Cuckoo | <i>Cacomantis pallidus</i> | 33 | | Unknown | Possible | Possible | Predominant insectivore but may take small vertebrates. |
| Horsfield's Bronze-cuckoo | <i>Chalcites basalis</i> | 17 | | Unknown | Possible | Possible | Predominant insectivore but may take small vertebrates. |
| Shining Bronze-cuckoo | <i>Chalcites lucidus</i> | 18 | | Unknown | No | No | Too small |
| Black-eared Cuckoo | <i>Chalcites osculans</i> | 21 | | Unknown | No | No | Too small |
| Wandering Albatross | <i>Diomedea exulans</i> | 135 | | Unknown | No | No | Eats at and from the sea. Don't bait on shoreline |
| Yellow-nosed Albatross | <i>Thalassarche chlororhynchos</i> | 82 | | Unknown | No | No | Eats at and from the sea. Don't bait on shoreline |
| Red-eared Firetail | <i>Stagonopleura oculata</i> | 13 | | Unknown | No | No | Too small |
| Zebra Finch | <i>Taeniopygia guttata</i> | 10 | | Unknown | No | No | Too small |
| Brown Falcon | <i>Falco berigora</i> | 50 | 30.1 ^A | Unknown | Unlikely | Unlikely | Unlikely to recognise bait as a food source |

| | | | | | | | |
|------------------------------------|--|----|-------------------|---------|----------|----------|---|
| Nankeen Kestrel/Australian Kestrel | <i>Falco cenchroides</i> | 35 | | Unknown | Unlikely | Unlikely | Unlikely to recognise bait as a food source |
| Australian Hobby | <i>Falco longipennis</i> | 35 | | Unknown | Unlikely | Unlikely | Unlikely to recognise bait as a food source |
| Sooty Oystercatcher | <i>Haematopus fuliginosus</i> | 52 | | Unknown | No | No | Unlikely to recognise bait as a food source. Feeds on shoreline. Don't bait shoreline |
| Pied Oystercatcher | <i>Haematopus longirostris</i> | 51 | | Unknown | No | No | Unlikely to recognise bait as a food source. Feeds on shoreline. Don't bait shoreline |
| White-backed Swallow | <i>Cheramoeca leucosterna</i> | 15 | | Unknown | No | No | Aerial insectivore. |
| Welcome Swallow | <i>Hirundo neoxena</i> | 15 | | Unknown | No | No | Aerial insectivore. |
| Tree Martin | <i>Petrochelidon nigricans</i> | 14 | | Unknown | No | No | Aerial insectivore. |
| Crested Tern | <i>Thalasseus bergii</i> | 48 | | Unknown | possible | Possible | May recognise bait as a food source |
| Caspian Tern | <i>Hydroprogne caspia</i> | 55 | | Unknown | possible | Possible | May recognise bait as a food source |
| Kelp Gull | <i>Larus dominicanus</i> | 60 | | Unknown | possible | Possible | May recognise bait as a food source |
| Pacific Gull | <i>Larus pacificus</i> | 66 | | Unknown | possible | Possible | May recognise bait as a food source |
| Silver Gull | <i>Chroicocephalus novaehollandiae</i> | 43 | | Unknown | possible | Possible | May recognise bait as a food source |
| Blue-breasted Fairy-wren | <i>Malurus pulcherrimus</i> | 15 | | Unknown | No | No | too small |
| Southern Emu-wren | <i>Stipiturus malachurus</i> | 19 | | Unknown | No | No | too small |
| Rufous Songlark | <i>Cincloramphus mathewsi</i> | 17 | | Unknown | No | No | too small |
| Malleefowl | <i>Leipoa ocellata</i> | 61 | 94.0 ^A | Unknown | Possible | Possible | Ground forager. May see bait as a food source |

| | | | | | | |
|---|--------------------------------------|------|---------|------------------------------------|------------------------------------|-------------------------------------|
| Western Spinebill | <i>Acanthorhynchus superciliosus</i> | 15 | Unknown | No | No | Frugivore, Nectivore |
| Spiny-cheeked Honeyeater | <i>Acanthagenys rufogularis</i> | 26 | Unknown | No | No | Frugivore, Nectivore |
| Red Wattlebird | <i>Anthochaera carunculata</i> | 36 | Unknown | No | No | Frugivore, Nectivore |
| Western Wattlebird | <i>Anthochaera lunulata</i> | 31 | Unknown | No | No | Frugivore, Nectivore |
| White-fronted Chat | <i>Epthianura albifrons</i> | 13 | Unknown | No | No | Frugivore, Nectivore |
| Purple-gaped Honeyeater | <i>Lichenostomus cratitius</i> | 19 | Unknown | No | No | Frugivore, Nectivore |
| White-eared Honeyeater | <i>Lichenostomus leucotis</i> | 22 | Unknown | No | No | Frugivore, Nectivore |
| Yellow-plumed Honeyeater | <i>Lichenostomus ornatus</i> | 16 | Unknown | No | No | Frugivore, Nectivore |
| Singing Honeyeater | <i>Lichenostomus virescens</i> | 22 | Unknown | No | No | Frugivore, Nectivore |
| Brown Honeyeater | <i>Lichmera indistincta</i> | 15 | Unknown | No | No | Frugivore, Nectivore |
| Yellow-throated Miner | <i>Manorina flavigula</i> | 27.5 | Unknown | No | No | Frugivore, Nectivore |
| Brown-headed Honeyeater | <i>Melithreptus brevirostris</i> | 14 | Unknown | No | No | Frugivore, Nectivore |
| White-naped Honeyeater | <i>Melithreptus lunatus</i> | 15 | Unknown | No | No | Frugivore, Nectivore |
| White-cheeked Honeyeater | <i>Phylidonyris niger</i> | 18 | Unknown | No | No | Frugivore, Nectivore |
| New Holland Honeyeater | <i>Phylidonyris novaehollandiae</i> | 18 | Unknown | No | No | Frugivore, Nectivore |
| White-fronted Honeyeater | <i>Purnella albifrons</i> | 18 | Unknown | No | No | Frugivore, Nectivore |
| Tawny-crowned Honeyeater | <i>Gliciphila melanops</i> | 17 | Unknown | No | No | Frugivore, Nectivore |
| Rainbow Bee-eater | <i>Merops ornatus</i> | 28 | Unknown | No | No | Insectivore |
| Magpie-lark | <i>Grallina cyanoleuca</i> | 30 | Unknown | Possible | Possible | May recognise bait as a food source |
| Restless Flycatcher | <i>Myiagra inquieta</i> | 21 | Unknown | No | No | Too small |
| Australian Pipit/Richard's Pipit/Groundlark | <i>Anthus novaeseelandiae</i> | 19 | Unknown | No | No | Too small |
| Varied Sittella | <i>Daphoenositta chrysoptera</i> | 12.5 | Unknown | No | No | Too small |
| Australian Bustard | <i>Ardeotis australis</i> | 150 | Unknown | Yes | Yes | May recognise bait as a food source |
| Grey Shrike-thrush | <i>Colluricincla harmonica</i> | 26 | Unknown | Possible | Possible | May recognise bait as a food source |
| Crested Bellbird | <i>Oreoica gutturalis</i> | 23 | Unknown | Possible, but unlikely due to size | Possible, but unlikely due to size | May recognise bait as a food source |
| Golden Whistler | <i>Pachycephala pectoralis</i> | 18.5 | Unknown | No | No | Predominantly insectivorous |
| Spotted Pardalote | <i>Pardalotus punctatus</i> | 10 | Unknown | No | No | Too small |

| | | | | | | |
|-----------------------------|-------------------------------------|------|---------|----------|----------|---|
| Striated Pardalote | <i>Pardalotus striatus</i> | 11.5 | Unknown | No | No | Too small |
| Australian Pelican | <i>Pelecanus conspicillatus</i> | 190 | Unknown | Possible | Possible | May recognise bait as a food source. Don't bait near water bodies |
| Southern Scrub-robin | <i>Drymodes brunneopygia</i> | 23 | Unknown | No | No | Too small |
| Eastern Yellow Robin | <i>Eopsaltria australis</i> | 16 | Unknown | No | No | Too small |
| Jacky Winter | <i>Microeca fascinans</i> | 14 | Unknown | No | No | Too small |
| Hooded Robin | <i>Melanodryas cucullata</i> | 17 | Unknown | No | No | Too small |
| Red-capped Robin | <i>Petroica goodenovii</i> | 12 | Unknown | No | No | Too small |
| Scarlet Robin/Pacific Robin | <i>Petroica multicolor</i> | 14 | Unknown | No | No | Too small |
| Great Cormorant | <i>Phalacrocorax carbo</i> | 92 | Unknown | Unlikely | Unlikely | Predominant fish eater. Don't bait near waterways |
| Black-faced Cormorant | <i>Phalacrocorax fuscescens</i> | 70 | Unknown | Unlikely | Unlikely | Predominant fish eater. Don't bait near waterways |
| Pied Cormorant | <i>Phalacrocorax varius</i> | 80 | Unknown | Unlikely | Unlikely | Predominant fish eater. Don't bait near waterways |
| Little Pied Cormorant | <i>Microcarbo melanoleucos</i> | 64 | Unknown | Unlikely | Unlikely | Predominant fish eater. Don't bait near waterways |
| Brown Quail | <i>Coturnix ypsilophora</i> | 22 | Unknown | No | No | Too small |
| Tawny Frogmouth | <i>Podargus strigoides</i> | 50 | Unknown | No | No | Unlikely to recognise the bait as a food source |
| Great Crested Grebe | <i>Podiceps cristatus</i> | 61 | Unknown | No | No | Feeds on fish and tadpoles. Don't bait near water |
| Hoary-headed Grebe | <i>Poliiocephalus poliocephalus</i> | 30 | Unknown | No | No | Feeds on fish and tadpoles. Don't bait near water |

| | | | | | | | |
|----------------------------------|--|----|-------------------|---------|----------|----------|--|
| Australasian Grebe | <i>Tachybaptus novaehollandiae</i> | 26 | | Unknown | No | No | Feeds on fish and tadpoles. Don't bait near water |
| White-browed Babbler | <i>Pomatostomus superciliosus</i> | 22 | | Unknown | Possible | Possible | Ground feeder - invertebrates spiders reptiles. May recognise bait as a food |
| Cape Petrel | <i>Daption capense</i> | 40 | | Unknown | No | No | Feeds at sea. Don't bait on shoreline |
| Great-winged Petrel | <i>Pterodroma macroptera</i> | 43 | | Unknown | No | No | Feeds at sea. Don't bait on shoreline |
| Soft-plumaged Petrel | <i>Pterodroma mollis</i> | 36 | | Unknown | No | No | Feeds at sea. Don't bait on shoreline |
| Western or Australian Ringneck | <i>Barnardius zonarius</i> | 37 | 10.8 ^A | Unknown | No | No | Granivorous |
| Purple-crowned Lorikeet | <i>Glossopsitta porphyrocephala</i> | 17 | | Unknown | No | No | Granivorous |
| Budgerigar | <i>Melopsittacus undulatus</i> | 20 | | Unknown | No | No | Granivorous |
| Elegant Parrot | <i>Neophema elegans</i> | 24 | | Unknown | No | No | Granivorous |
| Rock Parrot | <i>Neophema petrophila</i> | 23 | | Unknown | No | No | Granivorous |
| Scarlet-chested Parrot | <i>Neophema splendida</i> | 22 | | Unknown | No | No | Granivorous |
| Ground Parrot Subsp flaviventris | <i>Pezoporus wallicus flaviventris</i> | 32 | | Unknown | No | No | Granivorous |
| Mulga Parrot | <i>Psephotus varius</i> | 31 | | Unknown | No | No | Granivorous |
| Regent Parrot | <i>Polytelis anthopeplus</i> | 41 | 11.8 ^A | Unknown | No | No | Granivorous |
| Chestnut-backed Quail-thrush | <i>Cinclosoma castanotum</i> | 24 | | Unknown | No | No | Insectivore |
| Eurasian Coot | <i>Fulica atra</i> | 38 | | Unknown | Unlikely | Unlikely | Water bird. Don't bait near watercourses |
| Purple Swamphen | <i>Porphyrio porphyrio</i> | 48 | | Unknown | Unlikely | Unlikely | Water bird. Don't bait near watercourses |
| Australian Spotted Crake | <i>Porzana fluminea</i> | 23 | | Unknown | Unlikely | Unlikely | Water bird. Don't bait near watercourses |
| Spotless Crake | <i>Porzana tabuensis</i> | 21 | | Unknown | Unlikely | Unlikely | Water bird. Don't bait near watercourses |
| Banded Stilt | <i>Cladorhynchus leucocephalus</i> | 45 | | Unknown | Unlikely | Unlikely | Predominantly water feeders. Don't bait near water bodies. |

| | | | | | | |
|-----------------------------|--------------------------------------|----|---------|----------|----------|--|
| Black-winged Stilt | <i>Himantopus himantopus</i> | 38 | Unknown | Unlikely | Unlikely | Predominantly water feeders. Don't bait near water bodies. |
| Red-necked Avocet | <i>Recurvirostra novaehollandiae</i> | 48 | Unknown | Unlikely | Unlikely | Predominantly water feeders. Don't bait near water bodies. |
| Grey or New Zealand Fantail | <i>Rhipidura fuliginosa</i> | 17 | Unknown | No | No | Aerial insectivore |
| Willie Wagtail | <i>Rhipidura leucophrys</i> | 22 | Unknown | No | No | Aerial insectivore |
| Red-necked Stint | <i>Calidris ruficollis</i> | 16 | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |
| Sanderling | <i>Calidris alba</i> | 21 | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |
| Sharp-tailed Sandpiper | <i>Calidris acuminata</i> | 21 | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |
| Bar-tailed Godwit | <i>Limosa lapponica</i> | 46 | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |
| Eastern Curlew | <i>Numenius madagascariensis</i> | 65 | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |

| | | | | | | | |
|-----------------------------|------------------------------|------|-------------------|---------|----------|----------|--|
| Whimbrel | <i>Numenius phaeopus</i> | 43 | | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |
| Grey-tailed Tattler | <i>Tringa brevipes</i> | 27 | | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |
| Common Greenshank | <i>Tringa nebularia</i> | 34 | | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |
| Common Sandpiper | <i>Actitis hypoleucos</i> | 22 | | Unknown | Unlikely | Unlikely | Predominantly beach / mudflat feeders. Don't bait near water bodies or beaches |
| Little Penguin | <i>Eudyptula minor</i> | 45 | | Unknown | No | No | Feeds at sea. Don't bait on shoreline |
| Southern Boobook / Morepork | <i>Ninox novaeseelandiae</i> | 36 | | Unknown | Unlikely | Unlikely | May not recognise bait as a food source |
| Silvereye | <i>Zosterops lateralis</i> | 12.5 | | Unknown | No | No | insectivore. Frugivore |
| Painted Button-quail | <i>Turnix varius</i> | 19 | | Unknown | No | No | Too small |
| Little Button-quail | <i>Turnix velox</i> | 14 | | Unknown | No | No | Too small |
| Barn Owl | <i>Tyto alba</i> | 40 | 21.8 ^A | Unknown | Unlikely | Unlikely | Unlikely to recognise bait as a food source |

REPTILES

| NAMES VARIOUS | Scientific name | Length (cm) | 1080 Sensitivity (mg/kg) | PAPP Sensitivity (mg/kg) | Potential for bait consumption | Potential for pellet consumption | Reason for risk / Risk mitigation |
|---------------------------------------|---------------------------------------|--------------------|---------------------------------|---------------------------------|---------------------------------------|---|--|
| Ornate Dragon | <i>Ctenophorus ornatus</i> | 8 | | Unknown | No | No | Too small |
| Claypan Dragon | <i>Ctenophorus salinarum</i> | 7 | | Unknown | No | No | Too small |
| Bardick | <i>Echiopsis curta</i> | 60 total | | Unknown | Unlikely | Unlikely | May not recognise bait as food |
| Dugite Subsp affinis | <i>Pseudonaja affinis affinis</i> | 150 total | | Unknown | Unlikely | Unlikely | May not recognise bait as food |
| Western Crowned Snake | <i>Drysdalia coronata</i> | 40 total | | Unknown | Unlikely | Unlikely | May not recognise bait as food |
| Marbled Gecko | <i>Christinus marmoratus</i> | 7 | | Unknown | No | No | Too small |
| Bynoe's Gecko | <i>Heteronotia binoei</i> | 5 | | Unknown | No | No | Too small |
| Three-lined Knob-tail | <i>Nephrurus levis</i> | 8 | | Unknown | No | No | Too small |
| Thick-tailed Gecko | <i>Underwoodisaurus millii</i> | 8 | | Unknown | No | No | Too small |
| Marble-faced Delma | <i>Delma australis</i> | 8 | | Unknown | No | No | Too small |
| Western Three-lined Skink | <i>Acritoscincus trilineata</i> | 6 | | Unknown | No | No | Too small |
| Cryptoblepharus pulcher Subsp clarus | <i>Cryptoblepharus pulcher clarus</i> | Unknown | | | | | |
| Common South-west Ctenotus | <i>Ctenotus labillardieri</i> | 6 | | Unknown | No | No | Too small |
| Barred Wedgesnout Ctenotus | <i>Ctenotus schomburgkii</i> | 4.5 | | Unknown | No | No | Too small |
| King's Skink | <i>Egernia kingii</i> | 20 | | Unknown | No | No | Too small |
| Southern Sand-skink | <i>Egernia multiscutata</i> | 8 | | Unknown | No | No | Too small |
| South-western Crevice-skink | <i>Egernia napoleonis</i> | 12 | | Unknown | No | No | Too small |
| Lowlands Earless Skink Subsp peronii | <i>Hemiernis peronii peronii</i> | 5.5 | | Unknown | No | No | Too small |
| Southern Slider | <i>Lerista dorsalis</i> | 6.5 | | Unknown | No | No | Too small |
| South-western Slider Subsp intermedia | <i>Lerista microtis intermedia</i> | 5 | | Unknown | No | No | Too small |
| Shrubland Morethia Skink | <i>Morethia obscura</i> | 4.5 | | Unknown | No | No | Too small |

| | | | | | | | |
|-------------------------------------|---------------------------------|-----------|--------|---------|-----|----------|---|
| Bobtail Skink Subsp rugosa | <i>Tiliqua rugosa rugosa</i> | 25 | 800.0A | Unknown | Yes | Possible | Carion eater. Likely to view baits as food source. Lay baits during cooler months when not active |
| Southern Blind Snake | <i>Ramphotyphlops australis</i> | 50 total | | Unknown | No | No | Termite / Ant eater. Unlikely to identify bait as food |
| Heath Monitor / Rosenberg's Monitor | <i>Varanus rosenbergi</i> | 100 total | 235.0A | 3 CRC | Yes | Possible | Carnivore. Ground feeding. May perceive bait as food. Bait in cooler months when less active. Highly tolerant to 1080 |