


**BUSHLAND PLANT SURVEY RECORDING SHEET 1- use pencil only**

BUSHLAND AREA DAVEY'S FARM - KONNONGARRING SITE NUMBER Kon 8  
 DATE TRIP 31/8/96 RECORDERS M.P., B.M.  
 DATE TRIP \_\_\_\_\_ RECORDERS \_\_\_\_\_  
 DATE TRIP \_\_\_\_\_ RECORDERS \_\_\_\_\_  
 BOTANIST Cate Tausz

**1. LOCATION of the QUADRAT**

From 'Bushland Plant Survey' written by B. Keighery (1994) and published by the Wildflower Society of WA (Inc.), PO Box 64 Nedlands WA 6008.

Mud Map Draw a sketch of the location of the site below.




**Road Location**

Geographic Location Latitude 31 00.483 S Longitude 116 47.161 E Altitude 357 ± 43  
 Reference Map 31 00 - 548 116 47.031 ± 65 M

Photograph Photographer's Name \_\_\_\_\_ Photo No \_\_\_\_\_

Topographic position Circle position of site on the transect (alter the transect if necessary)



**2. SITE DATA** Circle the correct response.

Slope flat gentle steep Aspect N NE E SE S SW W NW

Surface Soil Sandy loam Colour L Brown  
 Exposed rock type \_\_\_\_\_ % surface \_\_\_\_\_

Sub-surface Soil Loam to clay Colour Brown  
 Rock type \_\_\_\_\_ depth to rock \_\_\_\_\_

Drainage well mod poor depth water \_\_\_\_\_ cm Wet all year winter/spring

Litter 30-70 % cover \_\_\_\_\_ Bare Ground 10-30 % cover \_\_\_\_\_  
 Depth cm 2-3 cm



*Allocasuarina huegeliana* (Rock Sheoak) low open woodland (over *Santalum spicatum* open shrubland (c. 2%) over *Dampiera laevidulacea*, *Opercularia jagrata* low open shrubland over *gossypoid + open herbland + v. open sedgeland*.

## BUSHLAND PLANT SURVEY RECORDING SHEET 2 - use pencil only

### 3. VEGETATION STRUCTURE AND COVER

From 'Bushland Plant Survey' written by B. Keighery (1994) and published by the Wildflower Society of WA (Inc.), PO Box 64 Nedlands WA 6008.

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class 2-10% 10-30% 30-70% over 70%

| TREES            |                                            |                                             | MALLEES                                        |         |          |
|------------------|--------------------------------------------|---------------------------------------------|------------------------------------------------|---------|----------|
|                  | over 30m                                   | 10-30m                                      | under 10m                                      | over 8m | under 8m |
| LIFE FORM        |                                            |                                             |                                                |         |          |
| COVER CLASS (%)  |                                            |                                             | less 2-10%                                     |         |          |
| DOMINANT SPECIES |                                            |                                             | <i>Allocas. hueg</i>                           |         |          |
| SHRUBS           |                                            |                                             | SHRUBS                                         |         |          |
|                  | over 2m                                    | 2m-1m                                       | under 1m                                       |         |          |
| LIFE FORM        |                                            |                                             |                                                |         |          |
| COVER CLASS (%)  |                                            | under 2%                                    |                                                |         |          |
| DOMINANT SPECIES |                                            | <i>Santalum spicatum</i>                    | <i>Dampiera opercularis</i>                    |         |          |
| GRASSES          |                                            | HERBS                                       | SEDGES                                         | OTHER   |          |
| LIFE FORM        |                                            |                                             |                                                |         |          |
| COVER CLASS (%)  | 30-70%                                     | 10-30%                                      | 2-10%                                          |         |          |
| DOMINANT SPECIES | <i>Stipa elegantissima</i><br><i>Vulpa</i> | <i>Asteraceae</i> SP<br><i>Clamaescilla</i> | <i>Lepidosperma</i> SP.<br><i>Schoenus</i> SP. |         |          |

### 4. VEGETATION CONDITION

|   |           | COMMENTS |
|---|-----------|----------|
| 1 | PRISTINE  |          |
| 2 | EXCELLENT |          |
| 3 | VERY GOOD | ✓        |
| 4 | GOOD      |          |
| 5 | DEGRADED  |          |



e if required

97 totals

- Column 4 identification check

From 'Bushland Plant Survey' written by B. Keighery (1994) and published by the Wildflower Society of WA (Inc.), PO Box 64 Nedlands WA 6008.

ADJ. 46 each 2000 on granules - AIC 48 Allocas Compektion -  
AIC 48 Allocas Compektion -

Allo-cao Cambodiana

Protein (w/ post)  
Protein (w/ pre)