


BUSHLAND PLANT SURVEY RECORDING SHEET 1- use pencil only

BUSHLAND AREA NABALLING ^{Wardy + Peter} Berrill Brown SITE NUMBER MB 4
 DATE TRIP 4.10.97 RECORDERS HELEN, KELD, ANN, JEFF + PETER
 DATE TRIP _____ RECORDERS _____
 DATE TRIP _____ RECORDERS _____
 BOTANIST _____

1. LOCATION of the QUADRAT

From 'Bushland Plant Survey' written by B. Keighery (1994) and published by the Wildflower Society of WA (Inc.), PO Box 64 Nedlands WA 6008.

Mud Map Draw a sketch of the location of the site below.




Road Location Wb DABBY RD Lipseig Gully.

Geographic Location Latitude 33°28'13.9 S Longitude 117°30'32 E Altitude _____
 Reference Map _____

Photograph Photographer's Name _____ Photo No _____

Topographic position Circle position of site on the transect (alter the transect if necessary)



2. SITE DATA Circle the correct response.

Slope flat gentle steep Aspect N NE E SE S SW W NW

Surface Soil Sandy loam (collapsed surface from water logging) Colour grey brown
 Exposed rock type _____ % surface _____

Sub-surface Soil duplex sandy loam with clay Colour _____
 Rock type _____ depth to rock _____

Drainage well mod poor depth water _____ cm Wet all year winter/spring

Litter 2-10% % cover _____ Depth <1 cm _____
 Bare Ground 2-30 % cover _____

BUSHLAND PLANT SURVEY RECORDING SHEET 2 - use pencil only

From 'Bushland Plant Survey' written by B. Keighery (1994) and published by the Wildflower Society of WA (Inc.), PO Box 64 Nedlands WA 6008

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class 2-10% 10-30% 30-70% over 70%

	TREES			MALLEES	
	over 30m	10-30m	under 10m	over 8m	under 8m
LIFE FORM					
COVER CLASS (%)					
DOMINANT SPECIES					

	SHRUBS		SHRUBS
	over 2m	2m-1m	under 1m
LIFE FORM			
COVER CLASS (%)			30-70% (40%)
DOMINANT SPECIES			Halosarcia sp.

	GRASSES	HERBS	SEDGES	OTHER
LIFE FORM				
COVER CLASS (%)	2-10%	10-30		
DOMINANT SPECIES	salt grass puccinellia	Cotula coronopifolia Spergularia diandra		

4. VEGETATION CONDITION

1	'PRISTINE'	COMMENTS Salt affected, dead acacia acuminata shrubs ~ 30 yrs since salt arrival. samphire spreading. herb layer of cotula (possibly a weed)
2	EXCELLENT	
3	VERY GOOD	
4	GOOD	
5	DEGRADED	

BUSHLAND PLANT SURVEY RECORDING SHEET 3 - use pencil only

5. SPECIES PRESENCE

Record on Sheet

Date 4/6/97

- Column 1 plant name.

- Column 2 plant number

• Column 3 flowering time- TICK if species flowering

- Column 4 identification check

From 'Bushland Plant Survey' written by B. Keighery (1994) and published by the Wildflower Society of WA (Inc.), PO Box 64 Nedlands WA 6008.

[illegible]

Apr 34: Schoenus subflavus