

SEED  
PRICE  
LIST



1997-1998

Department of Conservation and Land Management



## THE SEED CENTRE

The Department of Conservation and Land Management (CALM) Seed Centre supplies seed of native and exotic trees and shrubs. It is located within CALM's Plant Propagation Centre on Burnside Road, West Manjimup.

**Mailing Address:** Seed Centre  
Department of Conservation and Land Management  
Brain St  
MANJIMUP  
Western Australia 6258

**Telephone:** (08) 9772 1288

**Fax:** (08) 9772 1305

**International Fax:** +61 8 9772 1305

**E-mail:** markd@calm.gov.wa.au

**Business Hours:** Monday to Friday, 8.00am to 4.30pm

**Ordering Seed** (seed order form in centre pages):

- Your order should specify clearly the quantity of seed required (in grams) of each species, the date of ordering, your contact address and phone number.
- Supply of some species may be limited by current stocks. Please indicate alternative species.
- The minimum weight of seeds per specie is FIVE grams (5 gm). In the case of Banksia and Xylomelum species, the minimum quantity per specie is 10 seeds.
- Postage must be included with all orders. (See last page for postal charges.)
- The minimum total order, including postage, is \$10.00.
- Orders of \$100.00 or less must include payment before dispatch.
- Payment may be by cheque, credit card or money order made payable to 'Dept. of CALM'. Personal callers may pay in cash.

If you have any queries about ordering, please contact Monica Dalton or Mark Dalton at the Seed Centre.

## ADDITIONAL SERVICES

1. **Scarification.** Seed of some species belonging to the following genera requires scarification (i.e. mechanical abrasion of the seed coat) before sowing:

Acacia	Hardenbergia
Bossiaea	Hovea
Ceratonia	Kennedia
Chamaecytisus	Senna
Chorizema	Templetonia
Daviesia	

This service is available through the Seed Centre at no extra cost. Please indicate if you require this service by ticking the appropriate box on the order form.

2. **Export.** Seed that is intended for export overseas must be inspected by the Agriculture Department prior to dispatch. A 'Phytosanitary Certificate' must accompany the seed to its final destination. If you require a certificate, we can arrange an inspection (price on application).
3. **Import.** If an Import Permit is required, please provide us with the necessary documents.
4. **Pure Seed.** Seed listed in the price list as 'pure seed' is 99% pure.
5. **Provenance Data.** Provenance data may be issued on request.
6. **Individual Treelot Collections.** Individual treelot collections are available (see over).
7. **Special Collections from Natural Stands (SC)** The parent trees of those species listed as SC in the price list, have been carefully selected from natural populations for their good form and vigour. This seed has derived from ten (10) or more widely spaced, unrelated parent trees, therefore a broad genetic base is guaranteed. Seed from this type of collection would be suitable for the establishment of seed production areas, provenance trials and nursery sowing.

## PLEASE NOTE

The 'average height and width' figures given in the seed list indicate the potential size of the mature species in its natural environment under normal growing conditions. Should the site conditions (soil type, moisture, space, exposure to wind, or salinity) vary markedly, plant performance will differ. Unless otherwise stated, all *Eucalyptus* species are sold with seed and chaff.

### Individual Treelot Collections

Prices on application. Seed from this type of collection would be suitable for provenance and progeny trials. If you require any particular species not listed here, field collections can be arranged.

Species	Provenance
<i>Corymbia calophylla</i> 'rosea'	WELD 6
" " "	DINGUP
<i>Eucalyptus astringens</i>	HIGHBURY
" <i>camaldulensis</i> var. <i>obtusata</i>	ARROWSMITH LAKE
" " " "	MURCHISON RIVER
" <i>cornuta</i>	QUABICUP HILL
" <i>gomphocephala</i>	CERVANTES
" "	LAKE PRESTON
" <i>kochii</i> ssp. <i>kochii</i>	WILGIE HILL
" <i>kochii</i> ssp. <i>plenissima</i>	MT. GIBSON
" <i>longicornis</i>	LAKE JOHNSTON
" "	MT. GIBSON
" <i>occidentalis</i>	CALYERUP CREEK
" <i>platypus</i> var. <i>platypus</i>	CHILLINUP
" <i>salmonophloia</i>	MT. GIBSON
" "	LAKE JOHNSTON
" <i>salubris</i> var. <i>salubris</i>	MT. GIBSON
" " " "	LAKE CRONIN
" <i>sargentii</i> ssp. <i>sargentii</i>	KOORDA
" " " "	LAKE CARMODY
" " " "	LAKE HURLESTONE
" <i>suggrandis</i> ssp. <i>suggrandis</i>	EMU ROCK

Photographs of parent trees for the above species are available on request.

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>ACACIA</b>								
<i>acradenia</i>	.19	106	silky wattle	2-5	1.5-3	250	G,L	A
<i>acuminata</i>	.17	54	jam tree	6-10	3-5	380	L	AEG
<i>alata</i> var. <i>alata</i> *	.47	58	winged wattle	1-2	1-2	400	SLG	A
<i>ampleiceps</i>	.15	46	salt wattle	2-8	2-5	250	SC	JKW
<i>anaticeps</i>	.17	2	duck-leaved wattle	3-6	3-4	250	S	K
<i>ancistrocarpa</i>	.13	27	fitzroy wattle	3-5	2-4	250	SLG	AW
<i>anuera</i>	.11	74	mulga	4-10	2-6	130	L	EFHW
<i>baileyana</i>	.09	39	cootamundra wattle	5-8	5-8	550	SL	A
<i>bivenosa</i>	.19	36	two-veined wattle	3-4	4-5	400	SL	BJW
<i>browniana</i> var. <i>obscura</i>	.25	208		1-2	1-2	600	S	A
<i>celastrifolia</i>	.13	60	grey myrtle wattle	2-4	1.5-2	600	S,C	ASU
<i>citrinoviridus</i>	.21	23		7-8	3-6	200	SG	ACW
<i>cochlearis</i>	.43	128	coastal wattle	1-2	1-2	500	S	J
<i>coriacea</i>	.11	12	wirewood	3-10	2-4	200	S	ABE
<i>cowleana</i>	.11	88	hall's creek wattle	2-5	1.5-2.5	250	S	WE
<i>cyclops</i>	.11	26	western coastal wattle	2-4	2-4	500	S	BHJKVW
<i>decurrens</i>	.09	70	early black wattle	5-15	5-8	630	L	A
<i>dentifera</i>	.17	87	toothed wattle	2-4	3-4	300	SG	A
<i>divergens</i>	.33	400	shark's tooth wattle	1.5-2.5	1-2.5	800	SLG	A
<i>drummondii</i> subsp. <i>candolleana</i> *	.33	345	Drummond's wattle	0.5-2	1-2	700	SL	A
<i>drummondii</i> subsp. <i>drummondii</i> *	.63	270	dwarf golden wattle	0.5-2	1-2	700	SL	A

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>ACACIA (cont.)</b>								
<i>elata</i>	.19	24	cedar wattle	10-20	5-10	600	S,C	ABCEJ
<i>extensa</i>	.17	111	wiry wattle	1-3	2-3	650	SL	A
<i>farnesiana</i>	.09	9	scented wattle	3-7	2-3	200	LC	ANSW
<i>gilbertii</i>	.59	138	white-flowered wattle	1-2	1-2	800	SL	AJW
<i>holosericea</i>	.15	83	candelabra wattle	2-5	1.5-3	250	SL	ATW
<i>horridula</i>	.67	61		0.5-2	1-2.5	800	SL	AJ
<i>inaequilatera</i>	.21	15	camel bush	1-3	1-2	200	S	AWS
<i>insolita</i>	.43	85	solitary wattle	0.5-1.5	1.5-2	700	SG	A
<i>iteaphylla</i>	.15	31	Flinders Range wattle	3-5	3-6	300	SL	AB
<i>ixiophylla</i>	.31	227		2-3	1-2	300	S,LC	AB
<i>jennerae</i>	.13	23	coonavitra wattle	2-5	2-7	200	S,SL	AB
<i>jibberdingensis</i>	.21	30	Jibberding wattle	3-5	3-4	250	S	AB
<i>lasiocalyx</i>	.13	57	wilyurwur	3-5	4-6	300	SL	AW
<i>lateriticola</i>	.19	149	Dwellingup wattle	0.5-2	1-2	800	SG	A
<i>linophylla</i>	.13	12	wanderry mulga	3-7	4-6	150	S	
<i>longifolia</i> var. <i>longifolia</i>	.11	95	golden rods	3-10	2-8		S,C	ABEJKNSU
<i>maitlandii</i>	.21	92	spikey wattle	2-3	1-3	200	S	A
<i>mangium</i>	.45	90		15-25	5-8		LC	C
<i>mearnsii</i>	.13	85	black wattle	8-25	6-10	500	S,C	CEJSU
<i>melanoxylon</i> (TAS. provenance)	.19	58	tasmanian blackwood	5-30	4-15	890	L	BCDE
<i>merrallii</i>	.37	16	Merrall's wattle	0.3-2	1-4	350	S,SL,L,C	BJKU
<i>microbotrya</i>	.11	26	manna gum	4-6	2-4	380	L	ABOW

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
ACACIA (cont.)								
<i>myrtifolia</i>	.15	44	myrtle wattle	1-3	2-3	550	LCG	ABJ
<i>myrtifolia</i> var. <i>angustifolia</i>	.15	132	narrow-leaved myrtle wattle	1-3	2-3	600	LCG	A
<i>nerosa</i>	1.02	76	rib wattle	0.5-1	0.5	600	SLC	A
<i>papyrocarpa</i>	.45	108	western myall	3-10	3-6	200	SL,C	ABCN
<i>pentadenia</i>	.39	196	karri wattle	3-4	3-4	1000	SL	AW
<i>podalyriaefolia</i>	.09	26	Queensland silver wattle	3-5	3-4	510	L	A
<i>prominens</i>	.21	54	golden rain wattle	5-20	4-15	700	S,L,C	ABCEJST
<i>pruinocarpa</i>	.13	37	gidgee, black gidgee	3-10	3-5	250	SG	C
<i>pulchella</i>	.15	344	prickly moses	0.5-1.5	1-2	460	SLG	ABW
<i>pyrifolia</i>	.15	21	kanji bush	3-5	1-3	250	SG	AMW
<i>redolens*</i>	.25	128	vanilla wattle	1-4	3-8	400	S,C	JKPRW
<i>salicina</i>	.41	19	native willow	4-10	3-5	200	CL	BU
<i>saligna</i>	.09	51	western wattle	3-10	3-6	550	SLG	ABJKNQVW
<i>scirpifolia</i>	.19	44		2-5	2-3	350	S	AB
<i>sclerosperma</i>	.13	5	limestone wattle	3-5	4-5	200	SL	ABJW
<i>silvestris</i>	.17	48	red wattle	20-35	10-20	700	S,L,C	BEMU
<i>trachycarpa</i>	.19	16	sweet scent minni-ritchi	3-5	3-5	250	SG	ABTW
<i>translucens</i>	.27	85	poverty bush	0.5-2.5	1-3	250	SG	APW
<i>tumida</i>	.13	27	pindan wattle	3-8	2-4	250	S	AW
<i>urophylla</i>	.25	188	tall-leaved wattle	3-4	2-4	700	SL,G	AW
<i>wanyu</i>	.17	7	wanyu	2-4	2-3	200	SG	AM
<i>xiphophylla</i>	.15	16	snakewood	4-6	3-6	200	S	BW

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>ACTINOSTROBUS</b>								
<i>arenarius</i>	.29	66	sandplain cypress	3-5	2-4	580	S	ABF
<i>pyramidalis</i>	.25	59	swamp cypress	3-8	2-5	400	S	ABHKRW
<b>AGONIS</b>								
<i>flexuosa</i>	.33	600	WA peppermint	8-15	5-15	760	SL	ABCEJRTV
<i>linearifolia</i>	.45	3500	swamp peppermint	2-5	2-3	800	SL,C	BHJ
<i>parviceps</i>	.29		fine ti-tree	1-3	1-3	500	S,SL,L,C	BHJMW
<b>ALLOCASUARINA</b>								
<i>acutivalvis</i>	.21	384	black tammar	3-7	2-4	300	LG	BW
<i>campestris</i> subsp. <i>campestris</i>	.19	909	common tammar	1-4	1-3	300	SL,GC	ABNT
<i>corniculata</i>	.43	1670	tammar	2-5	2-3	250	SG	ABU
<i>decaisneana</i>	.17	60	desert oak	5-18	3-7	200	S	CEN
<i>decussata</i>	.23	180	karri oak	6-10	4-6	900	SL	BF
<i>fraseriana</i>	.23	217	sheoak	10-20	5-10	750	SG	BCEF
<i>huegeliana</i>	.21	555	granite sheoak	10-15	5-10	400	SL	ABCE
<i>humilis</i>	.21	625	dwarf casuarina	0.5-2	1.5-2	620	SLG	A
<i>lehmanniana</i>	.19	770	lehmann's sheoak	1-4	1-3	600	SL	B
<i>pinaster</i>	.29	108	compass bush	2-5	1-3	300	S,SL,L,CL	BTUW

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>ANIGOZANTHUS</b>								
<i>flavidus</i>	.55	1540	tall kangaroo paw	0.5-1	1	700	SL	A
<b>BANKSIA (Price per 100 seeds)</b>								
<i>ashbyi</i>	4.50	89	ashby's banksia	.2-6	2-4	300	S	ABMU
<i>attenuata*</i>	9.10		slender banksia	4-10	3-8	350	S	ABJKM
<i>audax</i>	5.80	33	bold banksia	1-2	1-2.5	250	S,SL	MU
<i>baueri*</i>	12.00	16	wooly banksia	1-2	2-3	350	S	AM
<i>baxteri*</i>	7.00	27	birds-nest banksia	3-4	3-5	400	S	ABMU
<i>burdettii*</i>	9.00	8	burdett's banksia	3-6	3-6	600	S	ABM
<i>candolleana*</i>	12.00	6	propeller-banksia	1-1.5	1-3	500	S	ABMW
<i>chamaephyton</i>	90.00	9	fishbone banksia	0.4	1-2	500	S,SG	AP
<i>coccinea*</i>	7.00	73	scarlet banksia	3-5	2-4	400	S,L	AJMU
<i>elderiana*</i>	7.80	22	swordfish banksia	1-4	1-4	240	S	ABMU
<i>gardneri var. hiemalis</i>	12.00	19	prostrate banksia	0-0.3	1-2.5	500	S	MP
<i>grandis*</i>	6.20	11	bull banksia	4-8	3-5	500	SLG	AEIJMU
<i>grossa</i>	10.00	16	coarse banksia	1-1.5	1.5-2	600	S	AM
<i>hookeriana*</i>	9.00	28	hooker's banksia	2-3	2-4	600	S	AM
<i>ilicifolia</i>	110.00	26	holly-leaved banksia	5-15	3-8	600	S,SL	ABJO
<i>incana</i>	19.00	19	hoary banksia	0.3-0.7	0.5-1.5	600	S	AM
<i>lanata</i>	19.00	24		0.3-1	0.5-2	500	S	AP
<i>leptophylla var. leptophylla</i>	5.00	37		1.5-2.5	1-2	500	S	ABM

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>BANKSIA (cont.)</b>								
<i>littoralis*</i>	7.00	52	swamp banksia	5-15	3-10	600	S,SL	ABW
<i>media*</i>	5.80	21	southern plains banksia	2-6	3-5	300	S,L,C	AMTUV
<i>menziesii*</i>	10.00	12	menzies' banksia	1-10	5-10	400	S	AM
<i>nutans var. nutans*</i>	7.00	26	nodding banksia	1-2	1-3	400	S	AMUW
<i>prionotes*</i>	4.30	36	acorn banksia	5-12	4-6	350	S	AJMU
<i>quercifolia</i>	5.00	58	oak leaved banksia	2-4	1.5-3	800	S	ABH
<i>sceptrum*</i>	6.90	20	sceptre banksia	2-7	2-5	350	S	ABJMT
<i>speciosa*</i>	9.00	11	showy banksia	3-6	3-8	350	S,C	ABJMU
<i>sphaerocarpa var. caesia*</i>	7.00	59	round-fruited banksia	1-2	2-4	400	S,C	ABMTU
<i>sphaerocarpa var. sphaerocarpa*</i>	12.00	21	fox banksia	0.5-2	2-4	500	SG	ABMTU
<i>victoriae*</i>	8.60	22	wooly orange banksia	2-5	2-5	350	S	AMTU
<i>violacea</i>	7.00	43	violet banksia	1-1.5	1-2	300	S	AJMU
<b>BOSSIAEA</b>								
<i>aquifolium</i>	.45	118	water bush	3-4	1-2	800	SG,L	AW
<i>laidlawiana</i>	.45	100	netic	5-6	1-2	800	SG,L	AW
<i>linophylla</i>	.47	355		2-3	2-3	800	SL	AT
<i>ornata</i>	.89	400	broad leaved brown pea	0.5-1	0.5-1.5	800	LG	AW
<b>BRACHYCHITON</b>								
<i>australis</i>	.13	12	broad-leaved bottle tree	3-7	3-5		SL	A
<i>diversifolius</i>	.09	9	northern kurrajong	7-18	3-5	300	SL	CJ
<i>gregorii</i>	.21	21	desert kurrajong	4-8	2-4	250	SL	ACKN

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>CALLISTEMON</b>								
<i>citrinus</i>	.21	8500	crimson bottlebrush	2-8	2-6	550	SL,CG	ABHJRTUV
<i>glauca</i>	.23	7100	Albany bottlebrush	2-4	1-3	610	SL	ABH
<i>phoeniceus</i>	.19	5200	lesser bottlebrush	2-4	3-5	350	S,LG	ABHKRV
<i>rigidus</i>	.25	3100	stiff bottlebrush	1-2.5	2-4	300	LC	AJKU
<i>salignus</i>	.23	2600	willow bottlebrush	5-15	3-5	500	SL,CG	AHJKTUV
<i>speciosus</i>			see <i>Callistemon glauca</i>					
<i>viminalis</i>	.25		weeping bottlebrush	1-12	1.5-6	550	S,CL,C	BJKT
<b>CALLITRIS</b>								
<i>columellaris</i>	.33	92	coastal cypress pine	15-20	5-15	380	SL	AJKRV
<i>drummondii</i>	.37	250	cypress pine	2-4	1-2	500	LC	A
<i>preissii</i>	.27	71	Rottneest island pine	10-20	5-8	760	S	ACEJ
<i>roei</i>	.33	90	Roe's cypress pine	3-5	1-3	350	SLG	AR
<b>CALOTHAMNUS</b>								
<i>blepharospermus</i>	.61	50		1-2	1-3	300	S	A
<i>chrysantherus</i>	.31		golden netbush	0.5-2.5	1-2	500	S,SL,L, CL	AU
<i>gilesii</i> *	.43	730	Giles' net-bush	2-4	2-4	250	SL	ABJUV
<i>homalophyllus</i>	.51	1000	Murchison claw flower	2-4	2-4	300	S,C	ABJU
<i>quadrifidus</i> *	.21	200	one-sided bottlebrush	2-4	2-5	630	SL	ABJMR
<i>rupestris</i>	.21	350	cliff net-bush	1.5-3	2-3	630	SLG	ABV

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>CALOTHAMNUS (cont.)</b>								
<i>sanguineus</i>	.65	300	blood-red net-bush	0.5-2.5	1-3	760	S	AJRVU
<i>validus</i>	.29		Barren's claw flower	2-3	2-3	500	SG	ABJ
<b>CASUARINA</b>								
<i>crinata</i> subsp <i>crinata</i>	.17	295	black oak	8-25	5-10	200	SG,LC	CEHJKV
<i>cunninghamiana</i>	.19	1600	river oak	10-35	10-12	500	SL,G	BCEHNW
<i>obesa</i>	.17	650	swamp oak	6-30	4-12	450	C	CEHKW
<b>CERATONIA</b>								
<i>siliqua</i>	.09	6	carob tree	8	8-10	500	SL	CINX
<b>CHAMAECYTISUS</b>								
<i>proliferus</i>	.05	45	tree lucerne	4-5	5	380	SL	BN
<b>CHORILAENA</b>								
<i>quercifolia</i>	1.70	470	chorilaena	1-4	1-3	1000	L	AT
<b>CHORIZEMA</b>								
<i>ilicifolium</i>	1.89	500	holly flame pea	0.5-3	1-3	800	SG	A
<b>CLEMATIS</b>								
<i>pubescens</i>	.25	400	western clematis climber	5-10	5-10	800	SL	AR

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>CORYMBIA</b>								
<i>calophylla</i> (pure seed)	.19	10±2	marri	13-25	10-13	500	SLG	ACEIJ
<i>citriodora</i> (cv)	.19	110±45	lemon-scented gum	15-30	10-25	650	SLG	CDLR
<i>citriodora</i> (pure seed)	.85	180						
<i>citriodora</i> (QLD provenance)	.45	110±45	lemon-scented gum	15-30	10-25	650	SLG	CDLR
<i>dichromophloia</i>	.37	45	variable-barked bloodwood	8-12	5-8	250	LG	AEC
<i>eximia</i>	.41		yellow bloodwood	3-15	2.5-12	600	S,SL,L	ABE
<i>ficifolia</i>	.41	45±15	red-flowering gum	6-15	5-20	1000	SG	ABCIJ
<i>ficifolia</i> (pure seed)	1.15							
<i>ficifolia</i> (SC)	.85							
<i>haematoxylon</i>	.25	20	mountain marri	6-12	4-8	600	SG	BEI
<i>maculata</i>	.19	107±40	spotted gum	20-35	10-20	750	SLG	CDEFIRV
<i>maculata</i> (pure seed)	.80	155						
<i>ptychocarpa</i>	.67	50	swamp bloodwood	8-15	3-10	500	SGC	AH
<b>CUPRESSUS</b>								
<i>arizonica</i>	.27	160	Arizona cypress	12	6	510	SL	ABCET
<b>DAVIESIA</b>								
<i>cordata</i>	.59	50	bookleaf	0.6-2	1-2	600	S,SL,L,G	AM

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>DRYANDRA</b> (per 100 seeds)								
<i>carlinoides</i> *	12.00	80	pink dryandra	0.6-1.5	1-1.5	500	S,SL,L	AM
<i>cuneata</i>	12.00	85	wedge-leaved dryandra	1.5-3	1-2	350	S,SL,G	M
<i>ferruginea</i> *	17.00	56		0.3-2	0.4-2	350	S,SL,L, CL	AM
<i>formosa</i>	7.80	143	showy dryandra	1.5-8	2-5	500	S,SL,L,CL,C,G	AMT
<i>nobilis</i> subsp. <i>fragrans</i>	11.00	60	golden dryandra	1.5-4	2-4	450	S,SL,L,CL,C,G	AMT
<i>quercifolia</i> *	11.00	84	oak-leaved dryandra	1-4	1-3.5	500	S,SL,CL	AM
<i>sessilis</i> *	7.00		parrot bush	2-6	1.5-3.5	500	S,C	MJU
<i>shuttleworthiana</i>	45.00	22	shuttleworth's dryandra					
<i>speciosa</i> *	11.00	47	shaggy dryandra	0.3-1.5	1-3.5	500	S,SL,L,CL,G	AM
<b>EUCALYPTUS</b>								
<i>accedens</i>	.23	60	powderbark wandoo	8-25	6-15	750	GC	CDEFIQ
<i>agglomerata</i>	.45	90	blue-leaved stringybark	20-30	10-15	700	L	DG
<i>albida</i>	.71	250	white-leaved mallee	2.5-5	4-6	350	SG	ABM
<i>alligatrix</i>	.21	348	Argyle apple	8-18	8-15	800	SL	AMRUV
'anceps'			see <i>Eucalyptus phenax</i>					
<i>angustissima</i>	.71		narrow-leaved mallee	1.5-4	3-7	300	S,SL	BIKL
subsp. <i>angustissima</i>								
<i>angulosa</i>	.17	40	ridge-fruited mallee	2-6	3-7	250	SL	BI
<i>annulata</i>	.19	170	open-fruited mallee	2.5-8	4-7	300	SL	ACIKLVW
<i>arachnaea</i> subsp. <i>arachnaea</i>	.21	150	black-stemmed mallee	3-5	4-6	300	LC	BCW



Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>arachnaea</i> subsp. <i>arachnaea</i> (SC)	1.89	260	black-stemmed mallee	3-5	4-6	300	LC	BCW
<i>argophloia</i>	.29	150	Queensland western white gum	20-35	10-15	700	LC	EGI
<i>argyphaea</i>	.41	130	silver mallet	10-15		300	SL	EQ
<i>aspersa</i>	.47	65	forest mallee	2-3		700	SG	
<i>aspratilis</i>	.19	200		2-3	3-4	250	SL	AB
<i>astringens</i>	.23	130±43	brown mallet	5-25	4-10	510	L	CDEFIKQV
<i>astringens</i> (pure seed)	3.82	1100						
<i>baueriana</i>	.35	240	blue box	12-25	8-15	600	LC	C
<i>botryoides</i>	.17	300±214	southern mahogany	12-40	10-25	630	SL	CEGIJKV
<i>botryoides</i> (pure seed)	4.20	1750						
<i>brachycorys</i>	.27	40	cowcowing mallee	3-5	3-6	200	LC	ACK
<i>brachyphylla</i>	.41			4-5		250	L	A
<i>brevistylis</i>	.63	50	Rate's tingle	20-40	10-20	1000	L	CDEF
<i>brookeriana</i>	.45		Brooker's gum	10-40		850	SG	C
<i>brockwayi</i>	.33	338±160	dundas mahogany	12-24	8-11	280	LG	BCEF
<i>brockwayi</i> (pure seed)	6.85	1200						
<i>buprestium</i>	.17	4	apple mallee	2-6	3-6	300	S	ABT
<i>burdettiana</i>	.55	130	Burdett's mallee	3-9	3-10	380	SLC	ABCJV
<i>burracoppinensis</i>	.21	55	burracoppin mallee	3-8	3-6	250	S	AB
<i>caesia</i> subsp. <i>caesia</i> * (cv)	.29	140	gungurru	4-10	3-8	380	L	AUM
<i>caesia</i> subsp. <i>magna</i> * (cv)	.25	70	silver princess	5-15	4-8	380	LC	AUM
<i>calpicola</i> (cv)	.63	35	Hamelin Bay mallee	2.5-5	3-5	760	S	ABJV

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>calophylla</i>			see <i>Corymbia calophylla</i>					
<i>calycogona</i> var. <i>calycogona</i> *	.27	60-388	gooseberry mallee	2.5-8	3-7	250	SL	ABIKV
<i>camaldulensis</i> var. <i>camaldulensis</i> (Lake Coorong, VIC)	.31	698±465	river red gum	15-50	15-35	250	SL	CEHIKV
<i>camaldulensis</i> var. <i>camaldulensis</i> (Lake Hindmarsh, VIC)	.27	698±465	river red gum	15-50	15-35	250	SL	CEHIKV
<i>camaldulensis</i> var. <i>camaldulensis</i> (cv)	.11	698±465	river red gum	15-50	15-35	200	SL	CEHIV
<i>camaldulensis</i> var. <i>obtusa</i> (Arrowsmith, WA)	.21	950	river red gum	12-20	12-30	400	S	CEHIV
<i>camaldulensis</i> var. <i>obtusa</i> (cv)	.11	698±465	river red gum	15-50	15-35	200	SL	CEHIV
<i>camaldulensis</i> var. <i>subcinerea</i> (Silverton, NSW)	.27	698±465	river red gum	15-50	15-35	250	SL	CEHIKV
<i>camaldulensis</i> var. <i>subcinerea</i> (pure seed)	5.45							
<i>campaspe</i> *	.19	180	silver-topped gimlet	5-11	4-7	250	CL	ACEFIKM
<i>campaspe</i> (pure seed)	6.20							
<i>capillosa</i> subsp. <i>capillosa</i>	.21	200	Wheatbelt wandoo	10-30	10-15	250	SL	CEFGI
<i>capillosa</i> subsp. <i>capillosa</i> (pure seed)	4.78							
<i>celastroides</i> subsp. <i>celastroides</i>	.27	650	snap and rattle	3-20	3-10	250	SLC	AEI
<i>ceratocorys</i>	.23	22				250	S	
<i>cinerea</i>	.29	430	Argyle apple	6-18	4-15	600	S,SL,L,CL,C	AHLMO
<i>citriodora</i>			see <i>Corymbia citriodora</i>					
<i>cladocalyx</i> (cv)	.17	90	sugar gum	15-35	15-20	380	SL	CDEFIUW

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>cladocalyx</i> 'Nana' (cv)	.15	50	dwarf sugar gum	8-10	8-15	380	SL	BCITU
<i>cladocalyx</i> 'Nana' (pure seed)	3.20	900						
<i>clelandii</i>	.19	310	Cleland's blackbutt	6-12	5-8	250	L	ABCE
<i>clivicola</i> (SC)	1.52	200	green mallet	8-12	4-6	350	CL,G	A
<i>cloeziana</i>	.15	120±87	gympie messmate	20-35	10-20	550	SL	DFG
<i>concinna</i>	.33	200-400	Victoria Desert mallee	3.5-9	5-9	200	SL	ABI
<i>conferruminata</i>	.13	70	Bald Island marlock	6-9	7-10	500	SL	ABCJV
<i>conferriminata</i> (pure seed)	1.60	590						
<i>coolabah</i> var. <i>rhodoclada</i>			see <i>Eucalyptus vitrix</i>					
<i>cornuta</i>	.15	180	yate	6-30	5-10	500	SL	ACHIJKV
<i>cornuta</i> (pure seed)	3.00							
<i>corrugata</i>	.27	150	rough-fruited mallee	6-15	4-8	250	SLG	ABCI
<i>crebra</i>	.21	520	narrow-leaved ironbark	20-30	10-15	550	SCL	ACDEIV
<i>crucis</i> subsp. <i>crucis</i> * (cv)	.51	50	silver mallee	3-8	4-10	280	SL	AMU
<i>crucis</i> subsp. <i>lanceolata</i>	.41		Southern Cross silver mallee	5-15	4-10	250	SL,L	AMU
<i>cupularis</i>	.41	50	Hall's Creek white gum	6-10	3-5	400	S	AC
<i>cylindrocarpa</i>	.25	170	woodline mallee	3-10	4-6	200	SL,LC	EI
<i>decipiens</i> subsp. <i>adesmophloia</i>	.19	130	redheart	3-12	5-8	350	SL	BITV
<i>delegatensis</i>	.33	104±52	alpine ash	15-40	10-30	700	L,C	BCDU
<i>densa</i> subsp. <i>densa</i>	.51	215		6-8	4-6	400	SL,L,CL ,L	B
<i>desmondensis</i> * (cv)	.25	110	desmond mallee	3-6	3-6	450	SL	A
<i>dichromophloia</i>			see <i>Corymbia dichromophloia</i>					
<i>dielsii</i>	.21	200	cap-fruited mallee	4-7	5-7	400	C	ACIK

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>dielsii</i> (SC)	.91	290	cap-fruited mallee	4-7	5-7	400	C	ACIK
<i>diptera</i>	.19	140	two-winged gimlet	5-8	5-7	300	SCL	ACIJK
<i>diptera</i> (SC)	.91	180	two-winged gimlet	5-8	5-7	300	SCL	ACIJK
<i>diversicolor</i>	.63	80	karri	30-70	8-15	900	SL	DEI
<i>diversicolor</i> (pure seed)	9.50	600						
<i>diversifolia</i> *	.13	35	soap mallee	3-10	3-10	250	S	BCIJ
<i>dolichorhyncha</i> *	.21	45	fuchsia mallee	3-7	3-5	350	SL	ABIUV
<i>dolichorhyncha</i> * (SC)	.35							
<i>doratoxylon</i>	.29	180	spearwood mallee	3-7	3-7	400	SG	ABKR
<i>drummondii</i>	.23	30	Drummond's mallee	4-8	4-8	400	SL	ABI
<i>dundasii</i>	.31	320	dundas blackbutt	7-20	5-10	260	GL	ACFIV
<i>ebbanoensis</i> subsp. <i>ebbanoensis</i>	.17	90	sandplain mallee	3-9	4-8	250	SGL	ABI
<i>eremophila</i> subsp. <i>eremophila</i> *	.17	265	sand mallee	4-8	5-8	220	SL	ACIKV
<i>erythrocorys</i> *	.19	10	illyarrie	4-9	3-8	480	SL	AJ
<i>erythrocorys</i> (pure seed)	2.50	85						
<i>erythrocorys</i> * (SC)	.80							
<i>erythronema</i> var. <i>erythronema</i>	.21	230	red-flowered mallee	4-9	4-7	350	L	AIQT
<i>erythronema</i> var. <i>marginata</i>	.25	150	red-flowered mallee	4-9	4-7	300	L	AIQ
<i>eudesmioides</i> subsp.	.19	50	mallalie	2.5-4	4-6	350	SL	BI
<i>eudesmioides</i>								
<i>eudesmioides</i> subsp. <i>pallida</i>	.41		pallid mallee	2.5-4	3-4	300	S	BIM
<i>ewartiana</i>	.25	65	Ewart's mallee	3-6	3-5	200	SL	A
<i>eximia</i>			see <i>Corymbia eximia</i>					
<i>falcata</i>	.23	260	silver mallee	3-15	3-7	340	SG	ACGIQ

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>ficifolia</i>			see <i>Corymbia ficifolia</i>					
<i>flavida</i>	.21	110	yellow-flowered mallee	6-8		250	L	AI
<i>flocktoniae</i>	.15	100±44	merritt	6-15	3-6	250	SL	ABFIKQV
<i>foecunda</i>	.29	460±210	narrow-leaved red mallee	3-8	4-8	800	S	ABIV
<i>formanii</i>	.45	110	Forman's mallee	4-12	4-8	250	LS	ABM
<i>forrestiana</i> subsp. <i>forrestiana</i> *	.19	45	fuchsia mallee	3-7	3-5	350	SL	ABIUV
<i>forrestiana</i> subsp. <i>forrestiana</i> * (SC)	.25							
<i>forrestiana</i> subsp. <i>dolichorhyncha</i>			see <i>Eucalyptus dolichoryncha</i>					
<i>fraseri</i>	.25	235	balladonia gum	6-20	4-10	250	L	ACI
<i>gardneri</i> subsp. <i>gardneri</i>	.19	140	blue mallet	6-9	4-8	350	GLS	ACIQRU
<i>globulus</i> subsp. <i>globulus</i> (S. TAS provenance)	.29	78±55	Tasmanian blue gum	15-70	10-25	600	L	DEFILRV
<i>globulus</i> subsp. <i>globulus</i> (pure seed)	2.60	350						
<i>globulus</i> subsp. <i>bicostata</i>	.25	109±76	southern blue gum	10-45	8-20	700	L	DIRV
<i>globulus</i> subsp. <i>bicostata</i> (pure seed)	2.60	470						
<i>gomphocephala</i> (cv)	.19	81±37	tuart	10-35	8-15	600	S	CDEFIJV
<i>gomphocephala</i> (pure seed)	2.85	460						
<i>gongylocarpa</i>	.25	60	desert gum	4-20	3-15	150	S	CEF
<i>gracilis</i> *	.21	520	yorrell	3-18	4-10	200	SL	ABEFIKUV
<i>grandis</i>	.21	628±331	rose gum	20-55	10-25	900	L	DE

# SEED ORDER FORM

Department of Conservation & Land Management  
 Seed Centre  
 Brain Street, MANJIMUP 6258  
 WESTERN AUSTRALIA

Phone: (08) 9772 1288  
 Fax: (08) 9772 1305  
 Fax: INT +61 (08) 977 21305

DATE: \_\_\_/\_\_\_/\_\_\_

CUSTOMER / INSTITUTE NAME: \_\_\_\_\_ TELEPHONE: \_\_\_\_\_

POSTAL ADDRESS: \_\_\_\_\_ FAX: \_\_\_\_\_

TOWN / CITY: \_\_\_\_\_ STATE / COUNTRY: \_\_\_\_\_ POSTCODE: \_\_\_\_\_

Office use	Grams reqd.	Species	Cost/gram	Total \$

<b>MEANS OF DELIVERY (please tick)</b> <input type="checkbox"/> Standard Post <input type="checkbox"/> Express Post <input type="checkbox"/> Airmail <input type="checkbox"/> Courier	<b>OTHER SERVICES</b> <input type="checkbox"/> Scarification (to those species that require it) <input type="checkbox"/> Propagation information
---	--

<b>COST OF SEED</b>	
<b>POSTAGE &amp; HANDLING</b>	
<b>TOTAL COST</b>	

Enclosed is my Cheque/Money Order or debit my credit card for \$ \_\_\_\_\_

Bankcard     Mastercard     Visa

Card Number

Card Holders Name: \_\_\_\_\_ Expiry Date: \_\_\_\_\_

Signature: \_\_\_\_\_

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>grandis</i> (pure seed)	4.70	3300						
<i>gratiae</i>	.19	290	Lake Grace gum	6-8	4-5	300	SL	IK
<i>griffithsii</i>	.29	160	Griffith's grey gum	4-12	5-8	250	SL	ABC
<i>grossa</i> *	.17	320	coarse-leaved mallee	2-6	3-6	250	SL	ABV
<i>guilfoylei</i>	.25	40	yellow tingle	15-35	10-20	1100	L	CDEFGI
<i>gummii</i>	.87	392±249	cider gum	10-25	6-15	800	S,L,C	BCEMNUN
<i>haematoxylon</i>			see <i>Corymbia haematoxylon</i>					
<i>halophila</i>	.27	140	salt lake mallee	2.5-4	4-6	500	SC	BHK
<i>horistes</i>	.50	120	oil mallee	5-7	3-10	400	S	BL
<i>horistes</i> (pure seed)	7.65	2200						
<i>hypochlamydea</i> subsp. <i>hypochlamydea</i>	.27	70	rough-barked mallee	4-6	4-5	250	S	BTW
<i>hypochlamydea</i> subsp. 'smooth bark'	.43	450	smooth-barked mallee	3-4		250	S,SL	ITW
<i>incerata</i>	.71	380	Mount Day mallee	3-4	3-4	250	SL	ABM
<i>incrassata</i>	.17	25	ridge-fruited mallee	3-8	4-7	250	SG	ABIV
<i>jacksonii</i>	.27	60	red tingle	40-70	15-25	1150	SL	DEI
<i>jucunda</i>	.21	30	yuna mallee	6-8	4-6	300	S	BI
<i>jutsonii</i>	.21	150	Jutson's mallee	1.5-8	4-8	250	SL	BI
<i>kessellii</i> subsp. <i>kessellii</i>	.23	80	Kessell's mallee	2-6	4-6	300	S	ABIW
<i>kingsmillii</i> *	.27	80±47	Kingsmill's mallee	4-8	3-6	200	SL	AI
<i>kochii</i> subsp. <i>kochii</i>	.27	230	oil mallee	5-7	4-8	350	S	ABEIL
<i>kochii</i> subsp. <i>plenissima</i>	.25	220	oil mallee	5-7	4-8	250	SL	BEIL
<i>kondininensis</i>	.33	310	kondinin blackbutt	8-15	5-10	300	LC	CIKW

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>kruseana</i> *	.35	215	book-leaf mallee	2-6	3-6	250	SL	AMU
<i>kumarlensis</i>	.63	380	Kumarl gum	10-12		250	L,CL,C	CDHK
<i>laeliae</i>	.27	180	Darling Range ghost gum	10-18	8-12	800	SGC	ACI
<i>lanepolei</i>	.25	47±25	salmonbark wandoo	10-15	6-12	600	SG	ACI
<i>lansdowneana</i> subsp. <i>albopurpurea</i>	.41	320	purple-flowered mallee	3-10	4-8	310	SL	ABJV
<i>largiflorens</i>	.25	701±474	black box	10-20	8-15	200	CL	ABEGHIKV
<i>lata</i>	.39	200	round-leaved mallee	2-3	2-3	250	S,SL,L	AIM
<i>latens</i> (cv)	.71		North Bannister mallee	2-3		500	SLG	
<i>lehmannii</i>	.27	70	Lehmann's mallee	2.5-5	4-5	450	S	ABIJV
<i>lehmannii</i> (pure seed)	8.70	520						
<i>leptopoda</i> subsp. <i>arctata</i>	.23	310	tammin mallee	2.5-6	3-5	200	S	ABE
<i>leptopoda</i> subsp. <i>leptopoda</i>	.23	310	tammin mallee	2.5-6	3-5	200	S	ABE
<i>lesouefii</i>	.25	160	Goldfield's blackbutt	6-15	3-8	230	L	ACEIK
<i>leucoxylon</i> subsp. <i>leucoxylon</i>	.21	235±109	yellow gum	10-30	6-20	400	SL	ACEFIJTUV
<i>leucoxylon</i> subsp. <i>megalocarpa</i> (cv)	.19	396±280	large-fruited yellow gum	4-9	5-10	400	SL	ABEIJTU
<i>leucoxylon</i> subsp. <i>megalocarpa</i> (pure seed)	4.55	1300						
<i>leucoxylon</i> subsp. <i>pruinosa</i> (cv)	.23	300	inland blue gum	8-10	5-10	400	SL	ABEIJTU
<i>leucoxylon</i> 'rosea' (cv)	.23	200	pink-flowered yellow gum	3-4	4-5	400	SL	ABIT
<i>livida</i>	.29	115	mallee wandoo	3-4	2-3	250	SL	AI
<i>longicornis</i>	.25	280±158	red morrel	8-30	5-20	250	CL	CEFIKV
<i>loxophleba</i> subsp. <i>gratiae</i>			see <i>Eucalyptus gratiae</i>					

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>loxophleba</i> subsp. <i>lissophloia</i>	.23	220	smooth barked York gum	6-8	4-6	250	SL	IK
<i>loxophleba</i> subsp. <i>loxophleba</i>	.21	630±350	York gum	6-18	5-12	300	SLC	CEFIKLV
<i>loxophleba</i> subsp. <i>loxophleba</i> (pure seed)	4.20	3200						
<i>loxophleba</i> subsp. <i>supralaevis</i>	.25	400	York gum	10-15	5-10	250	L, LC	CEFIKLV
<i>macrandra</i> * (cv)	.19	30	long-flowered marlock	3-8	4-10	400	SLC	ABIJUV
<i>macrocarpa</i> subsp. <i>elachantha</i> *	.21	20	small-leaved mottlecah	2-3	3-10	350	S	ABM
<i>macrocarpa</i> subsp. <i>macrocarpa</i> *	.21	25	mottlecah	3-4	4-12	350	S	ABM
<i>maculata</i>			see <i>Corymbia maculata</i>					
<i>mannensis</i> subsp. <i>vespertina</i>	.65	120	western mallee	4-7	4-5	250	S,SL	BC
<i>marginata</i> subsp. <i>marginata</i> *	.17	16±9	jarrah	6-40	6-25	600	SLG	CDEFGI
<i>marginata</i> subsp. <i>marginata</i> (pure seed)	1.65	65						
<i>megacarpa</i>	.25	40	bullich	2-30	4-20	750	SL	AVCHI
<i>megacornuta</i>	.17	90-207	warted yate	5-12	5-8	400	SLG	ABCIJTW
<i>melanophitra</i>	.41	450	black mallet	8-12	4-6	500	SL	CI
<i>melanoxylon</i>	.37	330	black morrel	7-24	5-15	250	LC	ACEI
<i>melliodora</i>	.21	354±220	yellow box	10-30	8-25	500	SL	ACDEIRUV
<i>microcarpa</i>	.35	730±616	grey box	10-25	10-15	400	SCL	BCDEHIV
<i>microcorys</i>	.15	200±139	tallowwood	25-40	10-20	1000	SLG	BCDEI
<i>muelleriana</i> * (cv)	.13	55±37	yellow stringybark	15-50	12-30	700	CL	CDFHIJV
<i>myriadena</i> subsp. <i>myriadena</i>	.23	850	snap and rattle	6-10	4-8	250	CL	BCIK
<i>newbeyi</i>	.29	200	Newbey's mallet	6-10	5-8	400	L,C	ACIJ
<i>nicholii</i>	.89	491±196	narrow-leaved peppermint	12-16	5-12	700	CL	CRU

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
'nutans'* (mallee form)	.23	150	red-flowered moort	2.5-4	3-5	400	CL	ABIV
'nutans' (mallet form)	.45		red-flowered moort	6-10	3-5	350	L,CL	ACI
obliqua*	.19	88±47	messmate stringybark	4-90	6-35	500	L	ACDISV
obtusiflora	.21		dongara mallee	2-8	2-8	300	S,SL	ABIJ
occidentalis var. occidentalis*	.19	213±92	flat-topped yate	12-20	5-10	380	SL	CEHIKV
occidentalis var. occidentalis (pure seed)	9.35	1000						
occidentalis var. stenantha			see <i>Eucalyptus aspratilis</i>					
oldfieldii	.21	45	Oldfield's mallee	3-8	4-8	200	S	ABEI
oleosa var. oleosa*	.27	169±76	red mallee	5-12	4-10	200	SL	BEFIK
oleosa var. oleosa (pure seed)	9.80	2500						
oraria	.81	224	ooragmandee	2-6	2-5	400	SL	BJ
orbifolia*	.39	284	round-leaved mallee	3-6	3-8	250	SL	AIMUV
oreades	.23	87±35	Blue Mountains ash	20-30	10-20	750	CL	CDE
ovata	.37	546	swamp gum	8-30	8-20	600	SC	CDHIJV
pachyloma	.19	6	Kalgan Plains mallee	2.5-5	4-6	500	S	AI
paniculata	.40	411±304	grey ironbark	20-30	10-15	750	SL,C	BCEU
patellaris	.21	107±55	weeping box	12-18	5-12	300	S	ABCW
patens	.17	45	WA blackbutt	10-45	15-25	850	SL	CDI
pellita	.35	170±125	large-fruited red mahogany	7-30	5-18	900	S	DIJV
pendens	.81	12	Badgingarra mallee	2-6	1.5-5	600	SG	AV
petraea	.23	540	granite-rock box	6-14	6-12	300	SL	B
phaenophylla subsp. phaenophylla	.17	180	mallee white gum	3-7	4-5	320	S	BI

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
phenax	.21	80	Kangaroo Island mallee	3-8	3-6	200	SLC	EIJ
pileata	.23	230	capped mallee	3-6	4-8	250	SL,CL	ABI
pilularis	.15	55±33	blackbutt	25-70	8-20	900	SL	CDI
platycorys	.21	40	boorabbin mallee	3-6	4-6	230	SCL	ABIK
platypus var. heterophylla	.21	150	coastal moort	4-6	5-7	400	S	ABIJTW
platypus var. heterophylla (pure seed)	3.85	1800						
platypus var. platypus*	.19	165	moort	5-10	6-10	400	CL	ABIJKV
platypus var. platypus (pure seed)	3.40	1800						
pluricaulis subsp. pluricaulis (SC)	.91	90		3-5	3-5	400	S,G	AB
pluricaulis subsp. porphyrea	.63	100	purple-leaved mallee	3-4	4-6	400	SL	ABIMT
polyanthemos	.21	465	red box	10-25	5-15	500	G	CEIJMRUV
polybractea	.29	835	blue-leaved mallee	4-10	4-8	470	S,SL,L	BLRU
praetamissa	.59	204	Beaufort Inlet mallet	9-15	5-10	500	SL	CJ
preissiana subsp. preissiana	.15	70	bell-fruited mallee	2-5	3-10	400	LG	AIJUV
propinqua var. propinqua	.17	465±230	small-fruited grey gum	20-40	10-25	850	CL	CDI
pterocharpa	.35	150		7-11		250	SCL	I
pulverulenta	.57	325	silver-leaved mountain gum	6-10	2-5	800	S,CL	MU
ptychocarpa			see <i>Corymbia ptychocarpa</i>					
punctata var. punctata	.25	90±82	grey gum	20-30	10-20	700	L	CDEF
pyriformis*	.19	35	pear-fruited mallee	2-6	4-6	350	S	AEIV

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>ravida</i>	.21	80	silver-topped gimlet	6-20	5-12	250	L,CL	ACFIKQ
<i>redacta</i>	.55	130	breakaway mallet	7-12	5-7	500	SL	EFW
<i>redunca</i>	.39	120±104		3-5	4-6	480	S	BIJT
<i>regnans</i>	.47	181	mountain ash	55-75	15-30	750	L	DG
<i>resinifera</i> (cv)	.27	222±182	red mahogany	20-45	10-15	800	SLG	CDGI
<i>rhodantha*</i> (cv)	.67	30	rose mallee	2.5-4	3-6	400	S	AIMV
<i>rigidula</i>	.23	200	stiff-leaved mallee	3-5	5-6	200	SL	ABI
<i>robusta</i> (cv)	.17	450±250	swamp mahogany	20-30	10-25	1000	SLC	CFHIV
<i>rubida</i>	.25		candlebark	10-40	10-20	600	L,GC	BCEI
<i>rudis</i> subsp. <i>cratyantha</i>	.17		WA flooded gum	10-20	8-13	800	SL,L	ACEHIW
<i>rudis</i> subsp. <i>rudis</i>	.19	570	WA flooded gum	10-20	8-13	500	SL	ACEHIW
<i>rudis</i> subsp. <i>rudis</i> (pure seed)	11.50	2500						
<i>salicola</i>	.41	290	salt gum	10-15	9-12	200	L	CEHK
<i>salicola</i> (pure seed)	9.65	6000						
<i>saligna</i>	.23	532±297	Sydney blue gum	20-55	10-25	900	SL,L	DEIV
<i>saligna</i> (pure seed)	5.35	2500						
<i>salmonophloia*</i>	.25	600±304	salmon gum	12-30	8-20	250	L	CEGUUV
<i>salmonophloia</i> (pure seed)	6.40	2800						
<i>salubris</i> var. <i>glauca*</i>			see <i>Eucalyptus ravida</i>					
<i>salubris</i> var. <i>salubris*</i>	.21	275±97	gimlet	6-20	5-12	200	L,CL	ACFIQKV
<i>salubris</i> var. <i>salubris</i> (pure seed)	4.45	2500						
<i>sargentii</i> subsp. <i>fallens</i>	.81	230	salt river mallee	2-4	4-6	300	SL,L,CL	BKV
<i>sargentii</i> subsp. <i>sargentii</i>	.35	220±83	salt river gum	6-12	5-8	350	SL	ABHIKQUVW

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
EUCALYPTUS (cont.)								
<i>sargentii</i> subsp. <i>sargentii</i> (pure seed)	9.75	1500						
<i>sepulcralis*</i>	.63	25	weeping gum	5-10	3-6	500	S	AJ
<i>sheathiana</i>	.25	400	ribbon-barked mallee	4-8	4-6	250	S,SL	ABIW
<i>sideroxylon</i> 'rosea'(cv)	.41	500	red-flowered ironbark	10-30	10-20	450	SGC	ACDEJLMRUV
<i>sideroxylon</i> subsp. <i>sideroxylon</i> (cv)	.21	226±166	red ironbark	10-30	10-20	450	GC	ACDEJLMRUV
<i>sideroxylon</i> subsp. <i>tricarpa</i>			see <i>Eucalyptus tricarpa</i>					
<i>sieberi*</i>	.21	111±32	silvertop ash	15-35	10-20	700	SC	CDGIV
<i>spathulata</i> subsp. <i>grandiflora</i>			see <i>eucalyptus suggrandis</i> ssp. <i>suggrandis</i>					
<i>spathulata</i> subsp. <i>spathulata</i>	.23	440	swamp mallet	6-12	6-8	300	SL,CL	ABHIJKMQRUV W
<i>staeri</i>	.15	15	albany blackbutt	6-15	4-10	500	S	BIM
<i>stoatei</i>	.21	38	scarlet pear gum	3-5	3-4	350	SL	AEIV
<i>stowardii</i>	.47	45	flute-horned mallee	3-8	4-6	300	S	AI
<i>striatocalyx</i>	.29	250	kopi mallee	3-12	3-10	150	SL	ACKV
<i>stricklandii*</i>	.17	145±65	Strickland's gum	6-12	5-8	230	L,CL	ABEIU
<i>subangusta</i> subsp. <i>subangusta</i>	.21	200		4-5	3-4	300	S	BI
<i>suggrandis</i> subsp. <i>suggrandis</i>	.23	160	swamp gimlet	3-5	4-6	300	SL,GL	ABHITVW
<i>talyuberlup</i>	.29	70	pretty yate	3-6	3-6	600	CL	ABIT
<i>tenuis</i>	.79		slender-stemmed mallet	4-8		250	CL,C	A
<i>tereticornis</i>	.21	642±407	forest red gum	20-30	10-25	650	S,GL	BCEU



Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>EUCALYPTUS (cont.)</b>								
<i>tetragona*</i>	.19	25	tallerack	2.5-8	3-8	400	S	AIJMUV
<i>tetraptera*</i>	.35	40	four-winged mallee	2-6	3-8	400	S	AU
<i>todtiana</i>	.15	2	coastal blackbutt	5-16	6-10	450	S	ABIJ
<i>todtiana (SC)</i>	.25	5						
<i>torquata*</i>	.19	91±35	coral gum	6-12	5-10	250	L	AEIJKU
<i>torquata (pure seed)</i>	3.50	800						
<i>transcontinentalis</i>	.23	114±45	redwood	3-20	3-10	230	SL	AEJKW
<i>tricarpa</i>	.51		red ironbark	25-35		450	S,SL,L, CL,G	CDF
<i>uncinata</i>	.29	805	hook-leaved mallee	1-4	1.5-5	380	SL,L	AEIJW
<i>viminalis</i> subsp. <i>cygnetensis</i>	.35	250	rough-barked manna gum	10-20	8-12	500	SL	ADEIR
<i>viminalis</i> subsp. <i>viminalis</i>	.23	349±237	manna gum	30-50	10-15	500	SL	ADEIRV
<i>vitrix</i>	.41		coolabah	10-15	8-15	200	S,SL	ACEH
<i>wandoo</i> subsp. <i>wandoo</i>	.23	300±100	wandoo	8-30	6-20	400	SLG	CDEFGIQV
<i>wandoo</i> subsp. <i>wandoo</i> (pure seed)	6.90	2500						
<i>websteriana*</i>	.51	175	Webster's mallee	3-6	3-6	250	LS	ADJ
<i>woodwardii*</i>	.19	125	lemon-flowered gum	6-15	3-8	240	SL	AIKU
<i>woodwardii</i> X <i>torquata</i>	.33	30	torwood	6-12	5-10	250	L	AEIJU
<i>woollsiana</i>	.29	1077±77	grey box	10-25	8-12	400	CL,SL	CEGI
<i>xanthonema</i> subsp. <i>xanthonema</i>	.25	280	yellow-flowered mallee	2-4	2-4	400	SL	BITW
<i>yilgarnensis</i>	.27		yilgarn mallee	4-5		250	CL	AK
<i>youngiana</i>	.29	44	yarldarba	4-12	5-12	200	S,SL	AEIV
<i>zopherophloia</i>	.55	94	blackbutt mallee	4-6	4-6	400	S,SL	BJ

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>GOMPHOLOBIUM</b>								
<i>marginatum</i>	3.37	450		0.2-0.6	0.3-1	600	S,SL	PU
<i>tomentosum</i>	1.89	670		0.6-1.3	0.5-1	500	S,SL	P
<b>HAKEA</b>								
<i>amplexicaulis</i>	4.70	20		1-3	1.3-3	600	S,SLG	AU
<i>baxteri</i>	4.70	15	southern fan-leaf	1.5-5	1-3	400	SG	ATU
<i>brooksiana</i>	1.89	25	Brook's hakea	1-2	1-2	300	S,LS	KT
<i>bucculenta*</i>	1.47	60	red pokers	2.5-7	2-6	300	S,L	ABTUW
<i>ceratophylla</i>	3.10	45	staghorn hakea	0.8-2.5	1-3	800	S,LS	MTU
<i>coriacea</i>	1.52	100	pin spike hakea	4-7	2-5	300	SLG	AIT
<i>corymbosa</i>	1.58	55	cauliflower hakea	1-2	1.2-3	400	S,SL,L	ABJTS
<i>cyclocarpa</i>	1.58	15	curved-fruit hakea	1-2.5	1-3	700	SG	T
<i>falcata</i>	2.40	80	sickle hakea	1-3	0.5-2	600	SL	BHT
<i>flabellifolia</i>	1.58	6	wedge hakea	0.6-1.3	0.6-1.5	500	S	A
<i>invaginata</i>	1.89	240		1-2.5	0.6-2	300	S,LS	T
<i>lasianthoides</i>	4.12	72	woolly-flowered hakea	2-5	1.5-4	800	SL	ABUV
<i>laurina</i>	1.01	54	pincushion hakea	3.5-8	3-10	500	C,G,S	ABMTW
<i>lehmanniana</i>	4.70	119	blue-flowered hakea	0.6-1.5	1-2	400	S,SL,G	AT
<i>lissocarpa</i>	2.00	42	honeybush	0.6-2	1.5-3	800	SG	AIU
<i>multilineata*</i>	1.58	70	grass-leaved hakea	3-5	3-7	250	S,GL	AMUW
<i>neurophylla</i>	2.34	30	pink-flowered hakea	1-2.5	1-2	500	S,G	AU
<i>oleifolia</i>	2.50	77	olive-leaved hakea	6	3	800	S,L	ABJTUW

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>HAKEA (cont.)</b>								
<i>orthorrhyncha</i>	2.84	33	birds beak hakea	1.5-5	1.5-4	300	S,SL	ABT
<i>pandanincarpa</i>	.81	12	screw-pine hakea	2-5	2-6	400	SL	BMT
<i>petiolaris</i>	2.34	63	sea-urchin hakea	1.5-10	2-6	300	G	AMTU
<i>platysperma</i>	.91	2	cricket ball hakea	1.5-3	2-4	350	S,G	T
<i>prostrata</i> (upright form)	2.46	20	harsh hakea	4	1.5	300	S	AMSTU
<i>ruscifolia</i>	3.80	66	candle spike hakea	0.5-2	1-2	400	S	ASTU
<i>undulata</i>	1.52	90	wavy-leaved hakea	1.5-3.5	1.5-3	600	S,GC	AMU
<i>varia</i>	2.00	68	variable-leaved hakea	1-4	1.5-4	700	S,SL	BHT
<b>HARDENBERGIA</b>								
<i>comptoniana</i>	.19	40	native wisteria	Climber	N/A	800	S,L	AJT
<b>HOVEA</b>								
<i>chorizemifolia</i>	3.67	50	holly-leaved hovea	0.5-1.5	0.3-0.8	700	S,G	AP
<i>elliptica</i>	.17	54	tree hovea	2-4.5	1.5-3.5	1000	SL	AT
<i>trisperma</i>	.73	84	common hovea	0.3-0.6	0.3-1	900	S	A
<b>KENNEDIA</b>								
<i>carinata</i>	.37	70		Runner	1-2	900	SL	AP
<i>coccinea</i>	.19	40	coral vine	Climber	N/A	550	SLG	APV
<i>eximia</i>	.49	125	fine red creeper	Runner		400	S,SL	AP
<i>prostrata</i>	.17	45	scarlet runner	Runner	1.5	300	SLG	AP

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
<b>KUNZEA</b>								
<i>baxteri</i>	.21	2500	Baxter's kunzea	1-2	2-3	600	SG	ATV
<i>ericifolia</i>	.41		pondil	2.5-5	2-3.5	700	S,SL	HMRU
<i>pulchella</i>	.60		granite kunzea	2-4	2-5	250	S,SL	AMU
<b>LAMBERTIA (Price per 100 seeds)</b>								
<i>ilicifolia</i>	14.00	75	holly-leaved Lambertia	1-1.5	0.6-1.5	400	S,SL,G	AT
<b>LEPTOSPERMUM</b>								
<i>erubescens</i>	.29	1500	pink tea tree	2-3	2	400	SLG	BJ
<i>laevigatum</i>	.07	500	coastal tea tree	3-4	2	500	S	BEIJV
<i>spinescens</i>	.25		spiny tea tree	0.6-2	0.5-1.5	400	S	B
<b>LOPHOSTEMON</b>								
<i>conferta</i>	.07	150	Queensland brush box	35-40	8	900	S,L	DIV
<b>MELALEUCA</b>								
<i>acerosa</i>	.23	2500	coastal honey-myrtle	0.6-3.5	2-5	400	S	ABJW
<i>acuminata</i>	.19	6230	scented honey myrtle	1-3	2-3	400	L	BHKS
<i>adnata</i>	.45	5400	hummock honey-myrtle	1-2.5	1-2	250	SL,L	ABTU
<i>armillaris</i>	.13	1550	bracelet honey-myrtle	7-8	5-6	450	SL	ABEHK

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
MELALEUCA (cont.)								
<i>brevifolia</i>	.25	2300	mallee honey-myrtle	2-5	1.5-3	400	S,SL,L	BKW
<i>cardiophylla</i>	.23	720	umbrella bush	1-3	1-3	250	S,SL,L	JT
<i>cucculata</i>	.27	700		2-3	1-2	250	CL	A
<i>cuticularis</i>	.27	2500	saltwater paperbark	6-12	8	500	SL	BCHKUW
<i>densa</i>	.33		lemon honey-myrtle	0.6-3	0.6-3	500	S,G,C	H
<i>depressa</i>	.29			1-2.5	1-2.5	300	S,LS	AMTU
<i>diosmifolia</i>	.27	2100	green honey myrtle	1-1.5	1.5-4	500	S	ABH
<i>elliptica</i>	.33		granite honey myrtle	2-4	2-5	250	SL	ABTU
<i>fulgens</i> subsp. <i>fulgens</i>	.31		scarlet honey-myrtle	1-3	1.5-3	200	S,L,G	A
<i>glaberrima</i>	.29	3000	mauve honey myrtle	1-2.5	1-2.5	300	SL	ABT
<i>hamulosa</i>	.29	4900	creekline honey myrtle	3-4	2-3	500	C	BHK
<i>holosericea</i>	.47	3400	silky honey-myrtle	1-2	1-2	300	S,SG,CL	A
<i>huegelii</i>	.29	500	chenille honey-myrtle	3-5	1.5-3	300	S	ABJKMV
<i>hypericifolia</i>	.13	4800	hillock bush	3-6	2-3	450	SL	AHJMRTV
<i>incana</i> subsp. <i>incana</i>	.27	3500	grey honey-myrtle	2-3	2	600	SL	ABHJKMU
<i>lanceolata</i>	.17	1100	Rottnest tea-tree	5-8	3-5	300	S	BCHIKTV
<i>lateriflora</i> subsp. <i>lateriflora</i>	.45	1600	oblong-leaf honey-myrtle	1-6	1-4	300	SL,CL	ABKMTU
<i>lateritia</i>	.45	8500	robin redbreast bush	2-2.5	2-3	500	SL	ABHV
<i>leucadendra</i>	.41	1600	broad leaf paperbark	10-30	8-25	700	SL,L,CL	ACHJW
<i>nesophila</i>	.27	500	showy honey-myrtle	3-4	2-3	600	S	ABHKV
<i>pauciflora</i>	.55		mattress myrtle	0.2-1	0.8-2	800	SL	HJ
<i>pauperiflora</i> subsp. <i>fastigiata</i>	.29	260	inland honey-myrtle	3-5	3-5	300	SL	AB
<i>pentagona</i>	.35	5900	oval-leaf honey-myrtle	1-3	2-4	400	S,SL	AT

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
MELALEUCA (cont.)								
<i>preissiana</i>	.27	4800	paperbark	6-10	4-6	400	SL	ACH
<i>quinquineria</i>	.29	3200	broad-leaved paperbark	8-25	6	500	SL	ACHIJKV
<i>radula</i>	.43	2500	graceful honey-myrtle	1-3	1-2	400	S	AKV
<i>rhaphiophylla</i>	.81	2900	swamp paperbark	7-10	4-6	700	SL	BHM
<i>scabra</i>	.47	1500	rough honey-myrtle	1	2	800	SL	A
<i>sheathiana</i>	.25	1450	buri	3-5	2-3	250	SL	BT
<i>spathulata</i>	.27	2127	purple honey-myrtle	0.5-1	0.5-1	400	SL	ABK
<i>spicigera</i>	.41	9400		1-2	1-2	300	SL	A
<i>strobophylla</i>	.27	3350		1.5-12	1.5-8	250	SL,L	ABHKT
<i>subfalcata</i>	.55	4550	sickle-leaved honey-myrtle	2-4	2-3	250	SL	AB
<i>thymoides</i>	.35	850	sand wattle-myrtle	1	2	600	S	AJK
<i>thyoides</i>	.53	1770	salt lake honey myrtle	2-4	2-4	250	CL	KW
<i>uncinata</i>	.33	1420	broombush	1-3	2	250	SL	AHU
<i>uncinata</i> (broad leaf form)	.35	5000	broombush	1-3	2	250	SL	AHU
<i>viminea</i> subsp. <i>viminea</i>	.19	3500	mohan	3	3	400	SL	ABH

#### NUYTSIA

<i>floribunda</i>	.21	12	WA Christmas tree	6-10	3-4	750	S,LG	A
-------------------	-----	----	-------------------	------	-----	-----	------	---

#### PINUS

<i>halepensis</i> var. <i>halepensis</i>	.11	61	aleppo pine	15-24	4-5	650	LS,L	BDJ
<i>halepensis</i> var. <i>brutia</i>	.11		calabrian pine	10-25	4-5	400	LS,L	BD
<i>pinaster</i> (Plantation grade)	.15	22	maritime pine	27-36	4-10	510	S,SL	BCDW

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
PINUS (cont.)								
<i>pinaster</i> (Seed Orchard) F1.5	.41	22	maritime pine	27-36	4-10	510	S,SL	BCD
<i>pinea</i>	.10	1.2	stone pine	15-25	4-5	350	S,SL	ABC
<i>radiata</i> (Plantation grade)	.15	29	Monterey pine	25-45	6-12	760	SL,L	BCD
<i>radiata</i> (Seed Orchard) F1.5	.41	29	Monterey pine	25-45	6-12	760	SL,L	BCD
<i>taeda</i>	.13	40	loblolly pine	27-33	6-8	800	S,SL	HCD

#### PITOSPORUM

<i>phylliraeoides</i>	.17	50	weeping pittosporum	4-5	3-4	250	SL	ACW
-----------------------	-----	----	---------------------	-----	-----	-----	----	-----

#### SANTALUM

<i>acuminatum</i>	.03	0.35	quandong	4-5	2-3	510	LGS	ABX
<i>album</i> (cv)	.50		indian sandalwood	8-10	3-5		L,CL	AL
<i>spicatum</i>	.05	0.21	sandalwood	2-8	1-3	200	L	AULX

#### SENNA

<i>hamersleyensis</i>	.39	52		0.3-0.5	1	250	S	AW
<i>notabilis</i>	.09	58	cockroach bush	1-2	1	250	S	AW

#### STRANGEA

<i>cynanchocarpa</i>	.25	4		0.5-1	1-2	500	S	PW
----------------------	-----	---	--	-------	-----	-----	---	----

Species	\$/g	Average seeds/g	Common name	Average Height (m)	Average Width (m)	Min Rainfall (mm)	Soil Type	Uses
---------	------	-----------------	-------------	--------------------	-------------------	-------------------	-----------	------

#### TEMPLETONIA

<i>retusa</i>	.21	32	cockie's tongues	1-2	2-3	500	SL	AJV
---------------	-----	----	------------------	-----	-----	-----	----	-----

#### XANTHORRHOEA

<i>gracilis</i>	.29	60	mimidi			850	SG	A
<i>preissii</i>	.19	62	balga (blackboy)	0.3-3	1-2	700	S,SL,L,G	A

#### XYLOMELUM (Price per 100 seeds)

<i>angustifolium</i>	10.00	2.4	sandplain woody pear	2-4	1.5-3	350	S	AM
<i>occidentale</i>	10.00	2	woody pear	4-8	2-3	700	S,SL	AM

## POSTAGE AND PACKING CHARGES WITHIN AUSTRALIA

### Regular Parcels

Seed Weight	Within 50 km	South & Mid WA	North WA (6700-6799)	SA	NSW	Northern Territory	Melbourne	Other Vic	Tasmania	Qld
	\$	\$	\$	\$	\$	\$	\$	\$	\$	\$
Up to 200g Over 200g to 450g	3.30 4.50	3.30 4.50	3.30 4.50	3.30 4.50	3.30 4.50	3.30 4.50	3.30 4.50	3.30 4.50	3.30 4.50	3.30 4.50
Over 450g to 900g Over 900g to 1.9g	6.30 6.30	6.70 6.90	6.70 7.60	9.60 11.90	9.95 12.60	9.95 12.60	9.60 11.90	9.95 12.60	9.95 12.60	9.95 12.60
Over 1.9g to 2.9g Over 2.9g to 3.9g	6.30 6.30	7.10 7.30	8.60 9.30	14.50 16.50	15.55 18.10	15.55 18.10	14.50 16.50	15.55 18.10	15.55 18.10	15.55 18.10
Over 3.9g to 4.9g Over 4.9g to 5.9g	6.30 6.30	7.50 7.70	10.60 11.30	18.50 20.50	20.45 22.80	20.45 22.80	18.50 20.50	20.45 22.80	20.45 22.80	20.45 22.80

<b>EXPRESS POST</b>	
Seed Weight	All States \$
Up to 450g	5.00
Over 450g to 3kg	7.50

FOLD OUT FOR KEY



SOIL TYPE		KEY TO USES & FEATURES	
S	Sand	A	Ornamental plantings for parks and gardens
SL	Sandy Loam	B	Hedges or windbreaks
L	Loam	C	Shade trees
LS	Loamy sand	D	Sawmilling timber
C	Clay	E	Firewood
G	Gravel	F	Small timber for poles, rails etc
		G	Durable timber suitable for contact with the ground
<b>ABBREVIATIONS</b>		H	For planting in waterlogged areas
cv.	cultivated	I	Honey-producing plants
subsp.	subspecies	J	For planting on coastal areas
var.	variety	K	For planting in saline soils
sc.	special collections (see page 1)	L	Essential oils
		*	Sensitive to dieback
		M	Useful for cut flowers, foliage or fruit
		N	Stock fodder
		O	Gum or resin products
		P	Mat and ground-cover plants
		Q	Tan bark (products rich in tannin)
		R	Perfumed foliage
		S	Perfumed flowers
		T	Screening plants
		U	Frost-tolerant plants
		V	Wind-resistant plants
		W	Soil stabilisation
		X	Edible fruits