

# LIBRARY

Department of Biodiversity,  
Conservation and Attractions

This PDF has been created for digital preservation. It may be used for research but is not suitable for other purposes. It may be superseded by a more current version or just be out-of-date and have no relevance to current situations.

## Lake Altham Waterbirds


Lake Altham is a shallow naturally saline wetland located 35 km south of Lake Grace in the chain of wetlands that includes Lakes Grace, Pingrup and Chinocup. The wetland has a history of waterbird data and may be an important waterbird site when the lake is full.



*View across the shallow Lake Altham (photo by S.A.Halse)*


### Species Richness

Since monitoring began, Lake Altham has not been a site of great importance to waterbirds in the wheatbelt. A total of 7 species have been recorded at relatively low abundances. However, prevailing rainfall conditions have meant that the lake has not held water after late winter except in 1998 when the lake dried after spring sampling.



*Species richness at Lake Altham.*

Multi-dimensional scaling Ordination (SSH) of waterbird species abundance indicates that the waterbird assemblage at lake Altham is different from most monitoring wetlands and differs substantially from year to year. Because the annual waterbird assemblages are comprised of only a small number of species, observed differences are principally the result of changes in the relative abundances of species. There is no apparent trend from year to year.



*MDS Ordination (SSH) of range standardized abundance of waterbird species.*

### Further Reading

Cale, D.J., S.A.Halse and C.D.Walker (2004) Wetland monitoring in the Wheatbelt of Western Australia: site descriptions, waterbird, aquatic invertebrate and groundwater data. *Conservation Science W. Aust* 5: 20-135


Halse, S.A., D.J. Cale, E.J. Jasinska and R.J. Shiel (2002) Monitoring change in aquatic invertebrate biodiversity: sample size, faunal elements and analytical methods. *Aquatic Ecology* 36: 1-16

Jaensch, R.P., R.M. Vervest and M.J. Hewish (1988) Waterbird surveys of wetland nature reserves in south-western Australia: 1981-85. Report No. 30, Royal Australasian Ornithologists Union, Melb.


# Salinity Action Plan Wheatbelt Wetlands Monitoring Programme

## Wheatbelt Wetlands Monitoring


The Salinity action Plan Wheatbelt Wetlands Monitoring programme commenced in 1997 with the sampling of five wetlands and was expanded to include 13 in 1998 and finally a total of 25 wetlands in 1999. These monitoring wetlands have been sampled every second year since commencement, such that half of the wetlands are sampled in alternating years. Wetlands first sampled in 1997 have now been sampled 4 times. While this actually yields few data points and interpretation is, at this stage, imprecise it is expected that as the project continues and further data points are collected an increasingly accurate estimate of wetland trends will be achieved. Faunal sampling includes; waterbird species richness and abundance, aquatic invertebrate species richness and abundance and water-chemistry. Sampling of these parameters is directed toward tracking trends in biodiversity of the wetlands individually and as a group to reflect the status of wheatbelt wetlands generally. This brief note presents data for waterbird surveys up to 2003 and is intended as an annual mechanism for reporting data from this project.



**MEAN Z-score for WATERBIRD RICHNESS at all lakes during year of sampling**



**MEAN Z-score for WATERBIRD ABUNDANCE at all lakes during year of sampling**



## Waterbird Richness and Abundance in the Wheatbelt

The number of species present (Richness), is a valuable measure of biodiversity and abundance is indicative of the productivity of wetlands. The mean z-score for waterbird richness and abundance is calculated in the same way. At each wetland the normal deviate (z) is calculated for each year, from the entire dataset for that wetland. The mean z-score is the average of these annual z scores over all wetlands. Thus, the mean z-score can be used to measure the overall trend in monitored wetlands over time. Values below zero reflect lower than average species richness or waterbird abundance. Over the period of monitoring there has been a decline in waterbird species richness and abundance at the monitored wetlands. This has coincided with a decline in rainfall following 1999 with 2000-2002 showing 'average' to 'very much below average' rainfall over the study area. It is too early to ascribe the reduction in species richness and abundance to low rainfall, although it is intuitive that the lower water levels and higher salinities associated with low rainfall are likely to lead to fewer species using wetlands.