

LIBRARY

Department of Biodiversity,
Conservation and Attractions

This PDF has been created for digital preservation. It may be used for research but is not suitable for other purposes. It may be superseded by a more current version or just be out-of-date and have no relevance to current situations.

Fungus of the Month - December 2011



© Richard Robinson

Hypomyces chrysospermus

Species of *Hypomyces* are common parasites of other fungi. They appear to be host specific or infect closely related species. *Hypomyces chrysospermus* infects bolete fungi and their relatives, and is commonly called the bolete eater. Boletes are fungi that produce mushrooms with a layer of pores on the underside of the cap (instead of the more usual gills).

The life cycle of *H. chrysospermus* is complex but intriguing, with three distinct stages. The first stage is a white cottony mould that engulfs the bolete mushroom. Next a distinctive bright yellow powdery coating develops (*left*). The white and yellow stages are anamorphic, or asexual, forms of the parasite. The final stage is a well formed but delicate creamy grey to light khaki skin-like covering embedded with small dark pimples (*right*). The pimples are the necks of tiny flask-like structures (*see inset*) which are the actual fruit bodies of the parasite and where the sexual spores are produced. The final stage is rarely seen, or more often ignored, because it generally develops after the host has decayed beyond recognition and has become slimy and very smelly.

The scientific name refers to the parasitic nature and distinctive asexual stage of the fungus. *Hypo*:- below, or under, *-myc*: fungus, mushroom, *chrys*:- golden, *sperm*:- seed.

Richard Robinson, DEC Science Division, Manjimup



Department of
Environment and Conservation

