

LIBRARY

Department of Biodiversity,
Conservation and Attractions

This PDF has been created for digital preservation. It may be used for research but is not suitable for other purposes. It may be superseded by a more current version or just be out-of-date and have no relevance to current situations.



Ramaria ochraceosalmonicolor

Richard Robinson, Science Division, Manjimup, richard.robinson@dec.wa.gov.au



© Richard Robinson

Ramaria ochraceosalmonicolor does not have a common name, but could be referred to as the salmon coral fungus. However, it does vary considerably in colour from various shades of yellow through to orange. The salmon coral fungus is common in eucalypt forests and woodlands throughout southern Australia, appearing amongst the litter from early to late autumn. It is conspicuous, not only by its colour and coral-like shape, but also by its size.

Several fruit bodies generally develop together and can form clusters up to 20cm in diameter and 15cm tall. The colour tends to be more intense in young specimens. The branching pattern is generally irregular, but the tips of branches tend to be rounded. The base of the stems is nearly always white.

There are many species that belong to the group commonly referred to as coral fungi. Some species form simple or branched, club-like structures (see *Clavaria miniata*; Factsheet 32/2009), the salmon coral fungus has a dense but open coral-like structure, and other species may be more compact and cauliflower-like.

The scientific name refers to its shape and colour. *Ram-*: branch, *-aria*: pertaining to or possessing, *ochreac-*: ochraceous, *salmon-*: salmon coloured.