

# LIBRARY

Department of Biodiversity,  
Conservation and Attractions

This PDF has been created for digital preservation. It may be used for research but is not suitable for other purposes. It may be superseded by a more current version or just be out-of-date and have no relevance to current situations.



## *Hydnum repandum* — hedgehog fungus

Richard Robinson, Science Division, Manjimup, [richard.robinson@dec.wa.gov.au](mailto:richard.robinson@dec.wa.gov.au)



*Hydnum repandum* produces a mushroom-like fruit body with spines on the underside of the cap rather than gills. The spines and the fact that it occurs throughout the world, gave rise to the British common name of hedgehog fungus. In Western Australia, *Hydnum repandum* is found in karri forest, the wetter regions of jarrah forest and coastal woodlands.

Caps are up to 6cm diameter often irregular in shape. They may be circular and broadly convex becoming flat with a wavy margin or irregular with a lobed and wavy margin. The surface is dull greyish orange and generally dry to moist. Spines are creamy-white, about 3mm long and sometimes extend a short distance down the stem. Stems are up to 4cm long, dull orange with a white powdery coating, which is easily removed by rain or when handled.

Often this fungus fruits in groups in and under litter on the forest floor, and may be quite small and misshapen compared to the specimens photographed. There is also a similar chestnut or reddish brown unnamed species (*above inset*) found in southern Australia, including the southwest of Western Australia.

The scientific name refers to the habit and appearance of the mushroom. *Hydn-*: is the ancient term for an edible mushroom, *hydroid* is also the technical term for toothed, *repand-*: folded backwards or upturned.