




FOREST FOCUS

NUMBER 21 APRIL 1979



**DIEBACK HYGIENE
ECOLOGICAL APPROACH TO DIEBACK CONTROL**



▲ *Once the disease has progressed to the stage of overstorey collapse it is highly unlikely that any cultural techniques will be able to restore native forest. (See article on page 7.)*

(Les Harman)

Back Cover

A three-year-old stand of shrub and tree species which were direct seeded on to a bauxite mine site. Legumes have been used to stabilise the mine surface to prevent erosion, but it is

possible that they could be used to suppress jarrah dieback and permit re-establishment of jarrah on bauxite pits.

Les Harman

DIEBACK HYGIENE

... first steps

There can be plenty of debate about forest disease risk areas being known as Quarantine. The word is probably misused in this context, but it has been misused to good effect.

Forest disease risk areas were proclaimed for a number of reasons:

- To give time for the symptoms of dieback to become visible . . . a process that can take up to three years to occur.
- To gain a pause in the spread of the fungus in areas of low infection by machinery and activity, so that methods for operating in the forest without fungal spread could be devised.
- To allow more time for promising laboratory research findings to be tested under field conditions.
- To "buy" time for the evolution of photographic techniques for detection of the disease in its early stages and in small pockets of infection.

Control of vehicle movement for a period of three or more years was the first step. The results of this exercise are encouraging, and although they are partly due to energetic implementation of access controls, the greatest achievement has resulted from tremendous public co-operation.

Success rate

On the 14th of January, 1976, the results of a year's planning fitted into place. Detailed mapping, planning, discussion with forest users, preliminary job descriptions and sites for warning signs had been prepared, and within 48 hours the entire northern Forest Disease Risk Area was established on the ground. A similar programme was executed for the second or southern area and was expedited in similar vein in February 1978.

The creation of verbal and physical constraints to forest entry is one thing . . . the success of this action is another.

Necessary access to quarantined areas is controlled by an entry permit, but non-permit or illegal penetration required special treatment. Control of illegal entry is achieved by a combination of ground patrol, continuing publicity, and aerial reconnaissance. Although it is not possible to exclude all unwanted vehicles from the risk area, data collated from patrol of key areas and a comparison with pre-quarantine visitor rates showed that a success rate of up to 90 per cent was achieved. Of course there have been exceptions, and the philosophy of enforcement by education and persuasion has been gradually hardened toward illegal entry of a persistent nature. Nevertheless, the department is gratified with the ready acceptance of quarantine and its constraints, by the vast majority of forest users.

WHAT DID THE TIME BUY?

Jarrah dieback research

Studies of the suppressive effect of native leguminous species on *Phytophthora cinnamomi* have been the major area of research. In two field studies, significant suppression of sporulation under understories of *Bossiaea aquifolium* and *Acacia pulchella* was recorded; this was the first field demonstration of this

FOREST FOCUS

Number 21, April 1979

Published for Mr. B. J. Beggs, Conservator of Forests, Forests Department of Western Australia, 54 Barrack Street, Perth.

Articles in this publication may be freely reprinted—preferably with acknowledgement, except in the case of commercial promotion material, when permission *must* be obtained.

Typesetting and page preparation by Filmset.

Offset plates by Art Photo Engravers Pty. Ltd.

Printed in Western Australia by the Government Printing Office.

AT ISSN 0049-7320

Compiled by Dale Watkins



New Minister Appointed

Following a re-allocation of portfolios in State Cabinet, the Hon. David Wordsworth was appointed Minister for Forests and Lands. He was previously Minister for Transport.

Mr. Wordsworth was elected Member of the Legislative Council seat of South Province in 1971, and subsequently represented Western Australia in an Australian Inter-Parliamentary Delegation invited to Britain to confer with Members of Parliament at Westminster on trade, defence and other matters.

He has always maintained a strong interest in civic affairs and economic factors concerning primary producers. He has had printed a booklet on the devaluation of the Australian dollar and its effects on farming and exports.



▲ *Utility entering the fungicide dip.* (Les Harman)

effect. Parallel laboratory and greenhouse studies have continued in an attempt to isolate the factors responsible for the suppression. (See page 7.)

Hygiene techniques

There is abundant proof that the use of machines in dieback affected forest will assist the fungus to spread into unaffected areas. There is also a direct relationship between size of machine, type of traction, season of the year, and the risk of new infection. Heavy, tracked machines in winter are the worst combination, while light, rubber-tyred machines

in summer pose the least risks. These facts, coupled with research findings by Mr. Frank Batini and others, have influenced the direction of research into hygienic methods of bush operations.

Operational studies are in various stages of completion with respect to washing down of vehicles, the practical, social and economic problems of summer stockpiling of forest produce, evolution of hard surfaced, disease free roads and the definition of the connection between climate and fungus sporulation.

It has been found that firewood, posts and S.E.C. poles can be cut and prepared in winter and extracted

in the dry months of the year, while the smaller sawmills at least, can effectively stockpile sawlogs in summer on a scale which allows for no winter cutting. Further refinements will be called for, and the larger mills still have to overcome practical problems, especially related to space at mill landings.

It has been shown that use of a high pressure water jet to remove soil particles from vehicles is a simple and effective precaution against the spreading by vehicles of infected material. A few simple precautions are needed with respect to planned drainage into a fungicide tank or an already diseased stream. There are



fixed washdown stations at all relevant Forests Department headquarters, and temporary washdown locations have been selected for important field sites.

Trials with a fungicide dip in a concrete basin or ford are continuing, and the technique appears to have potential for light vehicles with relatively small mud deposits on tyres and suspension.

Road maintenance and classification

The problems of summer stockpiling will not be easy to overcome, but it may be possible to extend the

“safe” logging period by careful classification of roads into “safe” and “unsafe” categories, and by the accurate pegging of affected and unaffected borders or boundaries according to a standard colour coding.

Small hardwood posts are erected at the perimeter of infections, painted yellow on the side facing into dieback and green on the side facing into healthy forest.

Routine grading and other road maintenance or repair work must not move from one type into the other without making sure that infected soil is clear of the vehicle.

▲ Wash-down station at the Department’s Dwellingup Division Headquarters.

(Les Harman)

“Eye Spy”

(colour dieback film)

It has been mentioned that the time lapse from new infection with *Phytophthora cinnamomi* until the first plant symptoms are seen can be up to three years, and that the Forest Disease Risk Areas were proclaimed in order to allow existing but undetected infections to emerge. During the period of quarantine, a new technique using 70 mm colour aerial


photography has been evolved which extends the scope of aerial detection of the disease.

The photography requires special techniques as does the interpretation of the photographs obtained:

- A 70 mm colour film in a Vinten camera is exposed at a height of 500 metres above the ground.
- Only days of complete high cloud cover can give the desired shadow free detail required.
- Plotting of flight lines requires a highly sophisticated aerial navigation method involving a computer built into the small aircraft used.
- A wide net has been needed to locate people who can effectively interpret the great detail revealed by these photographs.
- Photography will identify "spot infections" (individual blackboys, banksia trees, etc.) only in the absence of shadows and in good lighting conditions. The presence of shadows or burning within the previous three years drastically reduces their value.

Stereoscopy is the method used to obtain a three-dimensional image from a pair of photographs. In real life two eyes enable perception of distance or depth of a scene because each eye sees a slightly different view. In the use of aerial photographs, an adjacent pair—a stereo pair—of photographs is viewed simultaneously with both eyes. Where the pictures overlap an image of the third or vertical dimension is produced.

This photographic pair should be viewed by holding a piece of cardboard vertically along the join line and looking with one eye either side of the card. Most people can see in stereo but some practice may be needed to get the most out of these.



An Ecological Approach to the Control of Jarrah Dieback

by Dr. S. R. SHEA

Jarrah dieback is a plant disease caused by the introduced soil-borne fungus *Phytophthora cinnamomi*. This pathogen has the capacity to kill jarrah (*Eucalyptus marginata*) and many of the components of the forest almost throughout its geographic range. (See *Forest Focus* No. 14.)

In addition to the loss of timber, recreation and conservation values, the disease poses a threat to the major water supply system of the south-west of Western Australia. (See *Forest Focus* No. 19.)

Large areas of the forest have been quarantined to restrict spread of the fungus in contaminated soil and to permit detailed mapping of the distribution of the disease. Quarantine and/or stringent hygiene does provide short-term control but in the long term, particularly if access to quarantined forest areas is to be permitted, methods of controlling the fungus other than by its exclusion must be developed.

Options for control

Despite considerable advances in plant pathology research in the last decade, soil-borne plant pathogens cause massive crop losses throughout the world each year. In some agricultural or horticultural situations where it is economically feasible to practice intensive cultural techniques, control of some soil-borne plant pathogens has been achieved.

In forests, however, such techniques are usually not practicable. For example, it is relatively easy to kill *P. cinnamomi* with a fungicide, provided the fungicide can contact the fungus. In a diseased area of jarrah forest many fungal spores occur within thick roots and the soil would have to be frequently saturated with a fungicide before contact with the fungus occurred.

Even broad scale applications of fungicide in the jarrah forest would not eliminate all fungal spores and, particularly in catchments, fungicidal application could have damaging side-effects. Even if one spore remained viable in a treated forest area it is theoretically possible for the fungus to build up to pre-treatment levels.

Resistance to some plant diseases has been achieved by selection and breeding of resistant trees or plants. However, there is little evidence that there are strains of jarrah which are resistant to jarrah dieback. Even if successful, the development of resistant strains of jarrah would be very slow because of the relatively low growth rate of the species and the necessity to conduct extensive long-term field tests before resistance could be confirmed.

Replacement of jarrah with an alternative resistant tree species would partially compensate for the loss of some forest values. Extensive tree planting trials have shown that there are a number of eucalyptus and pine species which are resistant to the disease. In the valleys, where the disease is most prevalent and where the early trials are located, it is possible to grow some of these species to maturity where the soil moisture regime is favourable. It is only recently that trial plots containing a large number of tree species

have been established on dieback-infested upland forest sites. Upland sites, which represent between 60-80 per cent of the forest, have harsh conditions for tree growth and it is not possible at this stage of research to specify which species could be successfully used to replace jarrah. Rehabilitation of diseased forest areas with alternative tree species is an option, but further research is necessary before it will be possible to undertake broad scale rehabilitation. Even when the correct techniques for rehabilitation have been developed, replacement of the jarrah forest with alternative tree species would be very expensive.

Direct attack on the pathogen or improvement or replacement of the susceptible host are not the only methods by which a plant disease can be controlled. There are three components of a plant disease—a pathogen, a susceptible host and an environment suitable for the pathogen to attack and damage the host. It is possible for a pathogen and a susceptible host to co-exist if environmental conditions are unsuitable for the pathogen. Thus, another control option is to change environmental conditions in such a way as to inhibit fungal pathogenicity. In this article a research programme aimed at determining if it is possible to manipulate the jarrah forest environment so that the forest can be made compatible with *P. cinnamomi* is described. The approach is based on the idea that promotion of native leguminous species in the forest would create environmental conditions which are unsuitable for activity and survival of the fungus.



Virgin jarrah forest. *Banksia* occurred only as a relatively minor component of the understorey in the virgin forest.

(Les Harman)

Disease spread through *banksia* understorey. This species provides a highly susceptible food base for *P. cinnamomi*. It is unlikely that jarrah dieback will ever be controlled while a dense *banksia* understorey is present. (Les Harman)



The physical environment

P. cinnamomi requires relatively high soil temperature conditions and moist soil before it is able to attack susceptible plants. Prolonged drying of soil can kill the fungus.

In the jarrah forest the length of time when these soil physical conditions exist varies markedly on different sites. In the valleys which are moisture-gaining, soil moisture levels are high during considerable periods of the summer when soil temperatures are suitable for production of infective spores. These sites are highly susceptible to the disease. However, on the freely-drained lateritic and sandy soils which make up approximately 80 per cent of the forest, physical conditions for survival and infection by the fungus are only marginally suitable. In winter, soil temperatures are too low for the fungus to become active, while in summer, the soil is too dry. Optimum soil moisture and temperature levels for production of infective spores by the fungus occur only in spring and autumn.

Extensive trials both in the laboratory and in the field have shown that under a dense canopy and litter layer, soil temperatures are depressed in spring and autumn so that the periods during which there is a co-incidence of high soil moisture and high soil temperature—conditions which favour the fungus—are markedly reduced.

In some forests it is possible to change the canopy of the forest overstorey to create varying degrees of shade. Jarrah crowns, however, are not naturally dense and do not lend themselves to manipulation. However, native legume species can occur as a dense understorey in the jarrah and provide maximum shading.

Host susceptibility

The degree of host susceptibility has a very significant effect on the potential for controlling a plant disease by manipulation of the environment.

Although jarrah trees can be killed by *P. cinnamomi*, this species is not highly susceptible to the fungus. For example, in many disease areas it takes more than ten years after an infection has been initiated before all of the jarrah trees are killed. The fungus cannot invade the large roots of jarrah but causes death by repeated destruction of the fine root system.

Bull banksia (*Banksia grandis*), in contrast to jarrah, is highly susceptible to jarrah dieback and the presence of dense stands of this species in the forest is a major factor contributing to the spread and intensification of the disease. The fungus can move through very large banksia roots. In addition to providing a large food base, banksia roots protect the fungus from

adverse conditions such as low moisture levels, which may occur in the soil outside the roots.

The ability of the fungus to spread through the roots of banksias even when environmental conditions are adverse in the soil outside has a further effect. As the banksias die, more sun reaches the forest floor and soil temperatures are raised. This creates more favourable conditions for spore production, which predisposes less susceptible species to attack.

It is highly unlikely that control of jarrah dieback will be achieved by any method unless the density of the banksia component of the forest is markedly reduced. Such a reduction in banksia density would tend to return the forest to a more “natural” condition because bull banksia occurred only as a minor component of the understorey in the upland virgin jarrah forest.

Extensive tests have been carried out on the relative susceptibility of native legume species to jarrah dieback since it would be pointless to replace banksias with other highly susceptible species. Some of the native legumes are susceptible, but most of those which germinate profusely after fires are resistant. Therefore, replacement of the bull banksia understorey with native legumes would reduce the highly susceptible food base currently present in the forest, which provides a haven for *P. cinnamomi*.



The microbiological environment

Soil plant pathogens, except in sterile laboratory-created conditions, do not operate in a vacuum. The soil contains vast numbers of micro-organisms—some beneficial, some detrimental—which interact with both hosts and the pathogen. *P. cinnamomi*, in particular, is unusually affected by other micro-organisms because in the absence of certain soil bacteria, it will not produce sporangia and zoospores, the

spore types which are believed to be responsible for spread and infection.

It is obvious that on most jarrah forest sites the soil microbiological environment does not present an obstacle to the fungus and it is suspected that on many forest sites it favours *P. cinnamomi*.

If the soil microbiological environment can be changed it is possible that the microbiological conditions can be created which do not favour *P. cinnamomi*.

Preliminary studies suggest that a change from a banksia dominated understorey to one dominated by

▲ *This stand of prickly moses was regenerated by high intensity fire. Previously, the area had a dense banksia understorey and no live legume plants. The red plant is Kennedia coccinea. (Les Harman)*

certain legume species could create a soil microbiological environment which is less favourable for *P. cinnamomi* activity and survival. The composition and quantity of micro-organisms in the soil is markedly influenced by the type of plant growing in the soil. Therefore, it is not surprising that the soil micro-organisms associated with the roots of legume species, which have the ability to add nitrogen to the soil,

are different to those associated with banksia roots. Dr. N. Malajczuk from the C.S.I.R.O. has isolated micro-organisms associated with legume roots and banksia roots and examined their effect on *P. cinnamomi* in controlled laboratory studies. This work has shown that there are more than double the number of antagonistic micro-organisms in soils in which legumes have been grown, compared with soil micro-organisms isolated from soil in which banksias were growing.

More recently research which has been focussed on prickly mosses (*Acacia pulchella*) has shown that chemicals contained in volatiles emanating from roots and chemical extracts of the roots of this species have a highly suppressive effect on *P. cinnamomi*. This discovery parallels research carried out in other parts of the world which suggests that the chemical constituents of certain species is responsible for their resistance to *P. cinnamomi*.

Attempts are being made to isolate the chemical or chemicals responsible for suppression and to determine if they persist in the soil long enough to suppress the fungus.

Some pot trial studies have produced further evidence that legumes will improve resistance to the disease. It has been found when certain legumes are grown in pots which have been infested with *P. cinnamomi* that after a period of time the fungus cannot be recovered from the soil. When jarrah seedlings were grown in conjunction with legume species in pots the mortality of jarrah was significantly reduced when compared to the mortality of pots where jarrah has been grown with bull banksia.

The most significant research finding to this date resulted from a series of field trails which have shown that sporangial production and infection by *P. cinnamomi* was



markedly reduced beneath legume stands in comparison to jarrah forest where legume species were not present.

Improving forest vigour

There is increasing evidence, although it is still circumstantial, that native leguminous species may have additional beneficial effects on the jarrah forest ecosystem: studies carried out by Forests Department and C.S.I.R.O. research workers

▲ *Waterbush* (*Bossiaea aquifolium*) background. This stand of legumes regenerated following the 1961 Dwellingup wildfire.

(Les Harman)

have shown that native legume species have a relatively high capacity to fix atmospheric nitrogen. Some of the species tested have almost as much capacity to fix atmospheric nitrogen as some agricultural species such as clover. It is possible that over a period of years considerable quantities of nitrogen could be incorporated into the soil and eventually be used by other species, such as jarrah, which are unable to obtain their own nitrogen

Top: It is possible by measuring soil moisture and soil temperature regimes, to estimate the number of hours that physical environmental conditions are suitable for infection. When a legume understorey is present, the number of hours when both soil moisture and soil temperature are suitable for infection is markedly reduced.

(Les Harman)

Below: Field tests have shown that infection level is reduced when a legume understorey is present. (Les Harman)



from the atmosphere. This could significantly improve the health and vigour of the forest.

Jarrah does not respond to phosphate fertilisation but in field and pot trial studies significant increases in growth due to nitrogen fertiliser application have been demonstrated. Application of mineral nitrogen fertilisation to legume species inhibits their capacity to fix nitrogen from the atmosphere, however, they respond markedly to phosphate. It is possible that a single application of phosphatic fertiliser to legumes in the forest could be used to increase legume growth which, in turn, would result in higher additions of nitrogen and organic matter to the soil. The effect is likely to be sustained for a number of years because in forest ecosystems nutrients are constantly recycled. There is, therefore, the potential to raise the overall fertility of the jarrah forest ecosystem.


Manipulation of forest environment

It is relatively easy to manipulate the environment in agricultural or horticultural situations where it is practical and economic to employ intensive cultural techniques. For example, avocado growers in Queensland have been able to control *Phytophthora* root rot by large

In pot trial studies it has been found that some legume species, when interplanted with jarrah, increase jarrah's resistance to dieback. (Les Harman)




JARRAH SURVIVAL WHEN INTER-PLANTED WITH



additions of fowl manure and by intensive cultivation of agricultural legume species beneath the trees. But there are severe constraints when the technique must be applied in a susceptible forest extending over more than 1000000 hectares. Currently, there is only one tool available to jarrah forest managers which could be used on a broad scale—fire.

There is abundant evidence that fire is a natural factor in the jarrah forest environment. Apart from the demonstrated ability of the forest to tolerate fire, it is impossible to conceive that even in the absence of man, lightning did not cause ignition of the dry sclerophyll forests during the hot summer months. Although fire was a factor in the forest environment it is only possible to speculate on the frequency, intensity and season of the firing regime in pre-colonial times. In the 1950s and early 1960s broad scale mild prescribed burning, carried out principally in spring, was introduced in the forest. This was necessary to reduce the fuel hazard so as to give a better chance of controlling wild-fires. Perhaps more than in any other forest area in the world, necessity has led to the development of a highly organised and efficient hazard reduction burning system.


Without a hazard reduction burning programme, the forest inevitably would be subjected to severe destructive wildfire such as that which occurred in the forest areas surrounding Dwellingup in 1961. However, the technology of prescribed burning has now advanced to the stage where it is possible to use fire for management objectives in addition to hazard reduction. By using various combinations of fire frequency, intensity and season of burning the forest manager has the potential to change the structure and composition of the forest.




▲ *Sporangial breakdown caused by volatile components of legume roots.*
(Les Harman)




▲ *Viable sporangia releasing zoospores.*
(Les Harman)



▲ *Destruction of P. cinnamomi mycelium caused by micro-organisms associated with legume roots.*
(Les Harman)



▲ *Healthy mycelium.* (Les Harman)



▲ *In recent field tests it has been shown that spore production by P. cinnamomi is markedly reduced when in soil beneath an understorey of legumes, compared with spore production when in soil beneath an understorey of banksias as indicated in the graph.*



A. MYRTIFOLIA
FERTILIZED UNFERTILIZED

A. PULCHELLA
FERTILIZED UNFERTILIZED

A. EXTENSA
FERTILIZED UNFERTILIZED

▲ Stimulation of legume growth by phosphate fertiliser. Application of phosphate to legume stands could raise the overall fertility of the forest ecosystem. (Les Harman)



▲ Nodules found on legume roots. Native legume species can fix atmospheric nitrogen and this may significantly improve the fertility of the forest. (Les Harman)

Relationship between fire and native legumes

One of the most spectacular and rapid changes in the composition and structure of a forest was observed following the Dwellingup wildfire in 1961. Approximately 30000 hectares of forest regenerated with dense stands of native legume species. Following mild hazard reduction burning, these stands failed to regenerate and tended to be replaced by species more susceptible to the disease.

The association between fire and regeneration of leguminous species




▲ Clumped regeneration of acacia species following high intensity fire. Ant species collect acacia seed and store it in their nests. This burial of seed explains why it is necessary to burn the jarrah forest with high to medium intensity fire to stimulate acacia regeneration.

(Les Harman)



Ants collecting legume seed, which they bury in nests soon after it has fallen. This places it below the zone of heat penetration during normal mild fires and so does not germinate. During higher intensity fires heat penetration is sufficient to stimulate germination. (Les Harman) ▲



▼ Legume seedlings regenerating from a range of depths. Original location of seed is shown by the arrow. (Les Harman)

in forest ecosystems is not unusual. Legume seed has a hard coat and it generally will not germinate until it has been heated. For example, laboratory trials have shown that the best way to germinate jarrah forest legumes is to pour boiling water on the seed and allow it to

stand in the water for 15 minutes. In the long leaf pine (*Pinus palustris*) forests of the south-east of the United States of America, frequent mild fire has been used over a number of years to promote the regeneration of a shrubby leguminous layer.



The relationship between legume regeneration and higher intensity fire in the jarrah forest, although not unique, is unusual. Recent research has shown why hotter fires are necessary to regenerate legumes in the jarrah forest whereas in other forest, including the wet sclerophyll karri forests, regeneration of legumes occurs after mild fire.

Following the initial burning trials it was noticed that frequently legumes regenerated in clumps. This suggested that the seed was redistributed following seedfall. Subsequent studies showed that ants are the major agents responsible for legume seed redistribution. In addition to horizontal redistribution, ants also bury the seed. Careful examination of newly regenerated legumes have shown that some originated from as deep as 9 cm below the surface of the soil.

During normal mild spring burns soil temperatures rarely rise above 40°C in the top 1 cm of the soil. Legume seed which is below the surface of the soil does not receive enough heat to stimulate germination. During higher intensity burns, heat penetration is considerable and frequently a large proportion of the seed which has been buried by ants to varying depths in the soil, is stimulated.

Although the research carried out so far has contributed to the understanding of the effect of fire on legume regeneration and survival, the understanding of this relationship is still only partial. However, the results of the study to illustrate the complexity of the ecosystem and that apparently disjunct components such as legumes and ants are, in fact, closely inter-related.

Although mild hazard reduction burning is not the only factor affecting regeneration and survival of bull banksia it does favour this species. Mild burns in spring have little effect on the banksia understorey but they provide an excellent

seedbed for the seeds which are released from cones the following autumn. Thus, under a mild burning regime banksia population levels increase. Current research indicates that medium to high intensity fire in autumn and late summer when banksias are under drought stress, can kill over 50 per cent of the banksia understorey. The research suggests that series of these type of burns over a period of twenty years would reduce banksia numbers to low levels comparable with the virgin forests.


It would obviously not be feasible or desirable to carry out burning at the intensity of a wildfire. Apart from the danger to human life, the Dwellingup wildfire of 1961 caused extensive mortality of jarrah and many of the surviving trees were severely scarred. However, it is possible to carry out fires at intensities above those used commonly for hazard reduction but below the levels causing damage to the overstorey.

Over a period of years Forests Department research workers have developed a set of tables with which, over a range of meteorological and fuel conditions, it is possible to predict the intensity of fire resulting from a particular pattern of ignition. These tables were used to conduct a series of experimental high and medium intensity prescription burns which have shown that it is possible on many forest sites to bring about regeneration of native legumes where previously they were observed to occur only as a minor component of the understorey, and markedly reduce the density of the banksia understorey. These fires, although of higher intensity (and expense) than normal burns, have been controllable and except for jarrah saplings, which in pole stands will not form crop trees, there was no significant damage to jarrah trees. In fact, at least in the short term, growth of jarrah poles has been stimulated by these burns.

Although much more research is required it is conceivable that changes in the structure and composition of the forest (that is from a banksia-dominated understorey to one dominated with legumes) can be achieved on many forest sites by the strategic use of prescribed fire.

Implementation



Intensive research, both in Australia and overseas, is continually improving our knowledge of *P. cinnamomi* and diseases caused by the fungus. This research has shown that plant diseases caused by *P. cinnamomi* are affected by a number of factors and it is unlikely, particularly in a forest situation, that control will be achieved by any single



method. The use of leguminous species to control jarrah dieback is only one of a number of lines of research being investigated. This approach to control of jarrah dieback, however, is attractive because a change to a legume understorey will increase resistance to the fungus in several different ways and it has the potential to be applied on a

broad scale. It is important, however, to emphasise that this approach to control of jarrah dieback is still in the experimental phase. Despite the positive results that have been achieved so far, however, it is essential that the system be thoroughly tested in the field before it is adopted for broad scale application.

It is possible to accurately prescribe the intensity of fire as is illustrated by these photographs of experimental fires. The intensity, frequency and season of burning determine the effect of fire on an ecosystem. (Les Harman)





▲ Forest burnt 12 months previously with fire intensity which resulted in full crown scorch. Note crown vigour. (Les Harman)

▲ Adjacent forest burnt with a normal low intensity burn. Growth rates of scorched trees have been shown to be significantly higher than trees which have been burnt with low intensity fire. (Les Harman)


Although there is little doubt that a legume understorey will reduce the susceptibility of the forest to jarrah dieback, it is not known if the reduction in susceptibility will have practical significance. Prolongation of the survival of a horticultural crop such as avocado, which is being attacked by *P. cinnamomi* for ten years is a significant advance. However, in a forest crop such as jarrah the effect of a cultural technique must be permanent. Thus, it is necessary for careful long-term field trials to be carried out before the legume approach to control can be implemented on an operational scale.

It is also important to ensure that high intensity fire does not have serious adverse effects on the forest. For example, what will be the effect on native animals and wildflowers? What will be the effect of the

changed burning regime on the bee-keeping industry? Although the preliminary studies indicate that forest areas which have been subjected to high intensity fire are more vigorous than those which are protected or burnt mildly, it is not certain that this vigour can be maintained. It is also not known if there is a store of legume seed in the soils on all jarrah sites—it may be necessary to apply seed directly on some sites. High intensity prescription burning is also more hazardous and as a consequence more expensive than mild burning in spring. Will the costs be prohibitive? All these questions must be resolved before changes in existing management practices are made.

Even though this approach to control of jarrah dieback is being treated with caution, a number of large scale trials have been initiated

to determine the practical problem of high intensity burning. The decision to move to large scale trials, even though fundamental research has not been completed, has been made deliberately. This approach to control of jarrah dieback is based on the concept of forest manipulation—it cannot be implemented if there is no forest to manipulate. Once the disease extends to the overstorey mortality stage it would be difficult to reverse the decline in forest health.

The large scale trials being carried out will ensure that if it is found a legume understorey *will* confer permanent resistance of the forest to jarrah dieback, the technique can be implemented rapidly. The maintenance of stringent hygiene and in some forest areas, quarantine, will provide the maximum length of time to develop and test jarrah dieback control techniques. 

Re-establishment of jarrah on bauxite pits



▲ A four-year-old stand of myrtle wattle which was established on a bauxite pit by direct seeding. (Les Harman)

Forest ecological research is complex and requires a multi-disciplinary approach. A number of research workers from a variety of organisations have been assisting full or part-time on this particularly jarrah dieback research programme.

Soil microbiology. Dr. N. Malajczuk and Mr. D. Darling, C.S.I.R.O., Dr. A. Glen, Murdoch University, and Miss W. Lepard, Forests Department.

Field experiments. Messrs. R. J. Kitt, R. G. Fairman and M. L. Mason, Forests Department.

Laboratory experiments. Mr. K. J. Gillen and Miss H. J. Warren, Forests Department, and Mr. K. Russell, C.S.I.R.O.

Analysis of volatile chemicals from legumes. Drs. F. Whitfield and K. J. Shaw, C.S.I.R.O.

Plant chemistry. Mr. A. B. Hatch, Forests Department; Drs. R. Kagi, B. Alexander, K. Croft and R. Bradbury, Western Australian Institute of Technology.

Fire ecology. Messrs. J. McCormick, R. M. Buehrig and C. C. Portlock, Forests Department.

Ant ecology. Dr. J. D. Majer, Western Australian Institute of Technology.

Fire management. Mr. L. Robson, Forests Department.

Jarrah has not been used on any scale to rehabilitate bauxite mine sites since, during mining, it is probable that *P. cinnamomi* spores, if they were not already present in the soil, would be introduced during the mining process. A range of alternative tree species which are resistant to *P. cinnamomi* has been established on bauxite mine sites. While it is possible that these species may replace jarrah, it may be desirable to re-establish jarrah on bauxite mine sites, particularly if mining extends to salt-prone areas of the forest.

It is likely to be more difficult to achieve control of *P. cinnamomi* by use of an understorey of native legumes in comparison to unmined forest. For example, drainage is impeded on bauxite mine sites and this is likely to increase the susceptibility of trees to the disease. It is possible, however, that native leguminous species if interplanted with jarrah could confer on it resistance to the disease and thus permit re-establishment of jarrah on bauxite mined sites.

Part of the mining procedure involves removal and stockpiling of the topsoil. Much of the seed and lignotuberous material is killed during prolonged stockpiling so that the understorey and shrub layer is not re-established naturally on bauxite mine sites. Research by W. Tacey of Alcoa has shown that if the topsoil is

immediately replaced a considerable proportion of the seed stored in the soil remains viable. There are logistical problems associated with avoiding storage of topsoil, and even if these can be overcome it is probably still desirable to enrich the shrub and understorey layers on bauxite pits.

A series of trials was initiated by Forests Department research officers to determine if it was possible to re-establish a native understorey and shrub layer on bauxite pits to improve stability of the pit surface and increase the diversity of plant species on rehabilitated mine sites. Native legume species were primarily used because of the availability of seed, and their resistance to *P. cinnamomi*, and their ability to improve soil fertility by nitrogen fixation. The trials were highly successful and were expanded to include other native shrubs and a variety of tree species. Aerial seeding of shrub and understorey seed on bauxite pits has now been adopted on an operational basis.

Tests are currently being conducted, using some of the early legume trials on bauxite pits, to determine if the reduction in fungal activity, which has been demonstrated under native legume stands in the forest, can be reproduced on bauxite pits.

In trial areas jarrah seed has been incorporated into the mix of legume and shrub species which is applied to mined-over areas after they have been replanted with dieback-resistant trees. Establishment of jarrah as part of the shrub and understorey layer beneath the fast-growing planted species may provide the opportunity in the future, if control of jarrah dieback by legumes is achieved, to re-establish jarrah as the dominant species on bauxite pits.

Double Safety Award to Western Australian Forests Department

In November 1978, the Forests Department of Western Australia won Australia's top industrial safety award—the C.M.L. Award for Industrial Safety—in competition with state and national companies and other government departments.

This followed on the heels of equal success for the department in the state section of the C.M.L. Award in the previous September.

The C.M.L. Award trophies, together with the national award certificate.

(Brian Stevenson)



Forestscapes



It's an ill wind . . .

The old saying "It's an ill wind that blows no good", seems to apply even to Cyclone Alby which battered the state last April. The wildfires which were caused by the cyclone, apart from causing havoc in some areas, also led to the profuse regeneration of wildflowers.

◀ *This area of burnt jarrah forest is carpeted with blue lechenaultia.*

Custard orchid. ▶

▼ *Donkey orchids are scattered among the lechenaultia.* (All Les Harman)







INDEX

This index lists notable photographs and articles which have been published in *Forest Focus* Nos. 1 to 21.

Italic figures indicate photographs * indicates same photograph

- Acacia
Focus on an ecological approach to the control of jarrah dieback, by S. R. Shea **21**: 1-19
seed coat **13**: 6
- Acacia*
myrtifolia (myrtle wattle) **21**: 28
pentadenia (karri wattle) **18**: 4
podalyriifolia (Queensland silver wattle) **15**: 15*, **17**: 7*
pulchella (prickly moses) **21**: 10
urophylla **18**: 10
- Acanthiza chrysorrhoa* (yellow-tailed thornbill) **7**: 4
- Aerial incendiary operation **3**: 4
- Agonis juniperina* (Warren River cedar) **18**: 2
- Agro-forestry **16**: 13-14
trials in the south-west, Focus on, by F. H. McKinnell and F. Batini **20**: 1-16
- Agricultural clearing, Murray catchment **19**: 10
- Aleppo pine, see *Pinus halepensis*
- Amphibolurus ornatus* (ornate dragon, running lizard) **8**: 12
- Anigozanthos flavidus* (tall kangaroo paw) **10**: 15
- Anniversary of a landmark (One-tree Bridge) **13**: 10-11
- Ant-eater
banded, see *Myrmecobius fasciatus*
spiny, see *Tachyglossus aculeatus*
- Antechinus flavipes* (mardo, yellow-footed marsupial mouse) **10**: 7
- Anthochaera carunculata* (red wattle-bird) **10**: 15
- Ants collecting legume seeds **21**: 15
- Arboretum, Badgingarra **4**: 14
- Asterolasia pallida* (star flower) **17**: 2
- Bandicoot, short nosed, brown, see *Isodon obesulus*
- Banksia **21**: 8-9
root **14**: 4
woodlands **14**: 8
- Banksia*
attenuata **10**: 13, **18**: 14
fruits **13**: 2
dead **10**: 12
coccinea (scarlet banksia) **10**: 14
-
- Bushy kennedia*, also regenerated by hot fire.
(Les Harman)
- grandis* (bull banksia) **10**: 13
dead **10**: 12, **14**: 5
ilicifolia (holly-leaved) **18**: 15
littoralis (swamp) **10**: 14
- Barrabup Pool **18**: 6
- Batini, F. E.
Jarrah root rot **4**: 10-11
Trees for country areas **4**: 12, 14
- Batini, F. E., and Havel, J. J. Focus on land use conflicts in the northern jarrah forest **11**: 1-16
- Batini, F. E., and McKinnell, F. H. Focus on agro-forestry trials in the south-west **20**: 1-16
- Bauxite mining
Focus on loss of productive forest **4**: 1-9, 12, 16
Focus on managing jarrah forest catchments **19**: 1-20
Re-establishment of jarrah on bauxite pits **21**: 19, 28
Rehabilitation after, root development **19**: 14
seedlings **11**: 10
Spread of dieback **14**: 14
- Beaufortia sparsa* (swamp bottlebrush) **2**: 9, **13**: 5 (bottom, second from left), **18**: 5
- Bettongia penicillata* (woylie, brush-tailed rat-kangaroo) **10**: 5, **12**: 12
- Bibbulmun Bushwalking Track, by R. Gobby **13**: 12-15
- Big Brook Arboretum picnic seats **2**: 10
- Bird numbers in forest **12**: 10 (table)
- Birds of the jarrah forest, by P. Kimber **7**: 1, 4-7
- Blackboy, see *Xanthorrhoea preisii*
- Blackbutt
Coastal, see *Eucalyptus todtiana*
see *E. patens*
- Blackwood River **16**: 7
- Blackwood Valley, summary of main plantation operations **5**: 9
- Bluebush, see *Hovea elliptica*
- Boongul, see *Eucalyptus transcontinentalis*
- Boranup **18**: 7, 24
- Boronia resource survey **3**: 15
- Boronia megastigma* (brown or scented) **3**: 15, **13**: 5
- Bossiaea*
aquifolium (waterbush) **21**: 11
laidlawiana (netic) **18**: 10
ornata **17**: 2
- Bottlebrush
Albany, see *Callistemon speciosus*
Swamp, see *Beaufortia sparsa*
- Boulder Rock **5**: 13, **9**: 8, **15**: 16
- Bullich, see *Eucalyptus megacarpa*
- Bunnings Welshpool building **13**: 9
- Butterbark, see *Eucalyptus laevis*
- Bushfire survival, Focus on **6**: 9-12, 14-15
- Bushy kennedia, see *Kennedia stirlingii*
- Caladenia filamentosa* (red spider orchid) **14**: 2
- Caladenia*
flava (cowslip orchid) **13**: 8
sericea (silky blue orchid) **13**: 5
- Callcup dune **8**: 1-6, 10-11, 14-16
- Callistemon speciosus* (Albany bottlebrush) **18**: 5
- Canary Island pine, see *Pinus canariensis*
- Canoeing **11**: 8
- Cascades, Pemberton **2**: 7, **18**: 15 (captioned as Lane Poole Falls)
- Casuarina (sheoak) **18**: 18
- Casuarina huegliana* (granite) **11**: 12
- Cedar, native, see *Agonis juniperina*
- Cercartetus concinnus* (pigmy possum, munda, south-western pigmy possum) **10**: 11, **14**: 12
- Christensen, P. E. S.
Focus on a new concept in forestry—fauna priority areas **10**: 1-10
Focus on the role of fire in the south-west forest ecosystems **13**: 1-8, 16
Some ecological aspects of jarrah dieback **10**: 11-16
- Christmas tree, see *Nuytsia floribunda*
- Christmas tree well **7**: 15-16
- Chuditch, see *Dasyurus geoffroyi*
- Clematis pubescens* (Traveller's joy) **2**: 9, **13**: 5 (right, middle, bottom)
- C.M.L. Safety Award **21**: 20
- Cone, needle and bud of main pine species grown in Western Australia **5**: 10-11
- Conservation, Focus on forest **2**: 1-6
- Contour plantation road, Murray Valley **2**: 14*, **5**: 5*
- Controlled burning for forest conservation, Focus on **3**: 1-8
- Coral vine, see *Kennedia coccinea*

Crowea
angustifolia 13: 5 (left, bottom)
dentata (bush crowea) 2: 9
 Crowned snake, see *Denisonia coronata*

Daisy, blue, see *Olearia rudis*

Darling Range
 Focus on land use conflicts in the northern jarrah forest, by J. J. Havel and F. E. Batini 11: 1-16
 Recreation 5: 14

Dasyurus geoffroyi (Western native cat, chuditch) 10: 8

Deadman's trail footbridge, Manjimup 9: 15-16

Demansia nuchalis affinis (dugite) 10: 9

Denisonia coronata (crowned snake) 10: 16

Diamond tree fire lookout 3: 1, 3

Dieback hygiene . . . first steps, Focus on 21: 1-6

Dieback, see Jarrah dieback

Diuris setacea (bristly donkey orchid) 19: 19, 21: 20

Dodonaea attenuata (native hop) 17: 2

Donnelly River 18: 16, 17
 Deadman's trail footbridge 9: 15-16
 One-tree bridge 13: 10-11

Donnybrook sunklands, Focus on 16: 1-16

Dromaius novaehollandiae (emu) 10: 6

Dryandra an ecological oasis, Focus on 8: 7-9, 12-13

Dryandra
formosa (showy) 10: 14
praemorsa (urchin) 17: 2
sessilis (parrot bush) 10: 15

Dual-purpose fire control vehicle 1: 11

Dugite, see *Demansia nuchalis affinis*

Dundas mahogany, see *E. brockwayi*

Dunnart, see *Sminthopsis murina*

Dwellingup fire, 1961 6: 15

Early history of Jarrahdale, by E. Willis 7: 2-3, 10-14

Echidna, see *Tachyglossus aculeatus*

Ecological approach to the control of jarrah dieback, Focus on, by S. R. Shea 21: 7-19

Elythranthera brunonsis (purple enamel orchid) 13: 8

Emu, see *Dromaius novaehollandiae*

Environment, Trees are regulators of the 9: 15

Environmental costs of building materials 13: 9

Eopsaltria griseogularis (Western yellow robin) 8: 13*, 13: 4*

Errata 6: 3, 12: 15, 17: 14

Eucalypts
 a simplified key to 17 W.A. species 2: 11-13
 inland 15: 1-11

Eucalyptus
accedens (powderbark wandoo) 2: 2*, 8: 12*, 11: 9
 buds and fruits 2: 13

astringens (brown mallet) 8: 12-13
 buds and fruits 2: 13

brevistylis (Rates tingle) 18: 9

calophylla (marri)
 buds and fruits 2: 13
 Focus on the marri woodchip project 12: 1-16
 in dieback areas 14: 6
 and pasture 20: 15

cornuta (yate) buds and fruits 1: 13

diversicolor (karri)
 buds and fruits 2: 13
 effect of wildfire on karri crowns 3: 7
 Focus on the karri forest 1: 1-7, 13, 16
 Focus on the marri woodchip project 12: 1-16
 Focus on the role of fire in the south-west forest ecosystems 13: 1-8, 16
 Focus on southern recreation and conservation management priority areas 18: 1-24
 Forestscapes 6: 15-16
 Four Aces, Manjimup 9: 6
 Lefroy Brook, 100 years old 1: 4, 2: 10
 life cycle 17: 8-10
 masterpiece of engineering 4: 15
 ringbarked 12: 9, 20: 2
 seed trees 12: 16
 six-month-old regeneration 12: 5
 in Sunkland 18: 18

eremophila (tall/goldfields sand mallee) 6: 13*, 15: 8*

erythrocorys (illyarrie) 4: 13

erythronema (red-flowered mallee, white mallee) 15: 9

ficifolia (red flowering gum) 18: 2

flocktoniae (merrit) 15: 6

forrestiana (fuchsia gum) 1: 15

globulus (Tasmanian blue gum) 14: 13

gompophcephala (tuart) 18: 17, 20: 15
 buds and fruits 2: 13

guilfoylei (yellow tingle) 18: 9

haematoxylon (mountain marri) 16: 9

jacksonii (red tingle) buds and fruits 2: 13

laeliae (Darling Range ghost gum, butterbark) 11: 5

loxophleba (York gum) buds and fruits 2: 13

macrocarpa (mottlecah) 15: 2

maculata (spotted gum) 19: 13, 15

marginata (jarrah)
 buds and fruits 2: 13, 6: 3
 fifty-three years old 4: 16, 11: 10
 Focus on birds of the jarrah forest 7: 1, 4-7
 Focus on dieback hygiene and an ecological approach to the control of jarrah dieback 21: 1-19
 Focus on jarrah dieback—a threat to W.A.'s unique jarrah forest 14: 1-16
 Focus on jarrah forest 6: 1-8
 Focus on land use conflicts in the northern jarrah forest 11: 1-16
 Focus on loss of productive forest 4: 1-9, 12, 16
 Focus on managing jarrah forest catchments 19: 1-20

Focus on Sunklands multiple use land management 16: 1-16

Re-establishment of jarrah on bauxite pits 21: 19, 28

Southern recreation and conservation management priority areas 18: 1-24

forest devastated by wildfire 3: 5

high quality 16: 5

jarrah/marri regeneration 12: 8

jarrah root rot 4: 10-11

natural occurrence—map 7: 7*, 6: 2*

northern jarrah forest catchments—map 19: 4

open forest 10: 4

pole stand control burnt for 40 years 3: 2

regrowth after 1961 wildfire 3: 8

roots 14: 4, 19: 5

timber haulage 17: 13

vegetation types 19: 2

megacarpa (bullich) 11: 7, 18: 16, 18: 19: 2
 buds and fruits 2: 13

microcorys (tallowwood) 4: 4, 9, 19: 15

occidentalis (flat-topped yate) buds and fruits 2: 13

patens (W.A. blackbutt) 11: 8, 18: 17
 buds and fruits 2: 13
 "pterocarpa" 15: 2*, 17: 5*

rhodantha (rose mallee) 1: 15

rudis (flooded gum) 19: 2
 buds and fruits 2: 13

salmonophloia (salmon gum) 15: 10
 buds and fruits 2: 13

salubris (gimlet) 15: 10
 buds and fruits 2: 13

tetraptera (four-winged mallee) 2: 15

toytiana (pricklybark, coastal blackbutt) buds and fruits 2: 13

torquata (coral gum) 15: 6-7*, 11: 17: 2*

torquata and *E. woodwardii* hybrid (torwood) 15: 11

transcontinentalis (redwood, boongul) 15: 10

wandoo (wandoo) 3: 9-10, 16, 5: 13*, 10: 4, 11: 11*, 12: 13, 19: 2
 buds and fruits 2: 13

woodwardii (lemon-flowered gum) 15: 6

woodwardii and *E. torquata* hybrid (torwood) 15: 11

youngiana (large-fruited mallee) 15: 1

Facts and figures, 1970, forest area 3: 11, 14

Fauna priority areas, Focus on a new concept in forestry, by P. Christensen 10: 1-10

Featherflower, painted, see *Verticordia picta*

Fire control
 an early problem 3: 5-6
 vehicle 1: 11

Fire
 Forests, flora, fauna 3: 6-8
 in the south-west forest ecosystems, Focus on the role of, by P. Christensen 13: 1-8, 16
 lookout tower, Diamond tree 3: 1-2
 wildfire 17: 6

Firetail, red-eared, see *Zonaeginthus ocellatus*

- Firewood, goldfields **15: 5**
- First steps in W.A. paper manufacturing industry **1: 8-9**
- Flight Line One, film on prescribed burning in W.A. forests **7: 8-9**
- Flowering eucalypts **1: 14**
- Focus on agro-forestry trials in the South-West **20: 1-16**
- birds of the jarrah forest **7: 1, 4-7**
- bushfire survival **6: 9-12, 14-15**
- controlled burning for forest conservation **3: 1-8**
- country recreation grant **9: 6-9**
- dieback hygiene and an ecological approach to the control of jarrah dieback **21: 1-19**
- Dryandra: an ecological oasis **8: 7-9, 12-13**
- early history of Jarrahdale **7: 2-3, 10-14**
- forest conservation **2: 1-6**
- forest policy **17: 1-16**
- forest recreation **5: 12-16**
- inland eucalypts **15: 1-11**
- jarrah dieback, a threat to W.A.'s unique jarrah forests **14: 1-14, 16**
- jarrah forest **6: 1-8**
- karri forest **1: 1-7, 13, 16**
- land use conflicts in the northern jarrah forest **11: 1-16**
- loss of productive forest **4: 1-9, 12, 16**
- managing jarrah forest catchments **19: 1-20**
- marri woodchip project **12: 1-16**
- natural rounds here to stay **9: 1-5**
- a new concept in forestry—fauna priority areas **10: 1-10**
- role of fire in the South-West forest ecosystems **13: 1-8, 16**
- shifting sands **8: 1-6, 10-11, 14-16**
- southern recreation and conservation management priority areas **18: 1-24**
- Sunklands multiple use land management **16: 1-16**
- thirty-million dollar integrated forest products complex **5: 1-8**
- Forest areas of south-west W.A. **4: 2** (map)
- comparison with other countries **4: 2** (table)
- Forest communications **1: 12-13**
- Forest facts and figures **3: 11, 14**
- Forest policy, Focus on **17: 1-16**
- Forest recreation, by D. Spriggins **5: 12-16**
- Forests Minister approves grant for country works (recreation sites) **9: 6-9**
- Forestry in Western Australia, new edition **9: 14**
- Forestscapes **21: 20-22**
- Christmas tree well **7: 15-16**
- Donnelly log footbridge **9: 15-16**
- jarrah **4: 15-16**
- karri **6: 15-16**
- Four aces, Manjimup **9: 6**
- Four-wheel drive vehicle causing erosion **15: 13**
- Frankland River **18: 15**
- Gastrolobium bilobum* (heart-leaved poison) **10: 4**
- Gimlet, see *E. salubris*
- Gleneagle **5: 13, 16, 17: 7**
- Glossopsitta porphyrocephala* (purple-crowned lorikeet) **7: 4**
- Gobby, R., Bibbulmun bushwalking track **13: 12-15**
- Goldfields sand mallee, see *E. eremophila*
- Grevillea* **18: 5**
- endlicherana* (spindly) **17: 2**
- pilulifera* (woolly-flowered) **17: 2**
- wilsonii* (wilson's) **8: 13**
- Group settlement scheme, ring-barked karri **12: 9, 20: 2**
- Growing importance of pine **1: 9-10**
- Gum
- coral, see *E. torquata*
- Darling Range ghost, see *E. laeliae*
- flooded, see *E. rudis*
- fuchsia, see *E. forrestiana*
- lemon-flowered, see *E. woodwardii*
- red-flowering, see *E. ficifolia*
- salmon, see *E. salmonophloia*
- York, see *E. loxophleba*
- Hakea*
- laurina* (pin-cushion) **2: 15**
- lissocarpha* (honey-bush) **17: 2**
- Hamel nursery **2: 16, 4: 14, 5: 5**
- Hardenbergia comptoniana* (wild sarsaparilla, native wistaria) **2: 9**
- Hauling—past and present **3: 12-13**
- Havel, J. J., and Batini, F. E. Focus on land use conflicts in the northern jarrah forest **11: 1-16**
- Heart-leaved poison, see *Gastrolobium bilobum*
- Herbert, E., and Shea, S. Focus on managing jarrah forest catchments **19: 1-20**
- Hewett, P. N. Recreation characteristics of Western Australian forests **15: 12-16**
- Hibbertia cuneiformis* (cut leaf) **13: 5** (right top)
- Holyoake, 1961 **6: 14**
- Honeybush, see *Hakea lissocarpha*
- Honeyeater, new-holland, see *Phylidonyris novaehollandiae*
- Hop, native, see *Dodonaea attenuata*
- Hovea*
- chorizemifolia* (holly-leaved) **17: 2**
- elliptica* (tree hovea, bluebush) **13: 5** (right, bottom), **18: 11**
- Hydrological cycle **19: 5** (diagram)
- Hydromys chrysogaster* (water rat, beaver rat) **10: 10**
- Illyarrie, see *Eucalyptus erythrocorys*
- Information sheet series **12: 15**
- Inland eucalypts—a valuable biological resource, Focus on **15: 1-11**
- Iris morning, see *Orthrosanthus laxus*
- Isoodon obesulus* (quenda, short-nosed bandicoot, brown bandicoot) **2: 2*, 8: 12*, 10: 10**
- Isopogon dubius* **14: 2** (Incorrect should be *Kunzea micrantha*.)
- Jarra, see *E. marginata*
- Jarra dieback
- “A” class reserve near Pinjarra—conservation value destroyed by *Phytophthora cinnamomi* **14: 11**
- effect of **11: 8**
- Focus on dieback hygiene and an ecological approach to the control of jarrah dieback **21: 1-19**
- Focus on jarrah dieback a threat to W.A.'s unique jarrah forest **14: 1-14, 16**
- Focus on managing jarrah forest catchments **19: 1-20**
- in regrowth forest **17: 4**
- jarrah root rot **4: 10-11**
- rehabilitation **16: 11, 12**
- resistant species **4: 11, 14: 15**
- some ecological effects of **10: 11-16**
- in Sunklands **16: 4**
- washdown station **17: 6**
- Jarrahdale early history, by E. Willis **7: 2-3, 10-14**
- Kangaroo, western grey, see *Macropus fuliginosus*
- Kangaroo paw, tall, see *Anigozanthos flavidus*
- Karri, see *E. diversicolor*
- Kayaking **5: 14, 11: 8**
- Kennedia*
- coccinea* (coral vine) **17: 2, 21: 10**
- stirlingii* (bushy kennedia) **21: 22**
- Kimber, P. Birds of the jarrah forest **7: 1, 4-7**
- Kingia/jarra scrubby flats **16: 8**
- Kunzea micrantha* **14: 2**. (Incorrectly called *Isopogon dubius*.)
- Lane-Poole falls **18: 15**. (Incorrect, should be Cascades.)
- Land use conflicts in the northern jarrah forest, Focus on, by J. J. Havel and F. E. Batini **11: 1-16**
- Lechenaultia*
- biloba* (blue) **8: 13, 17: 2, 21: 20**
- formosa* (red) **8: 13**
- Lefroy Brook karri regeneration **1: 4, 2: 10**
- Legumes native, see *Acacia*
- Lesley picnic ground **9: 7**
- Lizard, running, see *Amphibolurus ornatus*
- Locomotives **7: 11**
- Logging operations **11: 9**
- Lorikeet, purple-crowned, see *Glossopsitta porphyrocephala*

- McKinnell, F. H., and Batini, F. Focus on agro-forestry trials in the South-West 20: 1-16
- Macropus*
eugenii (tammar, scrub wallaby) 10: 6, 13: 7
 habitat 13: 8
fuliginosus (western grey kangaroo) 10: 6
irma (western brush wallaby) 10: 5
- Macrozamia riedlei* (zamia palm) 14: 5
- Mallee
 four-winged, see *Eucalyptus tetraptera*
 large-fruited, see *E. youngiana*
 red-flowered/white, see *E. erythronema*
 rose, see *E. rhodantha*
 tall/goldfields sand, see *E. eremophila*
- Mallet, brown, see *E. astringens*
- Malurus*
elegans (red-winged wren) 2: 2*, 7: 4*, 8: 12*, 13: 5
splendens (splended, blue-banded) 7: 4, 13: 5*, 14: 2*
- Management priority areas 18: 1-24
 Sunklands 16: 1-16
- Managing jarrah forest catchments, Focus on, by S. Shea and E. Herbert 19: 1-20
- Maps
 Area of Bibbulmun tribe 13: 15
 Bibbulmun track 13: 13
 Dryandra 8: 8-9
 Forest areas of south-west W.A. 4: 2
 Jarrah—natural occurrence 6: 2*, 7: 7*
 Jarrah dieback distribution 14: 10
 Karri distribution 1: 6
 Management priority areas 18: 12-13
 Mining leases 4: 2, 11: 2
 Northern jarrah forest catchments 19: 4
 Pemberton area tourist map 2: 8-9
 Perup fauna priority area 10: 2
 Picnic sites, State Forest near Perth 5: 15
 South coast sand dunes 8: 2
 Sunklands locality plan 16: 2
 Wandoo distribution 3: 10
 Water/catchments 4: 2, 11: 2, 19: 5
 Woodchip area 12: 4
- Mardo, see *Antechinus flavipes*
- Margaret River 16: 8
 Rapids Pool 18: 6
- Maritime pine, see *Pinus pinaster*
- Marram grass 8: 5, 11, 14
- Marri
 see *Eucalyptus calophylla*
 mountain, see *E. haematoxylon*
- Masterpiece of engineering 4: 15
- Melaleuca* (paperbark) 18: 2
preissiana 19: 2
radula (graceful honey-myrtle) 17: 2
- Millionth acre of plantations 1: 14
- Mining, see bauxite
- Milyeannup
 high quality jarrah 16: 5
 Brook 16: 6
- Minister for Forests
 T. D. Evans 4: 3
 H. D. Evans 6: 2
 A. Ridge 17: 3
 M. J. Criag 18: 3
 D. Wordsworth 21: 3
- Mirbelia spinosa* 17: 2
- Monterey pine, see *Pinus radiata*
- Morelia variegata* (carpet snake) 10: 9
- Mottlecrah, see *Eucalyptus macrocarpa*
- Mount
 Cooke, Cuthbert, Randall, Vincent 11: 6
 Frankland 18: 1, 8, 16
- Mouse
 marsupial, see *Pseudomys* sp.
 common, see *Sminthopsis murina*
 yellow-footed, see *Antechinus flavipes*
- Multiple use 11: 1, 4; 16: 1-16; 18: 1-24
- Mundarda, see *Cercatetus concinnus*
- Murray
 River 11: 8
 Valley 15: 16, 17: 7
- Mymecobius fasciatus* (numbat, banded ant-eater) 2: 2*, 8: 13*, 11: 11*, 17: 1, 19: 19
- Myrtle, graceful honey, see *Melaleuca radula*
- Native cat, Western, see *Dasyurus geoffroii*
- Natural rounds—here to stay 9: 1-5
- Netic, see *Bossiaea laidlawiana*
- New concept in forestry—fauna priority areas 10: 1-10
- Nicholls, Lexie 4: 9
- Noolbenger, see *Tarsipes spencerae*
- North Dandalup River 11: 14
- Notechis scutatus occidentalis* (Western tiger snake) 10: 9
- Now for a few good words about wood . . . 13: 9
- Numbat, see *Myrmecobius fasciatus*
- Nuytsia floribunda* (Christmas tree) 7: 15-16, 9: 9, 18: 2
- Oil pollution 4: 9
- Olearia rudis* (rough daisy bush; blue daisy) 18: 5
- One million acres of softwoods 2: 14, 16
- One Tree Bridge 9: 9, 13: 10-11
- Orchards 11: 7
- Orchid
 bristly donkey, see *Diuris setacea*
 cowslip, see *Caladenia flava*
 custard, see *Thelymitra villosa*
 purple enamel, see *Elythranthera brunonis*
 red spider, see *Caladenia filamentosa*
 silky blue, see *Caladenia sericea*
- Ornate dragon, see *Amphibolurus ornatus*
- Orthranthus latus* (morning iris) 13: 5 (left, top)
- Over grazing 2: 6
- Paper manufacturing, first steps in W.A. 1: 8-9
- Paperbark, see *Melaleuca*
- Parrot, red-capped, see *Purpurecephalus spurius*
- Peet, G. Prescribed burning in W.A. forests 7: 9, 14-15
- Pemberton area tourist map 2: 8-9
- Perup
 fauna priority area 10: 2
 river 10: 7
- Petroica multicolor* (scarlet robin) 13: 4
- Phylidonyris novaehollandiae* (new holland honeyeater) 7: 4, 10: 15*, 14: 12*
- Phytophthora cinnamomi*, see jarrah dieback
- Pimelea rosea* (rose banjine) 2: 9
- Pinaster pine, see *Pinus pinaster*
- Pine
 the growing importance of 1: 9-10
 Harvey Weir 11: 9
 plantations, wildlife in 2: 6
 seedlings survival 9: 14
 species—cone, needle and bud 5: 10-11
- Pines and agriculture 16: 13-14, 20: 1-16
- Pinus*
canariensis (Canary Island) 5: 11
elliottii (slash) 5: 11
halepensis (Aleppo) 5: 10
pinaster (pinaster, maritime) 5: 1-10
 on Callcup dune 8: 5
 on dieback site 4: 11
pinea (stone, umbrella) 5: 11
radiata (radiata, Monterey) 5: 1-10, 16: 1-16, 17: 12
 Grimwade plantation 1: 9
 killed by *Phytophthora cinnamomi* 14: 8
- Plantations, millionth acre of 1: 14
- Platycercus icterotis* (Western rosella) 7: 4
- Playground equipment—pine 9: 2, 4-5
- Porcupine, see *Tachyglossus aculeatus*
- Possum
 honey, see *Tarsipes spencerae*
 pigmy, see *Cercatetus concinnus*
 ringtailed, see *Pseudocheirus peregrinus*
- Prescribed burning 3: 1-8
- Preservative treated pine poles 9: 1-5
- Pricklybark, see *E. todtiana*
- Prickly Moses, see *Acacia pulchella*
- Pseudocheirus peregrinus* (ringtailed possum) 10: 1, 9
- Pseudomys* sp. (marsupial mouse) 14: 2
- Purpurecephalus spurius* (king parrot) 7: 1
- Push-button age 9: 10-12
- Quarantine, Focus on dieback hygiene and an ecological approach to the control of jarrah dieback 21: 1-19
- Quarry 11: 5
- Queensland party sees W.A. forests, flowers, farms 2: 15
- Quenda, see *Isodon obesulus*
- Quokka, see *Setonix brachyurus*
- Radiata pine, see *Pinus radiata*
- Radio
 forest communications 1: 12-13
 tracking—fauna surveys 12: 12

- Rainbow trail leaflet **2: 7-10**
- Rat
beaver, see *Hydromys chrysogaster*
southern bush, see *Rattus fuscipes*
water, see *Hydromys chrysogaster*
- Rat-kangaroo, brush tailed, see *Bettongia penicillata*
- Rattus fuscipes* (southern bush rat) **10: 10, 13: 7**
- Recreation characteristics of Western Australian forests, by P. N. Hewett **15: 12-16**
- Recreation, forest **5: 12-16**
Forests Minister approves grant for country works **9: 6-9, 15-16**
- Redwood, see *E. transcontinentalis*
- Re-establishment of jarrah on bauxite pits **21: 19, 28**
- Repeater station **1: 12**
- Ringbarking
karri **12: 9, 20: 2**
Mundaring Weir **2: 3**
- Roadside flowering trees **2: 15**
- Robin
scarlet, see *Petroica multicolor*
western yellow, see *Eopsaltria griseogularis*
- Rock climbing **5: 12**
- Rockingham jetties **7: 2**
- Rose banjine, see *Pimelea rosea*
- Rosella, western, see *Platycercus icterotis* **7: 4**
- Safety Award **21: 20**
- Salt affected land **11: 10*, 14: 10*, 19: 20, 20: 2**
- Sand dunes, Focus on shifting sands **8: 1-6, 10-11, 14-16**
- Sarsaparilla wild, see *Hardenbergia comp-toniana*
- Sawmills—The push-button age **9: 10-12**
- Saws **9: 10-12, 17: 14**
- Selected flowering eucalypts of Western Australia **4: 13, 6: 12-13, 9: 13-14**
- Serpentine Dam **4: 6**
- Setonix brachyurus* (quokka) **10: 8*, 11: 6***
- Shannon Dam **18: 6**
- Shea, S. R.
Focus on an ecological approach to the control of jarrah dieback **21: 7-19**
Focus on jarrah dieback, a threat to W.A.'s unique jarrah forests **14: 1-14, 16**
- Shea, S. R., and Herbert, E. Focus on managing jarrah forest catchments **19: 1-20**
- Shelterbelts **20: 4**
- Sheoak, see *Casuarina*
- Shifting sands, Focus on, by B. J. White **8: 1-6, 10-11, 14-16**
- Signs for recreation sites **5: 12, 14**
- Slash pine, see *Pinus elliotii*
- Sleepers, track laying **6: 7**
- Sminthopsis murina* (dunnart, common marsupial mouse) **10: 7**
- Smoky Bear's story of the forest, new publication **1: 14**
- Snake
carpet, see *Morelia variegata*
crowned, see *Denisonia coronata*
Western tiger, see *Notechis scutatus occidentalis*
- Snigging **6: 6**
- Softwood plantations **5: 1-11, 16: 6-15**
- Softwoods, one million acres of **2: 14, 16**
- Some ecological aspects of jarrah dieback, Focus on, by P. Christensen **10: 11-16**
- South Dandalup catchment salinity **19: 16** (diagram)
Dam **19: 1**
River **2: 1**
- Southern recreation and conservation management priority areas, Focus on, by B. J. White **18: 1-24**
- Spotlighting party **10: 9**
- Spriggins, D. Forest recreation **5: 12-16**
- Starflower, see *Asterolasia pallida*
- Statistics—forest **3: 11, 14**
- Stone pine, see *Pinus pinea*
- Stream gauging station **19: 6**
- Sullivan Rock **5: 13**
- Summary of main plantation development operations for Blackwood Valley **5: 9**
- Sunklands multiple use land management, Focus on **16: 1-16, 18: 1-24**
- Swamp **10: 3, 11: 11**
treeless **19: 2**
Melaleuca preissiana **19: 2**
- Tachyglossus aculeatus* (echidna, spiny ant-eater, porcupine) **10: 8, 11: 13**
- Tallowwood, see *E. microcorys*
- Tammar, see *Macropus eugenii*
- Tarsipes spencerae* (honey possum, nool-benger) **10: 15, 14: 12**
- Tasmanian blue gum, see *E. globulus*
- Tetratheca viminea* (slender tetratheca) **17: 2**
- Thelymitra villosa* (custard orchid) **21: 21**
- Thornbill, yellow-tailed, see *Acanthiza chrysorrhoa*
- Timber-framed buildings **13: 9**
- Timber industry and the national economy **9: 14**
- Tingle
yellow, see *E. guilfoylei*
Rates, see *E. brevistylis*
red, see *E. jacksonii*
- Torwood, see *E. torquata* and *E. woodwardii*
- Trail bikes **14: 14, 15: 15**
- Transmission lines **4: 6**
- Transpiration studies **19: 14, 20: 16**
- Traveller's joy, see *Clematis pubescens*
- Trees are regulators of the environment **9: 15**
- Trees for country areas, by F. Batini **4: 12, 14**
- Trymalium ledifolium* **17: 2**
- Tuart, see *E. gomphocephala*
- Umbrella pine, *Pinus pinea*
- Verticordia picta* (painted featherflower) **17: 2**
- Wallaby
scrub, see *Macropus eugenii*
Western brush, see *Macropus irma*
- Wally's Nob **5: 4**
- Wandoo
botanical notes **3: 9-10, 16**
see *E. wandoo*
powderbark wandoo, see *E. accedens*
- Warren River **2: 2, 7, 8: 5-6**
- Washdown station, Dwellingup **21: 5**
- Waterbush, see *Bossiaea aquifolium*
- Water catchments **4: 2-9, 12**
map **4: 2, 19: 4**
- Wattle,
myrtle, see *Acacia myrtifolia*
see *Acacia*
- Wattle-bird, red, see *Anthochaera carunculata*
- What we get from trees—new publication **1: 14**
- Whim **7: 13**
- White, B. J.
Focus on shifting sands **8: 1-6, 10-11, 14-16**
Focus on southern recreation and conservation management priority areas **18: 1-24**
- Wildfire **3: 2, 5, 7-8, 17: 6**
- Willis, E. Early history of Jarrahdale **7: 2-3, 10-14**
- Wistaria, native, see *Hardenbergia comp-toniana*
- Woodchip project, Focus on marri **12: 1-16**
- Woodline train, Goldfields **15: 5**
- Woylie, see *Bettongia penicillata*
- Wren
blue, banded, splended, see *Malurus splendens*
red winged, see *Malurus elegans*
- Xanthorrhoea preissii* (blackboy) **14: 5**
- Yate
see *E. cornuta*
flat-topped yate, see *E. occidentalis*
- Yeagerup dunes **8: 1-6, 10-11, 14-16**
- Zamia palm, see *Macrozamia*
- Zonaeginthus oculus* (red-eared firetail) **7: 4**

