




INFORMATION SHEET

REPRINTED JANUARY 1977



EUCALYPTS

A Simplified Key to 17 W.A. Species

F. E. BATINI

Eucalyptus is by far the most important genus of Australian forest trees. Its members dominate 95 per cent of our forest area and spread out over much of the remainder of the country.

As would be expected in such a widespread genus, a great range in form is exhibited by members growing under different environmental conditions.

Identification of eucalypts is often not a simple task and a single characteristic is rarely sufficient for the purpose.

Over 600 species and variations have been named. Many different eucalyptus species have similar characteristics, and even the identifying characteristics on one particular tree or trees of the same species can vary: e.g. *E. loxophleba* (York Gum) in the key.

A positive identification can generally result only from the careful consideration of a number of features possessed by an individual tree. The most helpful of these are the general size and form; the nature of the bark on the trunk and branches; the adult leaf characters, including venation and oil dots; juvenile leaf forms; flower colour type and size, shape and structure of buds and fruits. Where naturally-occurring plants are concerned, the geographical location and habitat can be of great assistance in their identification as many species are strictly limited in their distribution.

In response to enquiries received by the Forests Department, this article includes a simple key for identification of 17 of Western Australia's more important species.


Example for using simplified key: Assuming we have a tree with a smooth, greyish-white bark, large fruits and exsert valves.

Under the major heading "Bark smooth", the specimen fits into the colour of bark category *greyish-white*. The large fruit and exsert valves would then place the specimen into the first of the next two categories, which means that it is *E. megacarpa*, or Bullich.

Explanatory notes

Eucalypt buds consist of the *operculum* which protects the developing stamens and which is eventually shed as these mature; and the *calyx* which develops into the fruit.

Operculum shorter than calyx



E. calophylla (Marri)


When the seeds are ripe, the valves in the fruit open and the seeds are released. The valves may either be *exsert* or *enclosed*.

Exsert valves




E. astringens (Brown Mallet)

Operculum longer than calyx



E. cornuta (Yate)


Enclosed valves




E. marginata (Jarrah)

Eucalypts—a simplified key to 17 W.A. species

BARK THICK, ROUGH AND PERSISTENT



BARK SMOOTH




BARK ROUGH ON TRUNK, SMOOTH ON LARGER BRANCHES


- Operculum much longer than calyx (4-5 times); distinctive beaked fruit *E. cornuta* (Yate)
- Operculum much longer than calyx (2-3 times), slightly ribbed, bell shaped fruit *E. occidentalis* (Flat-topped Yate)
- Operculum bluntly pointed, slightly longer than calyx *E. rudis* (Flooded Gum)
- Operculum shorter than calyx *E. loxophleba* (York Gum)

Buds and Fruits


(normal size)




E. marginata (Jarrah)



E. jacksonii (Red Tingle)




E. todtiana
(Coastal Blackbutt or Prickly Bark)



E. calophylla (Marri)




E. gomphocephala
(Tuart)




E. patens (Blackbutt)



E. loxophleba
(York Gum)




E. astringens (Brown Mallet)




E. salubris (Gimlet)




E. salmonophloia (Salmon Gum)




E. megacarpa (Bullich)




E. diversicolor (Karri)




E. accedens (Powderbark Wandoo)




E. wandoo (Wandoo)



E. cornuta (Yate)



E. occidentalis (Flat-topped Yate)



E. rudis (Flooded Gum)