

The flora of the Pyramid Lake - Mt Beaumont Districts, near Esperance, Western Australia

M.A. Burgman¹ and K.R. Newbey^{2*}

¹ Department of Botany, University of Western Australia, Stirling Highway, Nedlands, 6009
Present Address: Forestry Section, University of Melbourne, Creswick, Victoria 3363

² C/- Western Australian Herbarium, Department of Conservation and Land Management,
P.O. Box 104, Como, Western Australia 6152

Abstract

Burgman, M.A. and Newbey, K.R. The flora of the Pyramid Lake - Mt Beaumont Districts, near Esperance, Western Australia. *Kingia* 1(2): 217-253 (1990). A total of 1351 vascular plant taxa were identified from approximately 4000 collections made in the eastern Roe Botanical District between 1980 and 1984. Frequency and cover/abundance data were collected for each taxon in each of the eight landforms present. Life form, conservation status, representation on existing conservation reserves, and distribution within seven biogeographical regions were also recorded.

The flora was dominated by taxa from the Myrtaceae (17%), Proteaceae (8%) and Mimosaceae (8%). Dwarf shrubs (35%) were the dominant life form. Endemism was high (10%) and the number of exotic plants low (3%). The flora of the study area bears strongest relationship to the South-West Botanical Province, with 32% of the taxa also occurring there. Twenty per cent of the total taxa are also found in South Australia.

The flora contained 266 (20%) undescribed taxa and 149 taxa (11%) were considered to be rare, geographically restricted or very poorly known. Only a small proportion of the total flora (17%) and very few of the rare species were known to be adequately represented on existing conservation reserves or national parks. These facts emphasised the need for further detailed botanical studies before any land in the study area is considered for release for agricultural development.

Introduction

This study was undertaken to record the flora of land proposed for agricultural development north of Esperance. The Rural and Allied Industries Council (R.A.I.C. 1979) considered that there were 2.7 million hectares of vacant Crown Land available, which may be suitable for agricultural development in the Forrestiana-Lake Johnston and Salmon Gums-Israelite Bay areas. These areas included the Pyramid Lake and Mt Beaumont Districts, north of existing farmland between the No. 1 Rabbit Proof Fence, north-east of Ravensthorpe, and the western boundary of the Cape Arid National Park (Figure 1). The study area is largely within the Roe Botanical District, with a small south-eastern section within the Eyre Botanical District (Beard 1980). Farmland around Salmon Gums separates the two districts which together cover approximately 12,000 km². They are referred to collectively here as the study area. Since the R.A.I.C. (1979) report was released, some land in the Mt Beaumont District, and near Cascades in the Pyramid Lake District was made available for agriculture. However, the State Government presently has a moratorium on further land release.

* Deceased July 23, 1988




Figure 1. Map of the Study Area

The flora of the study area has not been documented. An earlier assessment of conservation reserves by the Conservation Through Reserves Committee (1974) did not recommend that any sections of the study area be considered for preservation, undoubtedly because very few collections of flora and fauna had been made there. A private overview survey (by KRN) of the study area recorded a number of rare species and a few new to science. This deficiency in biological data was recognized by staff of the Western Australian Wildlife Research Centre, and led to successful applications by S.D. Hopper for funds for botanical survey work from the Australian Biological Resources Study and from the Western Australian Government. Consultancies arising from these grants were awarded to us to undertake appropriate surveys.

The specific aims of this study were to:

- a) list all taxa recorded by us during four surveys carried out since 1980;
- b) assess the frequency and cover/abundance of each taxon on each landform;
- c) assess the distribution of each taxon within southern Western Australia;
- d) assess the conservation status of each taxon (re-assessing the conservation values for those taxa listed by Marchant and Keighery 1979, Leigh *et al.* 1981, Leigh *et al.* 1984, Rye and Hopper 1981, Patrick and Hopper 1982, Rye 1982);
- e) assess the status of plants recorded by us from the Frank Hann National Park (KRN), Peak Charles National Park (KRN) and conservation reserves within the study area (MAB);
- f) list those taxa considered to be rare or geographically restricted, for gazettal as rare flora;
- g) assess the completeness of our flora list and the extent of invasion of exotic taxa;
- h) present a life form spectrum for the taxa collected.

Biophysical Environment

Climate

The climate may be described as semi-arid Mediterranean (Specht and Moll 1983), with most of the rain falling between May and October. The climate at Salmon Gums in the north of the study area is only weakly Mediterranean and tends towards non-seasonality (Beard 1973a,b). As there are no daily recording meteorological stations within the study area, the spatial and temporal rainfall patterns are poorly known. Average annual and monthly rainfall, and maximum and minimum temperatures for nearby stations at Lake King, Grass Patch, Salmon Gums, Esperance, Balladonia and Ravensthorpe are provided by Hall *et al.* (1981) and Newbey (1983a). Estimated average annual rainfall varies from 670 mm near the southern coast to 300 mm inland. Snow is a rare event (one year in 20) and a few days each summer exceed 40°C.

Geology

Bedrock of Archaean granite underlies the study area, but it is largely covered by soil. The tectonically stable bedrock was eroded into a series of subdued uplands dissected by broad, saucer-shaped valleys. Most of the valleys drained to the south coast but a few on the north-eastern section of the study area drained east and north to the Nullarbor Plain (Van de Graaff *et al.* 1977). During the Cretaceous, rainfall declined and the valleys were reduced to a string of salt lakes. A marine transgression during the Eocene filled river valleys with sediments (spongolite), and most of the land surface was eroded flat (Cockbain 1968). With the fall in sea level to the present, rivers in the western half of the study area (the Young, Lort and Oldfield Rivers) have eroded the spongolite back to

bedrock. Granite exposures represent later intrusions of Proterozoic granite through the Archaean bedrock. They have since been eroded into rounded hills conspicuous on the marine plain (Morgan and Peers 1973; Thom *et al.* 1977).

Landforms and soils

Landform classification follows the system devised by K.R. Newbey and A.V. Milewski for the biological survey of the eastern goldfields of Western Australia (Newbey 1984). Each landform with its associated soils is described below. Soils weathered *in situ* from the granitoid basement are slightly acidic and often highly leached. Soils of the marine plain have a high CaCO₃ content, due to deposition of salts leached from the uplands. Aeolian and alluvial soils are associated with valley bottoms and salt lakes. Aeolian soil sheets, some with fossil dunes, are present on some sections of the marine plain.

Breakaway (B): In a few places in the study area, the underlying kaolinized granite is exposed on the ancient land surface. A breakaway consists of the following elements: rim, free face, scree slope, and pediment. The summit and rim consist of material more resistant to weathering than the underlying material and this controls the breakaway morphology. Shallow to skeletal gritty sands cover the summit and fill small pockets on the rim. Bedrock fragments in skeletal gritty sands cover the scree slopes. The pediment consists of finer material washed off the other elements. Soil drainage is excessive on the summit, rim, free face and scree slopes. However, the pediment obtains additional moisture from run-off from these elements. Soil pH is generally 6.0-6.5.

Drainage Line (D): This landform is confined to the Pyramid Lake District and occurs along the upper reaches of the Lort, Young and Oldfield Rivers. These saline watercourses, with ephemeral flows, drain to the south coast. Drainage lines dissect the sandplains and their common boundary is where the river banks change from erosional to depositional (colluvial). This varies from 2 to 10 metres above the river bed. Soils are colluvial, or alluvial on narrow river flats. They are usually sandy, and saline within a few metres of the river channel. Soil pH is generally 6.0-6.5.

Flat Plain (F1 and F2): Most of the study area east of Pyramid Lake consists of plain levelled by the Eocene Sea (see Marine Plain below) with an internal relief that rarely exceeds 5 m. A wide range of soils are present which can be divided into two relatively distinctive groups having unique vegetation and floristics (F1 and F2). Both may contain small freshwater swamps up to 250 m across that fill to a depth of 1 m only after very heavy rain.

F1 consists of nutrient poor and slightly acidic sands and sandy gravels. They are duplex soils with a sandy clay B horizon that is usually slightly to highly calcareous but occasionally slightly acidic. Fossil aeolian sand sheets, sometimes with dune relicts, are present in some areas.

F2 consists of loams to clays with significantly higher nutrient levels than F1 soils. The profile is calcareous to highly calcareous throughout with a clay loam B horizon often with a higher pH than the A horizon. Carbonate nodules are often present in the B horizon.

Granite Exposure (G): This landform is most common in the Mt Beaumont district of the study area and it is present in two phases related to the origin of the rock. Most common are the Proterozoic granite intrusions now present as rounded hills. Exposures of granite bedrock are less common, but both phases support similar vegetation on similar landform elements. Skeletal soil sheets may be present on the exposures and small hollows may contain ephemeral pools. Surrounding the exposure is an inner apron of soil less than 50 cm deep. Peripheral to this is an outer apron of soil 50-150 cm deep.

Sometimes, seepage occurs where soil has accumulated in faint drainage lines on the exposure. Soil pH is generally 6.0-6.5.

Salt Lake Features (L): Lakes in the study area have two origins. Some lakes occur on flat bottoms of paleo-drainage lines, interspersed with lunettes and saline flats. Lunettes range in composition from quartz rich sands to gypseous clays or nearly pure gypsum (Bowler 1982). The second type of lakes are linear, being separated by almost parallel vegetated sand dunes. Both types consist of lake floors and margins partially enclosed by peripheral dunes composed of sand or clay loam. Pure quartz dunes are formed under lake-full conditions, and the others by deflation of the adjacent lakes. Peripheral soils vary from saline to sub-saline and the pH may vary from highly acidic to highly alkaline. Lake floors are often devoid of vegetation.

Marine Plain (M): Surrounding Boyatup Hill is a marine plain covered by up to 2 m of fine siliceous sand. Below the sand is a B horizon of clay loam to 1 m thick that grades into spongolite bedrock. This plain differs from Flat Plain in having a slightly acidic B horizon and the extensive deep cover of sand. It is also in a higher rainfall zone.

Sandplain (S): Overlying the granite bedrock west of Pyramid Lake is an ancient landscape with a highly leached and slightly acidic soil profile. Slopes on the plain are gentle and internal relief is less than 30 m. The soil B horizon consists of sandy clay that is often mottled near its upper surface. The A horizon may consist of sand, sandy loam, clayey sand or clay loam. In some places the sand is present as colluvial sheets up to 1.5 m thick. Clay loams are usually associated with gilgai in localised areas.

Vegetation

Beard (1973a,b) mapped the main vegetation types of the study area using a structural classification with floristic sub-units. Monk *et al.* (1979) described the vegetation of the Frank Hann National Park, and Bennett (1983) outlined many vegetation types at Mt Beaumont, adjoining one of the areas recently released for agriculture in the study area.

The landform types described above support characteristic structural vegetation types. Breakaways have a complex of shrubs and annuals on skeletal soils, and mallees and trees are sometimes present. Drainage lines support a variety of vegetation, related to water logging and salinity of soils, from dwarf shrubland, to thicket and woodland. Similarly, salt lake floors are often bare, fringed by low halophytic shrubs. Tall shrubs are present on better drained soils, while mallees occur on soils with low salt content. Vegetation on Granite Exposures is also variable, related to soil depth and water run-off.

Flat Plains (F1) and Marine Plains have vegetation dominated by shrubs and sedges less than 1 m tall, although scattered tall shrubs and mallees are sometimes present. Flat Plains (F2) support denser mallee with fewer shrubs, and small patches of low woodland on highly calcareous soils. Scattered to dense mallees cover most of the Sandplains. In all mallee communities, the low and medium shrub strata tend to be more dense where the mallee cover is lower or more sparse. Woodlands are present on soils with higher nutrient status and numerous, small, ephemeral swamps support open woodlands.

Flora

No comprehensive plant collections were made in, or near, the study area until Newbey (1979) collected widely on the central south coast around the Fitzgerald River. More recently, Clements and George (1980) led an expedition which collected extensively for orchids. Ecological studies have

also been rare (Beard 1967, Hopper and Moran 1981, Hopkins and Robinson 1981, Crook and Burbidge 1982).

Methods

The flora list includes all plant taxa recorded by us mainly during four surveys:

- 1) a short preliminary survey was undertaken by KRN in November 1980 to overview the flora;
- 2) searches were made by KRN in the summer of 1982-83 for rare plants (Newbey 1983b);
- 3) basic data were recorded in October 1983 by KRN for a land use survey of the North Boyatup Hill area (Newbey unpublished data);
- 4) fourteen trips were undertaken in 1983 and 1984 by MAB to record as much as possible of the flora (Burgman 1985a).

Species lists were compiled at more than 300 sites subjectively selected to represent all of the main vegetation types of the study area. Opportunistic collections were made between sites and searches were made of unusual landforms. Data on each taxon were collected using classification schemes to record frequency and cover/abundance (Newbey and Hnatiuk 1984), life form (Newbey 1979), conservation status (Leigh *et al.* 1981) and representation on reserves (Rye and Hopper 1981).

No detailed searches were made for plants considered to be rare. Rather estimates of rarity were made on known population sizes and distributions, and on our general knowledge of the study area. Previous publications on rare plants in Western Australia (see above) have been restricted to published taxa. We have included undescribed taxa, referenced by voucher collections in the Western Australian Herbarium (PERTH). The representation of taxa on reserves was restricted to Frank Hann and Peak Charles National Parks, and to flora conservation reserves in the study area. Cape Arid National Park adjoins the study area in the east, and Cape Le Grand and Stokes Inlet National Parks are to the south. The floras of these areas have not been documented and to do so was beyond the scope of this study.

Results and Discussion

A total of about 4000 plant specimens were collected in the study area as part of the four surveys. From these, 1351 different vascular plant taxa were identified (Appendix 1) which represent approximately 15% of the State's named flora (Green 1985). Plant taxa in the study area consisted of five species of ferns and fern allies, and 1290 species, 24 subspecies and 32 varieties of flowering plants. Families with the most taxa were the Myrtaceae (231 taxa), Proteaceae (106 taxa), Mimosaceae (105 taxa), Asteraceae (75 taxa) and Epacridaceae (62 taxa). The genera with the most taxa were *Acacia* (105 taxa), *Eucalyptus* (77 taxa), *Melaleuca* (67 taxa), *Leucopogon* (37 taxa), *Grevillea* (28 taxa), *Hibbertia* (21 taxa) and *Pultenaea* (19 taxa).

Life form spectrum

One of the most notable features of the species list was the predominance of dwarf shrub species, which make up 35% of the total taxa (Table 1). Kwongan (dwarf shrubland) is the richest and most complex vegetation type in the study area (Burgman 1985a) and contributes greatly to its species richness. Eucalypts are dominated by mallee-form species. Mature mallee is the most extensive vegetation type, although it is relatively poor in numbers of species (Burgman 1985a). Only a few fire ephemerals were present in recently burnt mallee and kwongan vegetation.

Table 1. Proportion of taxa (%) in Appendix 1 contributing to different life forms (including data from Newbey 1979 and Raunkiaer 1934). A key to the life form codes is provided in Appendix 1. + = less than 1%

P										CH	H					G		T		P		
MM		M			N					CH	H					G		T		P		
ST	MT	LT	DT	TS	MA	DS	SS	MS	LS	HP	CL	MP	RP	PG	SC	SI	SL	AB	HY	AG	AS	PC
1	+	-	1	3	4	35	13	8	2	2	1	2	1	2	1	5	2	5	+	1	10	+
70						2						11					5		11		0	
This Study																						
+	+	+	1	4	2	27	17	6	4	1	1	3	3	2	3	6	2	6	+	1	10	1
63						3						16					6		11		1	
Newbey (1979)																						
46						9						26					6		13		0	
Raunkiaer (1934)																						

Phanerophytes are strongly represented in the life form spectrum, compared to the data of Raunkiaer (1934), at the expense mainly of chamaephytes and hemicryptophytes. This is also true when compared to Newbey's (1979) central south coastal flora, and reflects the large contribution made by dwarf shrubs to the species richness of the study area.

Undescribed taxa

An important feature of the species list is the large number of undescribed taxa. Some 266 (20%) of the total are undescribed, in manuscript, or of uncertain status. A few taxonomic groups are outstanding for the number of undescribed taxa in them, notably *Lepidosperma* (4 taxa out of a total of 16), Rhamnaceae (7/20), *Hibbertia* (11/21), and *Melaleuca* (28/67). Additionally, a few genera are currently under revision, including *Acacia* (38/105), *Darwinia* (9/11), *Eucalyptus* (23/77) and *Leucopogon* (21/37). Recent revisions of genera in Western Australia have highlighted that most undescribed taxa are rare or have restricted geographic ranges (e.g. George 1981). The same appears likely to be true of many unnamed taxa recorded in the study area.

Biogeographic relationships

There are 44 exotic species listed in Appendix 1 (3% of the total), including 15 species from the Poaceae and 9 from the Asteraceae. This differs from the 494 taxa collected by Cheal *et al.* (1979) in which 10% of the mallee vegetation of north-western Victoria were exotic. However, Specht (1972) found that, like the study area, the incidence of introduced plants was rare in the South Australian mallee.

The low number of exotics in Appendix 1 is evidence of the low level of invasion of introduced plants in the study area. *Hypochaeris glabra* was ubiquitous, though rarely abundant. Most others were annual grasses or short-lived perennials. Usually, they were restricted to disturbed sites, or to salt lakes or granite rocks, probably responding to the more mesic micro-climates or higher nutrient status of these habitats.

Of the taxa in Appendix 1, 275 (20%) are also found in South Australia. Of these, 123 (9%) have disjunct and apparently vicariant distributions across southern Australia (Jessop 1984). The close relationship between the floras of South Australia and Western Australia is well known (Green 1964, Beard 1969, Parsons 1970, Nelson 1974), and is reflected in these figures.

As many as 10% of the taxa collected here are endemic to the study area and a further 9% are restricted to the Roe Botanical District. The flora of the study area bears strongest relationship to the South-West Botanical Province, with 32% of the taxa also occurring there. The relationship between the study area and the south coastal Eyre Botanical District is also strong, with 21% of the taxa shared; relatively fewer plants (13%) are shared with the more arid Coolgardie Botanical District.

Conservation values

Some 149 taxa (11%) were considered to be rare, geographically restricted or very poorly known and only 50 of these were listed by any one of the authorities on rare plants in Western Australia (see above). There were 266 taxa (20%) considered to be good taxonomic species which could not be matched with any named species. While they may be species new to science, their status is uncertain and will remain so until the taxa to which they appear to belong are revised.

Seven species in Appendix 1 have been gazetted as rare (Government of Western Australia 1988). They are *Adenanthos ileticos*, *Billardiera mollis*, *Eremophila denticulata*, *E. serpens*, *Eucalyptus merrickiae*, *Kennedia beckxiana* and *Ricinocarpos trichophorus*.

There are 38 taxa in Appendix 1 considered to be endangered and 45 taxa considered to be vulnerable. In order to highlight them, they are listed separately in Appendix 2. Endangered species require immediate survey and implementation of conservation measures, and species classified as vulnerable will need the same attention should land be considered for release for development. Most endangered taxa are recently described or undescribed. Only five of them have been listed by any one of the authorities on rare plants. These omissions are a result of the rarity of the taxa and their uncertain taxonomic status, which highlights the need for more effort to be spent on surveying rare plants.

Of the 98 taxa in Appendix 1 also listed in the existing publications on Western Australian rare plants, 50 were considered here to be relatively widespread or abundant. Examples of this kind are *Callitris columellaris*, *Bossiaea leptacantha* and *Oxylobium microphyllum*.

Of the 1351 taxa in Appendix 1, 59% are known to occur on conservation reserves or national parks but only 17% are known to be adequately represented (Rye and Hopper 1981). There are only five endangered species known on reserves, one of which is adequately represented. Similarly, there are only 23 vulnerable species known on reserves, two of which are adequately represented. A further 35 rare or poorly known species have been recorded on reserves.

Factors other than distribution and abundance contribute to the biological importance of collections. Range extensions in such species as *Acacia warramaba* may represent a paucity of collections

whereas in others such as *Acacia sorophylla*, they represent isolated populations. They may represent populations of species previously widespread in the wheatbelt (e.g. *Comesperma acerosum*, *Dampiera carinata*, *Goodenia trichophylla*), or forms, varieties or subspecies new to science (Burgman 1985b).

Conclusions

It is important to emphasise that Appendices 1 and 2 are incomplete. As many as 266 taxa are of uncertain status and it is anticipated that a significant proportion of these will be new, rare or restricted. Furthermore, by comparing Appendix 1 with other lists from nearby areas, we estimate that 15-20% of the flora of the study area remains to be collected. Clearly, a great many more records of rare or undescribed taxa remain to be made.

The study area exhibits a high degree of endemism, a large proportion of the flora is not known to occur on reserves and less than 20% is known to be adequately represented. Furthermore, very little is known of the biogeography of the study area and information of this nature will have important implications for the design of nature reserves. These factors suggest that very detailed studies are necessary, before land is released for development, if the flora is to be conserved.

Acknowledgements

We both would like to thank D. Bell, A. Hopkins, W. Loneragan and S. Hopper for their comments on the manuscript. W. Loneragan and S. Hopper provided advice throughout the study and P. van der Moezel provided valuable discussion. We are grateful to B.R. Maslin for liaising with the Western Australian Herbarium on our behalf.

Professional staff at the Herbarium assisted with difficult identifications. Outside assistance was sought from A.A. Burbidge (*Beaufortia*), M. Trudgen (*Baekkea* and related taxa), M.I. Brooker (*Eucalyptus*), J. Powell (*Leucopogon*), S.D. Hopper (*Conostylis* and Orchidaceae), M.D. Crisp (*Daviesia*) and A.S. George (*Dryandra*).

C.E. Layman, S. McNee, B. Tanahill, A. Napier, and F. and J. Phillips provided assistance in the field and in the herbarium. B. Smith typed the manuscript. J.W. Green and A.A. Burbidge kindly provided access to facilities at the Western Australian Herbarium and the Wildlife Research Centre respectively. This study was funded through the W.A. Wildlife Research Centre (KRN) and by a grant to S.D. Hopper from the Australian Biological Resources Study (MAB).

References

- Beard, J.S. (1967). A study of patterns in some Western Australian heath and mallee communities. *Australian Journal of Botany* 15: 131-139.
- Beard, J.S. (1969). Endemism in the Western Australian flora at the species level. *Journal of the Royal Society of Western Australia* 52: 18-20.
- Beard, J.S. (1973a). The vegetation of the Ravensthorpe area, Western Australia. SI: 51-5. Vegmap Publications, Perth.
- Beard, J.S. (1973b). The vegetation of the Esperance and Malcolm areas, Western Australia. SI: 51-6, SI:51-7. Vegmap Publications, Perth.
- Beard, J.S. (1980). A new phytogeographic map of Western Australia. *Western Australian Herbarium Research Notes* 3: 37-58.
- Bennett, E.M. (1983). The vegetation and flora of vacant Crown Land in the Beaumont land release area, Esperance Shire. Western Australian Department of Fisheries and Wildlife, unpublished report.

- Bowler, J.M. (1982). Aridity in the late Tertiary and Quaternary of Australia. In: W.R. Barker and P.J.M. Greenslade (eds), Evolution of the flora and fauna of arid Australia. Peacock Publications, South Australia.
- Burgman, M.A. (1985a). The flora, ecology and biogeography of the eastern Roe Botanical District. Western Australian Department of Conservation and Land Management, unpublished report No. 5.
- Burgman, M.A. (1985b). Cladistics, phenetics and biogeography of populations of *Boronia inornata* Turcz. (Rutaceae) and the *Eucalyptus diptera* Andrews (Myrtaceae) species complex in Western Australia. Australian Journal of Botany 33: 419-431.
- Cheal, P.D., Day, J.C. and Meredith, C.W. (1979). Fire in the national parks of north-west Victoria. National Parks Service, Victoria.
- Clements, M.A. and George, A.S. (1980). Report of the Australian Orchid Foundation expedition to Western Australia. National Botanic Gardens, Canberra.
- Cockbain, A.E. (1968). The stratigraphy of the Plantagenet group of Western Australia. Western Australian Geological Survey, Annual Report 1967, pp. 61-63.
- Crook, I.G. and Burbidge A.A. (1982). Lake Magenta Nature Reserve. Western Australian Nature Reserve Management Plan No. 4. Department of Fisheries and Wildlife, Western Australia.
- Conservation Through Reserves Committee. (1974). Conservation reserves in Western Australia. Environmental Protection Authority, Perth.
- George, A.S. (1981). The genus *Banksia* L.f. (Proteaceae). Nuytsia 3: 239-473.
- Government of Western Australia. (1988). Government gazette of Western Australia, July 15, 1988.
- Green, J.W. (1964). Discontinuous and presumed vicarious plant species in southern Australia. Journal of the Royal Society of Western Australia 47: 25-32.
- Green, J.W. (1985). Census of the vascular plants of Western Australia, ed. 2. Western Australian Herbarium, Department of Agriculture, Perth.
- Hall, N., Wainwright, R.W. and Wolf, L.J. (1981). Summary of meteorological data in Australia. CSIRO Division of Forest Research, Report No. 6.
- Hopkins, A.J.M. and Robinson, C.J. (1981). Fire induced structural change in a Western Australian woodland. Australian Journal of Ecology 6: 177-188.
- Hopper, S.D. and Moran, G.S. (1981). Bird pollination and the mating system of *Eucalyptus stoatei*. Australian Journal of Botany 29: 625-638.
- Jessop, J.P. (ed.). (1984). A list of the vascular plants of South Australia, ed. 2. Adelaide Botanic Gardens and State Herbarium, and the Environmental Survey Branch, Department of Environment and Planning, Adelaide.
- Leigh, J., Boden, R. and Briggs, J. (1984). Extinct and endangered plants of Australia. MacMillan, Melbourne.
- Leigh, J., Briggs, J. and Hartley, W. (1981). Rare or threatened Australian plants. Australian National Parks and Wildlife Service. Special Publication No. 7.
- Marchant, N.G. and Keighery, G.J. (1979). Poorly collected and presumably rare vascular plants of Western Australia. Kings Park Research Notes 5, pp. 1-103.
- Monk, D., Hnatiuk, R.J. and George, A.S. (1979). Vegetation survey of the Frank Ham National Park. Western Australian Herbarium Research Notes 2: 23-49.
- Morgan, K.H. and Peers, R. (1973). Esperance-Mondrain Island, 1:250,000 Geological Series, explanatory notes. Geological Survey of Western Australia, SI: 51-6. Australian Government Publishing Service, Canberra.
- Muir, B.J. (1977). Vegetation and habitat of Bendering Reserve. In: Biological survey of the Western Australian wheatbelt, part 2. Records of the Western Australian Museum Supplement No. 3, pp. 3-142.
- Nelson, E.C. (1974). Disjunct plant distributions of the south-western Nullarbor Plain, Western Australia. Journal of the Royal Society of Western Australia 57: 105-117.
- Newbey, K.R. (1979). The vegetation of central south coastal Western Australia. M. Phil. thesis, Murdoch University, Western Australia.
- Newbey, K.R. (1983a). Rainfall of the Southern Cross - Mt. Ragged area, Western Australia. Ongerup Land Use Research Centre Bulletin No. 1, pp. 1-17.
- Newbey, K.R. (1983b). Some important plant species in natural areas north of agricultural areas between Ravensthorpe and Esperance. Private report to Western Australia Department of Fisheries and Wildlife.

- Newbey, K.R. (1984). Physical environment. *In*: The biological survey of the eastern goldfields of Western Australia, part 2: Widgiemooltha - Zanthus study area. Records of the Western Australian Museum Supplement No. 18, pp. 29-40.
- Newbey, K.R. and Hnatiuk, R.J. (1984). Vegetation and flora. *In*: The biological survey of the eastern goldfields of Western Australia, part 2: Widgiemooltha - Zanthus study area. Records of the Western Australian Museum Supplement No. 18, pp. 41-51.
- Parsons, R.F. (1970). Mallee vegetation of the southern Nullarbor and Roe Plains, Australia. Transactions of the Royal Society of South Australia 94: 227-242.
- Patrick, S.J. and Hopper, S.D. (1982). A guide to the gazetted rare flora of Western Australia, Supplement No. 1. Western Australian Department of Fisheries and Wildlife, Report No. 54.
- R.A.I.C. (1979). Rural and Allied Industries Council, rural land release policy in Western Australia. Premiers Department, Western Australia.
- Raunkiaer, C. (1934). The life forms of plants and statistical plant geography. Oxford University Press, London.
- Rye, B.L. (1982). Geographically restricted plants of southern Western Australia. Western Australian Department of Fisheries and Wildlife, Report No. 49.
- Rye, B.L. and Hopper, S.D. (1981). A guide to the gazetted rare flora of Western Australia. Western Australian Department of Fisheries and Wildlife, Report No. 42.
- Specht, R.L. (1972). The vegetation of South Australia, ed. 2. Government Printer, Adelaide.
- Specht, R.L. and Moll, E.J. (1983). Mediterranean type heathlands and sclerophyllous shrublands of the world: an overview. *In*: R.L. Specht (ed.), Heathlands and related shrublands of the world. B. Analytical studies, pp. 5-13. Elsevier, Amsterdam.
- Thorn, R., Lipple, S.L. and Sanders, C.C. (1977). Ravensthorpe 1:250,000 Geological Series, explanatory notes. Geological Survey of Western Australia, SI: 51-5. Australian Government Publishing Service, Canberra.
- Van de Graaff, W.J.E., Crowe, R.W.A., Bunting, J.A. and Jackson, M.M. (1977). Relict early Cainozoic drainage in arid Western Australia. Zeitschrift für Geomorphologie 21: 379-400.

Appendix 1. Flora list

Arrangement of taxa follows Green (1985). Taxa listed below but not appearing in Green (1985) are: *Brizula muelleri*, *Eucalyptus scyphocalyx*, *Glyceria fluitans*, *Grevillea coccinea* and *Westringia rigida*. Data listed are life form, representation on conservation reserves, assessment of frequency and cover/abundance on landforms, and distribution. Voucher specimens of most taxa have been deposited in PERTH. Unnamed taxa are referenced by MAB or KRN collecting numbers. An asterisk indicates an introduced taxon.

LF = Life Form (Newbey 1979)

	Symbol
A. PHANEROPHYTES	P
Mesophanerophytes (trees 5-50 m high)	MM
1. Small trees (5-15 m)	ST
2. Medium trees (15-30 m)	MT
3. Large trees (greater than 30 m)	LT
Microphanerophytes (trees and shrubs 2-5 m high)	M
4. Dwarf trees (less than 5 m)	DT
5. Tall shrubs (greater than 2 m)	TS
6. Mallees - tree form	MAT
shrub form	MAS
Nanophanerophytes (shrubs less than 2 m high)	N
7. Dwarf woody shrubs (less than 0.5 m)	DS
8. Small woody shrubs (0.5-1 m)	SS
9. Medium woody shrubs (1-1.5 m)	MS
10. Large woody shrubs (1.5-2 m)	LS
11. Herbaceous shrubs	HP
12. Climbers	CL
B. CHAMAEPHYTES	CH
13. Mat plants	MP
C. HEMICRYPTOPHYTES	H
14. Rosetted perennials	RP
15. Perennial grasses	PG
16. Colonial sedges	SC
17. Tufted sedges	SI
18. Sedge-like plants	SL
D. GEOPHYTES	G
19. Terrestrials	AB
20. Hydrophytes	HY
E. THEROPHYTES	T
21. Annual grasses	AG
22. Other annuals	AS
F. PARASITIC CLIMBERS	P
23. Parasitic climbers	PC

Appendix 1 (continued). Flora list

RES = Presence on conservation reserve

F	=	Frank Hann National Park (Newbey unpublished data)
P	=	Peak Charles National Park (Newbey unpublished data)
S	=	Nature reserves within study area (Burgman 1985a)
Assessment		
.	=	not recorded
P	=	present, inadequate representation (<1000 mature plants)
A	=	adequate representation (>1000 mature plants)

CS = Conservation status (based on code developed by Leigh *et al.* (1981) but applied by Burgman and Newbey)

1	=	taxa known only from type collection, or a single collection.
2	=	very restricted distribution, range <100 km.
3	=	taxa with distribution >100 km but occurring only in small populations.
E	=	endangered taxon in serious risk of disappearing from the wild state within one or two decades if present land use and other causal factors continue to operate.
V	=	vulnerable taxon not presently endangered but at risk over a longer period.
R	=	rare taxon not currently considered endangered or vulnerable.
K	=	poorly known taxon suspected to belong to one of the above categories.
C	=	present within national park or other proclaimed conservation area.

LANDFORM

B	=	Breakaway
D	=	Drainage line
F1	=	Plain (mainly sandy A horizon, neutral B horizon)
F2	=	Plain (various A horizon, calcareous B horizon)
G	=	Granite exposure
L	=	Salt lake feature
M	=	Marine plain
S	=	Sandplain

Frequency and Cover/abundance (canopy cover - see Muir (1977))

Frequency		Cover/abundance	
A	=	1 or 2 populations	1 = 1 or 2 plants
B	=	Few populations	2 = Few plants
C	=	Scattered populations	3 = Few plants to 1% canopy cover
D	=	Frequent populations	4 = 1-5% canopy cover
E	=	Common populations	5 = 6-30% canopy cover
			6 = 31-70% canopy cover

DIS = Distribution (Unnamed taxa assumed restricted to Western Australia)

1	=	Endemic to study area
2	=	Roe Botanical District (Beard 1980)
3	=	Eyre Botanical District
4	=	South-West Botanical Province (excluding Eyre Botanical District)
5	=	South-Western Interzone (Coolgardie Botanical District)
6	=	Southern Australia (mainly South Australia (Jessop 1984))
7	=	Naturalised aliens

Appendix 1 (continued). Flora List

LP	RES	LANDFORM										DIS	
		FFS	CS	B	D	F1	F2	G	L	M	S		
PTERIDOPHYTA													
4	ISOETACEAE												
HY	<i>Isoetes australis</i> S. Williams	C2	2,5
7	ADIANTACEAE												
DS	<i>Cheilanthes austrotenuifolia</i> H. Quirk & T.C. Chambers	.AA	..	C3	C3	6
DS	<i>Cheilanthes distans</i> (R. Br.)Mett.	.A.	A1	6
11E	ASPLENIACEAE												
DS	<i>Asplenium flabellifolium</i> Cav.	..P	B2	6
DS	<i>Pleurosorus rutifolius</i> (R. Br.)Fee	..PP	B2	6
16A	ZAMIACEAE												
SS	<i>Macrozamia riedlei</i> (Fischer ex Gaudich.)C. Gardner	A3	4
GYMNOSPERMAE													
17A	PINACEAE												
ST	* <i>Pinus pinea</i> L.	..P	A1	7
18	CUPRESSACEAE												
DT	<i>Callitris columellaris</i> F. Muell.	A2	6
TS	<i>Callitris preissii</i> Miq. ssp. <i>verrucosa</i> (Cunn. ex Vogel)J. Garden	AAA	C2	C4	..	6
TS	<i>Callitris roei</i> (Endl.)F. Muell.	P.A	A1	..	4
MONOCOTYLEDONAE													
26	JUNCAGINACEAE												
AS	<i>Triglochin calcitrapa</i> Hook.	..P	D4	6
AS	<i>Triglochin minutissima</i> F. Muell.	D4	4
AS	<i>Triglochin mucronata</i> R. Br.	..P	C3	6
AS	<i>Triglochin muelleri</i> Buchenau	..	3K	B2	6
31	POACEAE												
AC	<i>Agrostis</i> sp. A (MAB 2417)	A2	1
AC	* <i>Aira cupaniana</i> Guss.	AA.	..	B3	B3	..	B2	C3	7
PG	<i>Amphipogon amphipogonoides</i> (Steudel)Vick.	..P	B2	..	4
PG	<i>Amphipogon turbinatus</i> R. Br.	APP	B2	..	4
PG	<i>Amphipogon</i> sp. A (MAB 3909)	4
PG	<i>Aristida contorta</i> F. Muell.	..P	B4	6
AG	* <i>Avellinia michelii</i> (Savi)Parl.	..P	C2	7
AG	* <i>Briza minor</i> L.	..P	B2	7
AG	* <i>Bromus rubens</i> L.	..P	B2	7
PG	<i>Danthonia caespitosa</i> Gaudich.	..PA	B2	6
PG	<i>Danthonia occidentalis</i> Vick.	..P	B2	4
PG	<i>Danthonia setacea</i> R. Br. var. <i>setacea</i>	..A	B2	D2	6
PG	<i>Danthonia setacea</i> R. Br. var. <i>brevisetata</i> Vick.	C3	4
PG	<i>Danthonia</i> sp. A (MAB 2463)	A2	4
PG	* <i>Ehrharta calycina</i> Smith	..P	B2	7
PG	* <i>Ehrharta longiflora</i> Smith	..P	B2	7
PG	<i>Eragrostis dielsii</i> Pilger ex Diels & E. Pritzel	..P	B2	C2	6
AG	<i>Glyceria fluitans</i> R. Br.	..	2V	A1	1
AG	* <i>Hordeum leporinum</i> Link	B2	7
PG	* <i>Lolium perenne</i> L. (L. ? x <i>L. rigidum</i> Gaudin)	A2	7
AG	* <i>Lolium rigidum</i> Gaudin	B3	7
PG	<i>Neurachne alopecuroidea</i> R. Br.	..APA	C3	C2	6
AG	* <i>Parapholis incurva</i> (L.)C.E. Hubb.	..P	B2	7
PG	<i>Paspalidium gracile</i> (R. Br.)Hughes	A1	6
AG	* <i>Pentaschistis airoides</i> (Nees)Stapf	..AAA	..	C2	C3	B3	7
PG	<i>Poa drummondiana</i> Nees	..PP	..	A1	6
PG	<i>Poa</i> sp. A (MAB 4533)	..P	A1	2
AG	<i>Puccinellia stricta</i> (J.D. Hook.)C. Blom	..P	A2	6
PG	<i>Spartochloa scirpoidea</i> (Steudel)C.E. Hubb.	..FAA	D4	2,5
PG	<i>Stipa acrociliata</i> Reader	A2	2,5
PG	<i>Stipa compressa</i> R. Br.	..P	B3	2,5
PG	<i>Stipa elegantissima</i> Labill.	..P	B1	C1	6
PG	<i>Stipa eremophila</i> Reader	..P	B2	6
PG	<i>Stipa flavescens</i> Labill.	B2	6
PG	<i>Stipa hemipogon</i> Benth.	..A	A1	B2	6
PG	<i>Stipa juncifolia</i> Hughes	..P	C3	2,3
PG	<i>Stipa pycnostachya</i> Benth.	B2	B3	2,3
PG	<i>Stipa semibarbata</i> R. Br.	A2	6
PG	<i>Stipa trichophylla</i> Benth.	B3	B2	6
PG	<i>Stipa variabilis</i> Hughes	..A	C3	6
PG	<i>Stipa</i> sp. A (MAB 3570)	A2	2

Appendix 1 (continued). Flora List

LF	REF	LANDFORM										DIS
		FPS	CS	B	D	F1	F2	G	L	M	S	
PG	<i>Stipa</i> sp. B (MAB 3086)	B2	2
PG	<i>Triodia concinna</i> N. Burb.	..P	1K	A5	5
AG	* <i>Trisetaria cristata</i> (L.)Kerguelen	..P	B2	7
AG	* <i>Vulpia bromoides</i> (L.)Gray	..P	C2	7
AG	* <i>Vulpia myuros</i> (L.)C. Gmelin	A.A	B2	D4	7
AG	* <i>Vulpia</i> sp. A (MAB 3223A)	B2	7
32 CYPERACEAE												
SC	<i>Baumea rubiginosa</i> (Sprengel)Boeckler.	..P	A3	..	6
SI	<i>Caustis dioica</i> R. Br.	PAA	E4	B3	..	4
SI	<i>Cyathochaeta avenacea</i> Benth.	B2	A2	4
SI	<i>Cyathochaeta clandestina</i> (R. Br.)Benth.	B2	..	4
SC	<i>Gahnia ancistrophylla</i> Benth.	AAA	D3	D4	B4	A2	B3	6
SC	<i>Gahnia australis</i> (Nees)K.L. Wilson	A2	B3	..	4
SI	<i>Gahnia decomposita</i> (R. Br.)Benth.	..P	B3	..	4
SI	<i>Gahnia deusta</i> (R. Br.)Benth.	..P	A3	6
SI	<i>Gahnia drummondii</i> (Steudel)K.L. Wilson	P.P	A3	4
SC	<i>Gahnia lanigera</i> (R. Br.)Benth.	P.P	B3	4
SI	<i>Gahnia</i> sp. A (MAB 4431)	..P	2K	A2	1
AS	<i>Isolepis congrua</i> Nees	..P	A1	C2	6
AS	<i>Isolepis marginata</i> (Thunb.)A. Dietr.	..P	A1	6
SI	<i>Isolepis nodosa</i> (Rottb.)R. Br.	A2	6
SI	<i>Isolepis</i> sp. A (MAB 1515)	...	2B	..	A2	A2	1
SI	<i>Lepidosperma aphyllum</i> R. Br.	C4	..	2,3
SC	<i>Lepidosperma brunonianum</i> Nees	AAA	B3	B3	A4	4
SI	<i>Lepidosperma carphoides</i> F. Muell. ex Benth.	P..	A1	C2	..	6
SI	<i>Lepidosperma drummondii</i> Benth.	AAA	C2	..	C2	..	C3	..	4,5
SI	<i>Lepidosperma gracile</i> R. Br.	..P	A2	4
SC	<i>Lepidosperma leptophyllum</i> Benth.	2,3
SI	<i>Lepidosperma pruinosum</i> Kuek.	P.P	C3	..	A3	..	4,5
SI	<i>Lepidosperma resinum</i> (Nees)Benth.	A.A	B3	4,5
SI	<i>Lepidosperma</i> aff. <i>resinosum</i> (Nees)Benth.	(KRN 5232)	APP	C3	B3	B4	..	4
SI	<i>Lepidosperma squamatum</i> Labill.	B2	..	4
SI	<i>Lepidosperma tuberculatum</i> Nees	PPP	B2	..	4
SI	<i>Lepidosperma viscidum</i> R. Br. var. <i>viscidum</i>	PAP	D4	..	B1	A3	4
SI	<i>Lepidosperma viscidum</i> R. Br. var. <i>flaccidum</i> Kuek.	A2	..	4
SI	<i>Lepidosperma</i> sp. A (MAB 1265)	..P	B2	1
SI	<i>Lepidosperma</i> sp. B (MAB 4371)	..P	3K	A2	1
SI	<i>Lepidosperma</i> sp. C (MAB 1298)	A2	1
SI	<i>Lepidosperma</i> sp. D (MAB 1138)	B2	1
SI	<i>Mesomelaena preissii</i> Nees	AAA	D3	D3	..	4,5
SI	<i>Mesomelaena stygia</i> (R. Br.)Nees ssp. <i>stygia</i>	A.P	D3	D4	B4	2,3
SI	<i>Mesomelaena tetragona</i> (R. Br.)Benth.	..P	B4	C2	4
SI	<i>Mesomelaena</i> sp. A (MAB 3898)	B2	1
SI	<i>Schoenus armeria</i> Boeckler	P.P	C3	..	2
SI	<i>Schoenus asperocarpus</i> F. Muell.	..P	A4	2
SI	<i>Schoenus brevifolius</i> R. Br.	A3	..	4
SI	<i>Schoenus brevisetis</i> (R. Br.)Benth.	AAA	B3	B3	2,3
SI	<i>Schoenus caespititius</i> W. Fitzg.	P.P	3RC	A1	..	2,3
SI	<i>Schoenus curvifolius</i> (R. Br.)Benth.	B2	..	2,3
SI	<i>Schoenus grammatophyllus</i> F. Muell.	B2	..	2,3
SI	<i>Schoenus laevigatus</i> W. Fitzg.	..P	A2	2,3
SI	<i>Schoenus</i> aff. <i>laevigatus</i> W. Fitzg. (KRN 3953)	P..	B2	..	2,5
AS	<i>Schoenus nanus</i> (Nees)Benth.	PPP	A2	..	A2	..	6
SI	<i>Schoenus obtusifolius</i> (Nees)Boeckler	A1	..	2,3
AS	<i>Schoenus odontocarpus</i> F. Muell.	B2	4
SI	<i>Schoenus pleiostemonus</i> F. Muell.	C3	B3	2,3
AS	<i>Schoenus sculptus</i> (Nees)Boeckler	..P	A2	6
SI	<i>Schoenus</i> aff. <i>subfascicularis</i> Kuek. (KRN 4744)	A3	2,3
SI	<i>Schoenus subflavus</i> Kuek.	PPP	B3	..	2,3
SI	<i>Schoenus sublaxus</i> Kuek.	C3	..	2
SI	<i>Schoenus</i> sp. A (MAB 2374)	...	3K	A2	1
SI	<i>Schoenus</i> sp. B (KRN 9787)	A1	..	2
SI	<i>Schoenus</i> sp. C (MAB 3608)	A2	..	1
SC	<i>Tetraria capillarlis</i> (F. Muell.)J. Black	P..	A5	A2	6
SI	<i>Tetraria</i> aff. <i>microcarpa</i> S.T. Blake (KRN 8214)	A2	2,3
SI	<i>Tricostularia compressa</i> Nees	..P	C3	..	2,3
SI	<i>Tricostularia neesii</i> Lehm. var. <i>neesii</i>	P.P	C3	..	2,3
SI	<i>Tricostularia neesii</i> Lehm. var. <i>elatior</i> Benth.	A2	..	2,3
39 RESTIACEAE												
SI	<i>Anarthria gracilis</i> R. Br.	..P	C2	2,3
SI	<i>Anarthria humilis</i> Nees	..A	C2	D3	..	2,3
SC	<i>Anarthria laevis</i> R. Br.	..P	D4	..	2,3
SI	<i>Anarthria prolifera</i> R. Br.	B2	..	2,3
SI	<i>Anarthria scabra</i> R. Br.	..P	B4	..	4
SC	<i>Harperia lateriflora</i> W. Fitzg.	D3	B3	4
SI	<i>Hypolaena exsulca</i> R. Br.	..P	A2	D3	..	4

Appendix 1 (continued). Flora List

LF		FRF	CS	LANDFORM									DIS
				B	D	F1	F2	G	L	M	S		
SI	<i>Hypolaena fastigiata</i> R. Br.	P.P	A2	B2	..	6	
SI	<i>Lepidobolus chaetocephalus</i> F. Muell.	..A	D2	D2	2,3	
SI	<i>Lepidobolus preissianus</i> Nees	AAP	C3	4	
SI	<i>Leptocarpus humilis</i> Gilg	A2	4	
SI	<i>Loxocarya fasciculata</i> (R. Br.)Benth.	PPP	B2	B1	6	
SI	<i>Loxocarya myrioclada</i> Gilg	AAP	C3	B3	2,5	
SI	<i>Loxocarya</i> sp. A (MAB 3967)	A2	1	
SI	<i>Lyginia barbata</i> R. Br.	C3	4	
SI	<i>Restio crispatus</i> R. Br.	D2	A2	..	3	
SI	<i>Restio sphacelatus</i> R. Br.	.PP	D4	C3	B2	6	
40	CENTROLEPIDACEAE												
AS	<i>Aphelia brizula</i> F. Muell.	..P	A2	4	
AS	<i>Brizula muelleri</i> Hieron.	A2	4	
AS	<i>Centrolepis aristata</i> (R. Br.)Roemer & Schultes	A2	6	
AS	<i>Centrolepis glabra</i> (F. Muell. ex Sonder)Hieron.	A2	6	
AS	<i>Centrolepis humillima</i> F. Muell. ex Benth.	PPP	B2	B2	..	4,5	
AS	<i>Centrolepis pilosa</i> Hieron.	.P.	B2	4	
AS	<i>Centrolepis polygyna</i> (R. Br.)Hieron.	P.P	D3	6	
AS	<i>Centrolepis strigosa</i> (R. Br.)Roemer & Schultes	A2	6	
52	JUNCACEAE												
AS	* <i>Juncus bufonius</i> L.	..P	B2	7	
SI	<i>Juncus pallidus</i> R. Br.	A3	6	
54C	DASYPOGONACEAE												
DS	<i>Calectasia cyanea</i> R. Br.	P.A	B1	4	
SL	<i>Chamaeeros fimbriata</i> (F. Muell.)Benth.	A2	2,5	
SL	<i>Lomandra collina</i> (R. Br.)Ewart	PPP	A2	..	C2	6	
SL	<i>Lomandra</i> aff. <i>effusa</i> (Lindley)Ewart (MAB 4036)	..P	A1	..	1	
SL	<i>Lomandra hastilis</i> (R. Br.)Ewart	..P	A1	4	
SL	<i>Lomandra micrantha</i> (Endl.)Ewart ssp. <i>micrantha</i>	P.P	B2	6	
SL	<i>Lomandra micrantha</i> (Endl.)Ewart ssp. <i>teretifolia</i>	6	
SL	<i>Lomandra mucronata</i> (R. Br.)A. Lee	..P	A1	4	
SL	<i>Lomandra nigricans</i> T.D. MacFarlane	A1	4	
54D	XANTHORRHOACEAE												
SL	<i>Xanthorrhoea platyphylla</i> D.J. Bedford	P..	C2	4	
54E	PHORMIACEAE												
SL	<i>Dianella revoluta</i> R. Br.	APA	C1	C2	B3	C2	..	A1	..	6	
SS	<i>Stypantra grandiflora</i> Lindley	..P	C3	4	
DS	<i>Stypantra imbricata</i> R. Br.	PAP	D3	4	
54F	ANTHERICACEAE												
SL	<i>Agrostocrinum scabrum</i> (R. Br.)Baillon	.PA	B1	4	
SL	<i>Borya constricta</i> D.M. Churchill	PAA	E4	4	
SL	<i>Borya</i> sp. A (MAB 1611)	.P.	B2	2	
AB	<i>Chamaescilla corymbosa</i> (R. Br.)F. Muell. ex Benth.	..P	B3	B2	..	6	
AB	<i>Chamaescilla spiralis</i> (Endl.)F. Muell. ex Benth.	A2	..	A2	4	
SL	<i>Johnsonia acaulis</i> Endl.	A2	..	A1	4	
DS	<i>Laxmannia brachyphylla</i> F. Muell. ex Benth.	..P	A2	3	
DS	<i>Laxmannia minor</i> R. Br.	..P	A1	4	
DS	<i>Laxmannia paleacea</i> F. Muell.	..P	B2	4	
DS	<i>Laxmannia ramosa</i> Lindley	B1	4	
DS	<i>Laxmannia sessiliflora</i> Decne.	A1	6	
DS	<i>Laxmannia squarrosa</i> Lindley	A1	..	B1	4	
AB	<i>Stawellia gymnocephala</i> Diels	A1	3	
AB	<i>Thysanotus</i> aff. <i>baueri</i> R. Br. (KRN 8002)	A1	2,5	
AB	<i>Thysanotus manglesianus</i> Kunth	..P	B3	4,5	
AB	<i>Thysanotus parviflorus</i> N.H. Brittan	..P	D2	B2	..	3	
AB	<i>Thysanotus patersonii</i> R. Br. ssp. <i>patersonii</i>	PAP	C2	..	B1	6	
AB	<i>Thysanotus sparteus</i> R. Br.	A1	4,5	
54G	ASPHODELACEAE												
AB	* <i>Asphodelus fistulosus</i> L.	..P	A2	7	
AS	<i>Bulbine semibarbata</i> (R. Br.)Haw.	..PP	C2	..	D3	B3	6
54J	COLCHICACEAE												
AB	<i>Wurmbea cernua</i> T.D. MacFarlane	..P	C2	3	
AB	<i>Wurmbea sinora</i> T.D. MacFarlane	..P	A2	3	
AB	<i>Wurmbea tenella</i> (Endl.)Benth.	..PP	C2	4	
55	HAEMODORACEAE												
AB	<i>Anigozanthos humilis</i> Lindley	A2	4	
AB	<i>Anigozanthos rufus</i> Labill.	P.P	B2	2,3	

Appendix 1 (continued). Flora List

LF	RES	LANDFORM										DLS
		FPE	CS	B	D	F1	F2	G	L	M	S	
5L	Conostylis androstemma F. Muell. ssp. argentea J.W. Green	A..	B2	B2	4
SL	Conostylis bealiana F. Muell.	..P	B2	..	C4	3
SL	Conostylis breviscapa R. Br.	A2	4
SL	Conostylis phathyrantha Diels	..	3H	C2	2,3
MP	Conostylis seorsiflora F. Muell.	..P	B1	3
SL	Conostylis vaginata Endl.	A2	3
SL	Conostylis sp. A (KRN 6572)	..	2E	C2	1
56A	HYPOXIDACEAE											
AB	Hypoxis sp. A (KRN 9759)	..	2V	B4	1
60	IRIDACEAE											
SL	Patersonia juncea Lindley	..P	B1	4
SL	Patersonia lanata R. Br.	B1	3
SL	Patersonia occidentalis R. Br.	..P	B1	6
SL	Patersonia rudis Endl. ssp. rudis	A1	4
66	ORCHIDACEAE											
AB	Acianthus reniformis (R. Br.)Schltr.	B2	6
AB	Caladenia caerulea R. Br.	..P	B2	4
AB	Caladenia deformis R. Br.	..P	C2	6
AB	Caladenia dilatata R. Br. var. falcata Nicholls	..P	B2	6
AB	Caladenia discoidea Lindley	C2	4
AB	Caladenia douthchae O. Sarg.	..P	D2	B2	4
AB	Caladenia filamentosa R. Br. var. denticulata (Lindley) H.G. Reichb.	B2	4
AB	Caladenia filamentosa R. Br. var. tentaculata R.S. Rogers	B2	6
AB	Caladenia flava R. Br.	..P	B2	4
AB	Caladenia gemmata Lindley	..P	B2	4
AB	Caladenia hirta Lindley	B2	4
AB	Caladenia longicauda Lindley	..P	B2	4
AB	Caladenia reptans Lindley	B2	4
AB	Caladenia roei Benth.	..P	B2	4
AB	Caladenia saccharata H.G. Reichb.	B2	A2	4,5
AB	Caladenia sigmoidea R.S. Rogers	..P	A2	2,5
AB	Caladenia sp. A (MAB 3291)	B2	2
AB	Caladenia sp. B (MAB 3208)	..P	A2	2
AB	Diuris laxiflora Lindley	..P	A2	..	Al	4,5
AB	Diuris longifolia R. Br.	..PA	B2	6
AB	Elythranthera brunonis (Endl.)A.S. George	..P	B2	..	Al	4
AB	Eriochilus dilatatus Lindley	..P	C2	4
AB	Eriochilus scaber Lindley	..P	B2	4
AB	Leporella fimbriata (Lindley)A.S. George	..P	B2	6
AB	Lyperanthus nigricans R. Br.	..P	B2	..	Al	6
AB	Lyperanthus serratus Lindley	Al	4
AB	Microtis unifolia (G. Forster)H.G. Reichb.	..P	A3	6
AB	Paracaleana nigrita (Lindley)Blaxell ssp. nov. (MAB 3745)	B2	B1	..	3
AB	Prasophyllum fimbria H.G. Reichb.	Al	4
AB	Prasophyllum macrostachyum R. Br. var. rigens (H.G. Reichb.)A.S. George	C2	2,5
AB	Prasophyllum nigricans R. Br.	P.P	B2	6
AB	Prasophyllum ovale Lindley var. trigloch H.G. Reichb.	B2	4
AB	Pterostylis mutica R. Br.	..P	B2	6
AB	Pterostylis nana R. Br.	P.P	B2	6
AB	Pterostylis plumosa L. Cady	B2	6
AB	Pterostylis recurva Benth.	..P	B2	4
AB	Pterostylis aff. rufa R. Br. A (MAB 4463a)	..P	C2	2
AB	Pterostylis aff. rufa R. Br. B (MAB 3592)	B2	1
AB	Pterostylis aff. rufa R. Br. C (MAB 3554)	B2	1
AB	Pterostylis sargentii C.R.P. Andrews	P P	B2	..	Al	4
AB	Pterostylis vittata Lindley var. vittata	PPP	B2	6
AB	Thelymitra antennifera (Lindley)J.D. Hook.	B1	6
AB	Thelymitra campanulata Lindley	B2	4
AB	Thelymitra canaliculata R. Br.	B2	6
AB	Thelymitra crinita Lindley	..P	B1	4
AB	Thelymitra fuscolutea R. Br. var. fuscolutea	B1	6
AB	Thelymitra nuda R. Br.	PPA	B1	C1	..	6
AB	Thelymitra pauciflora R. Br.	B1	6
	DICOTYLEDONAE											
70	CASUARINACEAE											
SS	Allocasuarina acuarina (F. Muell.)L. Johnson	A2	2,3
LS	Allocasuarina campestris (Diels)L. Johnson ssp. campestris	AAA	D5	..	C4	4

Appendix 1 (continued). Flora List

LF	SUS	SUS	LANDFORM									DLS
			B	D	F1	F2	G	L	M	S		
TS	Allocasuarina corniculata (F. Muell.)L. Johnson	P.P	A4	4,5
TS	Allocasuarina helmsii (Ewart & M. Gordon)			
	L. Johnson	.PP	B2		6
DT	Allocasuarina huegeliana (Miq.)L. Johnson	PAP	D4		4,5
SS	Allocasuarina humilis (Otto & Dietr.)L. Johnson	APA	C3	C3		4,5
DS	Allocasuarina microstachya (Miq.)L. Johnson	APP	C2	4
MS	Allocasuarina scleroclada (L. Johnson)L. Johnson	P..	A1		2,5
SS	Allocasuarina thuyoides (Miq.)L. Johnson	APP	B2	D3	4
TS	Allocasuarina trichodon (Miq.)L. Johnson	.P.P	A2	..	4
DT	Casuarina cristata Miq. ssp. pauper (F. Muell. ex Miq.)L. Johnson	.P.P	B2		6
90 PROTEACEAE												
DS	Adenanthos barbigerus Lindley	.P.P	B2	..	4
MS	Adenanthos cuneatus Labill.	P.A	B3	..	2,3
DS	Adenanthos glabrescens E.C. Nelson ssp. glabrescens	A2	B2	..	2,5
MS	Adenanthos ileticus E.C. Nelson	B2		1
DS	Banksia blechnifolia F. Muell.	.P.P	3VC	B2		2
TS	Banksia elderiana F. Muell. & Tate	AAP	C4	C5		4,5
TS	Banksia media R. Br.	AAA	D5	C4	A2	D5	2,3
SS	Banksia nutans R. Br. var. nutans	B1	..	3
DS	Banksia petiolaris F. Muell.	.P.P	D3	C3	2,3
TS	Banksia pilostylis C. Gardner	.P.P	3VC	B4	D4	C3	2,3
SS	Banksia pulchella R. Br.	B3	..	2,3
DS	Banksia repens Labill.	.P.P	D3	..	2,3
TS	Banksia speciosa R. Br.	.P.P	B4	D5	3
SS	Banksia violacea C. Gardner	P.P.P	B2	4
DS	Conospermum distichum R. Br.	P.A	B3		4
DS	Conospermum floribundum Benth.	B3	..	4
SS	Conospermum leianthum E. Pritzel	P.P.P	B3	B2	..	2,3
MS	Conospermum teretifolium R. Br.	P.P.P	B1	..	2,3
DS	Dryandra armata R. Br.	.P.P	C4	B5	..	4
SS	Dryandra cirsioides Meissner	.P.P	C4		4
DS	Dryandra aff. cirsioides Meissner (MAB 1328)	A.P	B3		2
MS	Dryandra cuneata R. Br.	.P.P	B3		4
DS	Dryandra nivea (Labill.)R. Br.	.P.P	D3	A1	..	4
DS	Dryandra obtusa R. Br.	D3	..	3
DS	Dryandra aff. pteridifolia R. Br. (MAB 4593)	.P.P	3VC	A2	A2	..	2
DS	Dryandra tenuifolia R. Br.	.P.P	C3	..	2,3
SS	Franklandia fucifolia R. Br.	.P.P	C2		4
DS	Grevillea acuaria F. Muell. ex Benth.	AAP	C3	..	4,5
SS	Grevillea aneura McGillivray	P..	3VC	B2	B3	..	2
SS	Grevillea beardiana McGillivray	A2		2
TS	Grevillea caghiana McGillivray	AP.	B3		4,5
MS	Grevillea coccinea Meissner	PP.	A2	B2	..	2,3
MS	Grevillea didymobotrya Meissner ssp. didymobotrya	P.P	B2	..	4,5
DS	Grevillea disjuncta F. Muell. ssp. disjuncta	.P.P	A2	..	2
MS	Grevillea endlicheriana Meissner	.P.	B3	..	4,5
TS	Grevillea eriostachya Lindley ssp. excelsior (Diels)McGillivray	PPP	D4	4,5
MS	Grevillea extorris S. Moore	P.P.P	A1		4,5
DS	Grevillea haplantha F. Muell. ex Benth.	P..	A2	..	2,3
SS	Grevillea huegelii Meissner	AAA	B3	D3		6
MS	Grevillea integrifolia (Endl.)Meissner ssp. incrassata (Diels)McGillivray	PP.	B4	..	2
MS	Grevillea integrifolia (Endl.)Meissner ssp. integrifolia	A2	..	4,5
MS	Grevillea integrifolia (Endl.)Meissner ssp. shuttleworthiana (Meissner)McGillivray	.P.P	A2	..	4,5
DS	Grevillea nudiflora Meissner	.P.P	A1	..	2,3
MS	Grevillea oligantha F. Muell.	.P.P	C2		2,5
MS	Grevillea oncogyne Diels	APP	B2		2,5
LS	Grevillea paniculata Meissner	C4		2,3
DS	Grevillea patentifolia F. Muell.	A2	..	2,3
MS	Grevillea pauciflora R. Br. ssp. pauciflora	AAA	B1	C2	C3	B2	C3	6
SS	Grevillea pectinata R. Br.	.AP	C2	C3	C3	..	2,3
MS	Grevillea aff. pectinata R. Br. (KRN 9784)	C3	..	2
MS	Grevillea plurijuga F. Muell.	.P.P	B2		2
MS	Grevillea teretifolia Meissner	PPP	C2	..	4,5
TS	Grevillea tetragonoloba Meissner	B2		2
DS	Grevillea aff. treueriana F. Muell. (MAB 2305)	C3		1
TS	Grevillea wittweri McGillivray	B2		2
TS	Hakea adnata R. Br.	.P.P	C3	A3	C1	..	2,3
DT	Hakea arida Diels	B2		2,5
LS	Hakea brooksiana F. Muell.	D3	..	2,3
MS	Hakea cinerea R. Br.	.P.P	B3	D3	..	2,3
MS	Hakea clavata Labill.	.P.P	D4	..	4,5
SS	Hakea commutata F. Muell.	P.P	B3	D4	B2	..	4

Appendix 1 (continued). Flora List

LF		RES		LANDFORM								Dis		
		FPS	CS	B	D	F1	F2	G	L	M	S			
MS	<i>Hakea corymbosa</i> R. Br.	PPP	D4	S3	C4	2,3		
LS	<i>Hakea crassifolia</i> Meissner	P..	B3	B3	4		
SS	<i>Hakea aff. falcata</i> R. Br. (KRN 5966)	A..	B5	4,5		
DT	<i>Hakea laurina</i> R. Br.	PPA	B3	D5	B2	..	4		
MS	<i>Hakea lissocarpha</i> R. Br.	APP	B2	A2	4		
DS	<i>Hakea marginata</i> R. Br.	P..P	B2	4		
DT	<i>Hakea multilinea</i> Meissner	PP.	B2	4		
TS	<i>Hakea nitida</i> R. Br.	.P.	D2	B1	C2	2,5		
TS	<i>Hakea pandanica</i> R. Br.	..P	B4	C3	3		
TS	<i>Hakea prostrata</i> R. Br.	..P	B1	..	4		
TS	<i>Hakea rubriflora</i> Lamont	..P	B2	..	2,3		
MS	<i>Hakea ruscifolia</i> Labill.	A3	..	4		
LS	<i>Hakea subsulcata</i> Meissner	A2	..	4		
MS	<i>Hakea trifurcata</i> (Smith)R. Br.	..PP	C4	C3	D4	4		
MS	<i>Hakea varia</i> R. Br.	..P	B3	B3	..	4		
LS	<i>Hakea sp. A</i> (KRN 9766)	..	IE	A2	1		
DS	<i>Isopogon alpicornis</i> Diels	..P	A2	2		
DS	<i>Isopogon attenuatus</i> R. Br.	P..P	A2	D2	..	4		
SS	<i>Isopogon buxifolius</i> R. Br.	P..A	Y5	4		
DS	<i>Isopogon formosus</i> R. Br.	..A	A2	..	2,3		
MS	<i>Isopogon polycephalus</i> R. Br.	P..P	B2	..	2,3		
DS	<i>Isopogon teretifolius</i> R. Br.	A..A	B2	..	4		
DS	<i>Isopogon tridens</i> F. Muell.	..P	3V	E4	2		
MS	<i>Isopogon trilobus</i> R. Br.	A2	C3	B3	2,3		
MS	<i>Isopogon tripartitus</i> R. Br.	A2	C3	..	3		
TS	<i>Lambertia inermis</i> R. Br.	..A	C4	D4	..	2,3		
DS	<i>Persoonia striata</i> R. Br.	..P	C2	..	4,5		
LS	<i>Persoonia teretifolia</i> R. Br.	..P	B2	..	4,5		
DS	<i>Persoonia tortifolia</i> Meissner	..	2K	B2	4		
DS	<i>Persoonia sp. A</i> (MAB 3392)	..PP	3RC	B2	2		
SS	<i>Petrophile divaricata</i> R. Br.	C2	4		
SS	<i>Petrophile ericifolia</i> R. Br. var. <i>ericifolia</i>	..P	A2	B2	4		
SS	<i>Petrophile aff. ericifolia</i> R. Br. A (MAB 1325)	A2	1		
SS	<i>Petrophile aff. ericifolia</i> R. Br. B (MAB 2910)	A2	1		
SS	<i>Petrophile fastigiata</i> R. Br.	..P	B2	C2	3		
MS	<i>Petrophile heterophylla</i> Lindley	C2	B2	4		
DS	<i>Petrophile phyllicoides</i> R. Br.	2A	4		
DS	<i>Petrophile seminuda</i> Lindley	P..P	C3	4		
SS	<i>Petrophile squamata</i> R. Br.	..A	C2	C3	..	4		
DS	<i>Petrophile teretifolia</i> R. Br.	..A	D3	C3	2,3		
DS	<i>Stirlingia tenuifolia</i> (R. Br.)Steudel	PAA	D3	D3	..	2,3		
DS	<i>Synaphea favosa</i> R. Br.	..A	D4	C3	..	4		
DS	<i>Synaphea polymorpha</i> R. Br.	C3	4		
DS	<i>Synaphea sp. A</i> (MAB 1778)	B2	1		
DS	<i>Synaphea sp. B</i> (KRN 3761)	A7	..	4		
92 SANTALACEAE														
SS	<i>Choretrum glomeratum</i> R. Br. var. <i>chrysanthum</i> (F. Muell.)Benth.	A1	6		
SS	<i>Choretrum glomeratum</i> R. Br. var. <i>glomeratum</i>	..P	B2	B2	6		
MS	<i>Exocarpos aphyllus</i> R. Br.	..P	C3	A1	6		
TS	<i>Exocarpos sparteus</i> R. Br.	..P	C4	6		
SS	<i>Leptomeria pachyclada</i> Diels	..P	B1	C3	A2	..	4		
MS	<i>Leptomeria preissiana</i> (Miq.)A. DC.	APA	B1	C3	B2	B2	6		
SS	<i>Leptomeria scrobiculata</i> R. Br.	B1	C2	B2	B3	A2	4	
DS	<i>Leptomeria spinosa</i> (Miq.)A. DC.	..P	D3	A2	C2	4		
DT	<i>Santalum acuminatum</i> (R. Br.)A. DC.	A2	B2	..	6	
DT	<i>Santalum murrayanum</i> (Mitch.)C. Gardner	AAA	A2	B2	D3	..	C3	..	6
95 OLACACEAE														
DS	<i>Olax benthamiana</i> Miq.	PPP	C2	B3	..	4		
DS	<i>Olax sp. A</i> (KRN 9799)	A1	..	1		
97 LORANTHACEAE														
DT	<i>Nuytsia floribunda</i> (Labill.)R. Br. ex Fenzl	..P	C2	B2	C3	4		
103 POLYGONACEAE														
DS	<i>Muehlenbeckia adpressa</i> (Labill.)Meissner	PPP	B2	B2	C2	..	6	
HP	* <i>Rumex acetosella</i> L.	A2	7		
105 CHENOPODIACEAE														
DS	<i>Atriplex lindleyi</i> Moq. ssp. <i>lindleyi</i>	B2	C3	..	6	
DS	<i>Atriplex lindleyi</i> Moq. ssp. <i>inflata</i> (F. Muell.) Paul G. Wilson	B2	C3	..	6	
DS	<i>Atriplex paludosa</i> R. Br.	..P	C3	..	6		
DS	<i>Atriplex pumilio</i> R. Br.	C3	..	6		
AS	<i>Atriplex spongiosa</i> F. Muell.	PPP	A2	C3	..	6	
DS	<i>Atriplex vesicaria</i> Heward ex Benth. ssp. <i>variabilis</i> Parr-Smith	AAA	B2	C3	..	B2	..	6

Appendix 1 (continued). Flora List

LP	REFS	LANDFORM								DIF		
		PPS	CM	B	D	F1	F2	G	L		M	S
MP	Chenopodium desertorum (J. Black)J. Black ssp. microphyllum Paul G. Wilson	B3	6
DS	Enchylaena lanata Paul G. Wilson	PPP	B2	..	B2	..	B3	4
DS	Enchylaena tomentosa R. Br. var. tomentosa	FAA	C2	..	B2	B2	D4	6
AS	Eriochiton sclerolaenoides (F. Muell.)F. Muell. ex A.J. Scott	A1	4
DS	Halosarcia halocnemoides (Nees)Paul G. Wilson ssp. caudata Paul G. Wilson	D5	4,5
DS	Halosarcia halocnemoides (Nees)Paul G. Wilson ssp. halocnemoides	D5	6
SS	Halosarcia indica (Willd.)Paul G. Wilson bidens Paul G. Wilson	D4	6
DS	Halosarcia lepidosperma Paul G. Wilson	P.P	C4	6
DS	Halosarcia lylei (Ewart & J. White)Paul G. Wilson	P.A	C4	E5	6
DS	Halosarcia pergranulata (J. Black)Paul G. Wilson ssp. pergranulata	..P	D4	6
DS	Halosarcia pterygosperma (J. Black)Paul G. Wilson ssp. pterygosperma	P.P	D5	6
DS	Halosarcia syncarpa Paul G. Wilson	.AP	E5	6
DS	Maireana amoena (Diels)Paul G. Wilson	.AP	D4	5
DS	Maireana enchylaenoides (F. Muell.)Paul G. Wilson	P.P	D4	6
DS	Maireana erioclada (Benth.)Paul G. Wilson	..P	C3	6
DS	Maireana oppositifolia (F. Muell.)Paul G. Wilson	..P	C4	6
DS	Maireana trichoptera (J. Black)Paul G. Wilson	C4	6
DS	Rhagodia crassifolia R. Br.	..P	C3	6
DS	Rhagodia drummondii Moq.	PPP	C3	6
MS	Rhagodia preissii Moq. ssp. preissii	PAP	B2	C3	6
DS	Sarcocornia quinqueflora (Bunge ex Ung.-Sternb.) A.J. Scott	..P	D3	6
DS	Sclerolaena patenticuspis (Anderson)Ulbr.	..P	D3	6
DS	Sclerolaena uniflora R. Br.	..A	D4	6
DS	Sclerostegia moniliformis Paul G. Wilson	..P	D3	4
DS	Suaeda australis (R. Br.)Moq.	A2	6
DS	Threlkeldia diffusa R. Br.	..A	E4	6
106	AMARANTHACEAE											
HP	Hemichroa diandra R. Br.	B2	6
HP	Ptilotus holosericeus (Moq.)F. Muell.	P.	B1	2,5
HP	Ptilotus spathulatus (R. Br.)Poiret	A2	A2	..	6
HP	Ptilotus sp. A (MAB 4484B)	..P	B3	1
108	GYROSTEMONACEAE											
DS	Cypselocarpus haloragoides (F. Muell. ex Benth.) F. Muell.	...	3VC	B2	3
SS	Gyrostemon ditrigynus A.S. George	...	2E	A3	2
MP	Gyrostemon prostratus A.S. George	...	3E	A2	2
SS	Gyrostemon ramulosus Desf.	C3	6
MS	Gyrostemon sheathii W. Fitzg.	A1	2,3
110	AIZOACEAE											
MP	*Carpobrotus edulis (L.)L. Bolus	A3	7
MP	Carpobrotus modestus S.T. Blake	PAP	C2	C3	D1	..	6
MP	Carpobrotus rossii (Haw.)Schwantes	PPP	B1	C2	..	6
MP	Disphyma crassifolium (L.)L. Bolus	..AA	D4	6
DS	Gunniopsis glabra (Ewart)C. Gardner	..P	A1	5
DS	Gunniopsis intermedia Diels	..P	A1	5
111	PORTULACACEAE											
AS	Calandrinia calyptrata J.D. Hook.	PPP	D2	C2	..	6
AS	Calandrinia eremaea Ewart	..P	C2	6
AS	Calandrinia polyandra Benth.	..P	C2	6
AS	Calandrinia sp. A (MAB 2265)	A2	1
113	CARYOPHYLLACEAE											
AS	*Spergularia diandra Heldr. & Sart. ex Heldr.	C2	7
AS	*Spergularia rubra (L.)J.S. & C. Presl	..P	C2	7
131	LAURACEAE											
PC	Cassytha aurea J.Z. Weber	B2	4
PC	Cassytha glabella R. Br.	PAP	B1	..	B2 B2	6
PC	Cassytha melantha R. Br.	PAA	C1	D3	C2	..	B2	..	6
PC	Cassytha micrantha Meissner	...	3VC	A1	..	2,3
PC	Cassytha racemosa Nees	..P	B1	B1	C2 B2	4
138	BRASSICACEAE											
AS	*Brassica tournefortii Gouan	B2	7
AS	*Brassica sp. A (MAB 2277)	A2	7
AS	*Hymenolobus procumbens (L.)Nutt. ex Schinz & Thell.	B2	7

Appendix 1 (continued). Flora List

LF		KEE		LANDFORM								DIS	
		PPS	CS	B	D	F1	F2	G	L	M	S		
HP	<i>Lepidium rotundum</i> (Desv.) DC.	P..	B2	6
AS	<i>Menkea australis</i> Lehm.	.P.	B2	6
AS	* <i>Sisymbrium irio</i> L.	A2	7
AS	<i>Stenopetalum robustum</i> Endl.	B2	4
143	DROSERACEAE												
AB	<i>Drosera bulbosa</i> Hook.	..P	B2	4
AS	<i>Drosera glanduligera</i> Lehm.	PPP	C2	6
AB	<i>Drosera huegelii</i> Endl.	A1	4
AB	<i>Drosera leucoblata</i> Benth.	B2	4
AB	<i>Drosera macrantha</i> Endl.	APP	D2	..	D3	..	D3	..	4
AB	<i>Drosera menziesii</i> R. Br. ssp. <i>menziesii</i>	PPP	B2	B1	4
AB	<i>Drosera neesii</i> Lehm. ssp. <i>neesii</i>	A2	4
RP	<i>Drosera paleacea</i> DC.	..P	H2	4
RP	<i>Drosera pycnoblata</i> Diels	PP.	B2	4
AB	<i>Drosera ramellosa</i> Lehm.	..P	B2	4
AB	<i>Drosera subhirtella</i> Planchon ssp. <i>moorei</i> (Diels) N.G. Marchant	A2	2,5
AB	<i>Drosera zonaria</i> Planchon	P.P	B2	4
AB	<i>Drosera</i> sp. A. (MAB 3703)	.P.	3R	B2	2,5
149	CRASSULACEAE												
AS	<i>Crassula colorata</i> (Nees) Ostenf. var. <i>colorata</i>	..A	D3	D2	D3	6
AS	<i>Crassula decumbens</i> Thunb. var. <i>decumbens</i>	A2	B3	6
AS	<i>Crassula exserta</i> (Reader) Ostenf.	AAA	D2	6
HY	* <i>Crassula natans</i> Thunb. var. <i>minus</i> (Ecklon & Zeyher) Rowley	B3	7
AS	<i>Crassula pedicellosa</i> (F. Muell.) Ostenf.	.PP	B2	6
AS	<i>Crassula sieberiana</i> (Schultes & J.H. Schultes) Druce ssp. <i>tetramera</i> Toelken	B2	6
152	PITOSPORAEEAE												
CL	<i>Billardiera bicolor</i> (Putterl.) E.M. Bennett	.PP	B1	4
CL	<i>Billardiera coriacea</i> Benth.	PPP	B1	..	B1	4
CL	<i>Billardiera lehmanniana</i> F. Muell.	A1	4,5
SS	<i>Billardiera mollis</i> E.M. Bennett	...	2V	A1	A1	2,3
SS	<i>Cheiranthra filifolia</i> Turcz.	.PP	A1	4,5
DT	<i>Pittosporum phylliraeoides</i> DC. var. <i>microcarpa</i> S. Moore	PAP	A2	4,5
CL	<i>Sollya heterophylla</i> Lindley	PPP	B1	4
163	MIMOSACEAE												
DS	<i>Acacia acanthoclada</i> F. Muell.	P.P	B1	6
DT	<i>Acacia acuminata</i> Benth.	.AP	B4	B3	..	D5	4,5
MS	<i>Acacia ancistrophylla</i> C.R.P. Andrews	B3	A2	4
DS	<i>Acacia andrewsii</i> W. Fitzg.	PPP	B1	B2	4
TS	<i>Acacia assimilis</i> S. Moore	PAA	D5	4,5
DS	<i>Acacia</i> aff. <i>bidentata</i> Benth. A (MAB 1688)	P.P	A2	4
DS	<i>Acacia</i> aff. <i>bidentata</i> Benth. B (MAB 2496)	P.P	2VC	B1	1
SS	<i>Acacia binata</i> Maslin	P..	A2	2,3
DS	<i>Acacia brachyphylla</i> Benth.	A2	4
SS	<i>Acacia camptoclada</i> C.R.P. Andrews	AAA	B3	B3	2,5
DS	<i>Acacia</i> aff. <i>camptoclada</i> C.R.P. Andrews (MAB 1879)	A1	1
TS	<i>Acacia chrysellia</i> Maiden & Blakely	P..	A2	4,5
DS	<i>Acacia chrysocephala</i> Maslin	..P	A1	4
DS	<i>Acacia cochlearis</i> (Labill.) H.L. Wendl.	B1	4
SS	<i>Acacia</i> aff. <i>cochlearis</i> (Labill.) H.L. Wendl. (KRN 9764)	...	2K	B2	3
DS	<i>Acacia</i> aff. <i>congesta</i> Benth. (MAB 1991)	A2	1
DT	<i>Acacia conniana</i> Maslin	B4	3
DS	<i>Acacia congesta</i> Benth.	A2	2,3
TS	<i>Acacia crassiuscula</i> Wendl.	4
DE	<i>Acacia crassuloides</i> Maslin	PAP	3VC	B2	2,3
DS	<i>Acacia curvata</i> Maslin	..P	A3	2,3
TS	<i>Acacia cyclops</i> Cunn. ex Don	..P	B2	6
DS	<i>Acacia delphina</i> Maslin	B3	C3	2,3
DS	<i>Acacia densiflora</i> Morrison	P..	B3	4,5
DS	<i>Acacia dermatophylla</i> Benth.	B2	1
DS	<i>Acacia</i> aff. <i>dermatophylla</i> Benth. (MAB 1877)	P.P	C3	5
DS	<i>Acacia diaphylloidea</i> Maslin	A.P	A3	2,3
MS	<i>Acacia dielsii</i> E. Pritzel	..P	B3	4
DS	<i>Acacia erinacea</i> Benth.	AAA	C4	B2	6
DS	<i>Acacia excentrica</i> Maiden & Blakely	...	3K	A2	2,5
DS	<i>Acacia ferocior</i> Maiden	..P	B4	4
SS	<i>Acacia flavopila</i> A.S. George	A2	4
TS	<i>Acacia fragilis</i> Maiden & Blakely	..P	A2	4,5
LS	<i>Acacia</i> aff. <i>fragilis</i> Maiden & Blakely (KRN 7972)	A1	2
SS	<i>Acacia glaucoptera</i> Benth.	..P	C3	4
SS	<i>Acacia gonophylla</i> Benth. var. <i>gonophylla</i>	..P	B2	A1	2,3

Appendix 1 (continued). Flora List

LF		RES	LANDFORM										DIS		
			FPS	CG	B	D	F1	F2	G	L	M	S			
DS	Acacia aff. gonophylla Benth. (MAB 1156)	..P	A1	1
DS	Acacia gonophylla Benth. var. crassifolia Benth.	..A.	B2	B2	2,3
LS	Acacia hakeoides Cunn. ex Benth.	..P.	A2	4,5
MS	Acacia hemiteles Benth.	..P.	B3	4,5
SS	Acacia ixiophylla Benth.	..PA.	B3	4,5
DS	Acacia lachnophylla F. Muell.	..PAP	B3	4,5
DS	Acacia laticarpa Meissner	..P	B2	4
DT	Acacia lasiocalyx C.R.P. Andrews	..PAP	E4	B2	..	4,5
DS	Acacia lasiocarpa Benth. var. lasiocarpa	..P	A2	4
DS	Acacia lasiocarpa Benth. var. bracteolata Maslin	..P	A2	4
MS	Acacia latipes Benth.	..A.	A3	..	4
LS	Acacia leptoneura Benth.	..A.	B2	B3	..	4
SS	Acacia leptospermoides Benth. var. leptospermoides	..P.	A1	4
LS	Acacia ligulata Cunn. ex Benth.	..PP.	B2	6
MS	Acacia aff. lineolata Benth. (KRN 5421)	..PA.	A2	..	C3	H1	..	2,3
MP	Acacia maxwellii Maiden & Blakely	..P.	B2	2,3
SS	Acacia merrallii F. Muell. x A. sp. (MAB 2231)	..P.	A1	1
MS	Acacia multispicata Benth.	..AAA	B3	4,5
MS	Acacia aff. multispicata Benth. (MAB 3785)	..P.	A3	1
MS	Acacia myrtifolia (Smith)Willd.	..P	C5	A1	6
DS	Acacia nitidula Benth. (MAB 2229)	..P.	C4	B4	2,3
DS	Acacia nodiflora Benth. var. ferox E. Pritzel	..PAP	B4	4
LS	Acacia nyssophylla F. Muell.	..AAA	B2	6
DS	Acacia pachypoda Maslin	..PAP	B3	2,5
DS	Acacia phlebopetala Maslin var. phlebopetala	..P.	A1	2,3
DS	Acacia pilosa Benth.	..P.	A2	2,3
DS	Acacia pritzeliana C. Gardner	..P	3RC	A2	C3	A2	..	2
MS	Acacia pulchella R. Br. var glaberrima Meissner	..P.	B1	B1	4
TS	Acacia saligna (Labill.)H.L. Wendl.	..AP	C2	4
SS	Acacia saxatilis S. Moore	..PAP	B4	4
SS	Acacia aff. saxatilis S. Moore (MAB 2242)	..P.	A2	1
MS	Acacia sessilis M. Moore & Blakely	..P.	B3	4
DS	Acacia sorophylla E. Pritzel	..P	3VC	B3	2
SS	Acacia sphacelatus Benth.	..P.	A2	..	4
SS	Acacia spinosissima Benth.	..P.P	A1	..	4
MS	Acacia subcaerulea Lindley	..P	A4	2,3
DS	Acacia sulcata R. Br. var. sulcata	..P.	A3	2,3
SS	Acacia sulcata R. Br. var. platyphylla Maiden & Blakely	..A.	B3	..	A2	4
MS	Acacia triptycha F. Muell. ex Benth.	..P.	A3	4
SS	Acacia uncinella Benth.	..P.	A3	4,5
DS	Acacia unifissilis Court	..P.P	B2	4
DS	Acacia varia Maslin var. parviflora (Benth.)Maslin	..P.	A1	4
MS	Acacia viscidifolia Maiden & Blakely	..P.P	B3	2,3
LS	Acacia warramaba Maslin	..P.	A2	2,5
DS	Acacia sp. A (MAB 4591)	..P	B2	4
DS	Acacia sp. B (MAB 3858)	..P	2V	B2	2
DS	Acacia sp. C (MAB 3540)	..P	C2	4
DS	Acacia sp. D (MAB 2319)	..P	B2	4
DS	Acacia sp. E (MAB 4008)	..P	3VC	A2	..	4
SS	Acacia sp. F (MAB 2575)	..P.	B2	2,5
DS	Acacia sp. G (MAB 4562)	..P	2
DS	Acacia sp. H (MAB 1192)	..P.	A1	..	2
DS	Acacia sp. I (MAB 4287)	..P	2VC	C3	C2	1
DS	Acacia sp. J (MAB 2157)	..P	C3	4
DS	Acacia sp. K (MAB 1916)	..P.	3R	C3	1
DS	Acacia sp. L (MAB 3528)	..P.	A1	1
DS	Acacia sp. M (MAB 1575)	..P.	A2	1
SS	Acacia sp. N (MAB 2740)	..P.	1V	A2	1
DS	Acacia sp. O (KRN 9681)	..P.	2E	A1	1
DS	Acacia sp. P (KRN 9714)	..P.	3E	A4	..	1
DS	Acacia sp. Q (KRN 9764)	..P.	A2	..	1
DS	Acacia sp. R (KRN 9810)	..P.	1E	A3	1
DS	Acacia sp. S (KRN 7952)	..P.	A1	1
DS	Acacia sp. T (KRN 9786)	..P.	B2	..	1
DS	Acacia sp. U (KRN 8138)	..P.	2KC	A1	2
SS	Acacia sp. V (KRN 8141)	..P.	3KC	B1	2
TS	Acacia sp. W (KRN 8276)	..P.	2E	A3	1
DS	Acacia sp. X (KRN 8280)	..P.	1E	A1	1
164	CAESALPINIACEAE	C4	2,5
SS	Cassia cardiosperma F. Muell.	..PAP	C3	6
SS	Cassia nemophila Cunn. ex Vogel var. nemophila	..PAP	6
LS	Cassia picurocarpa F. Muell. var. angustifolia Symon	..P.	B2	6
MS	Labichea lanceolata Benth. ssp. brevifolia (Meissner)J.H. Ross	..AP	C4	4

Appendix 1 (continued). Flora List

LF	REA	FPS	GS	LANDFORM								DIS		
				B	D	F1	F2	G	L	M	S			
165	PAPILIONACEAE													
DS	<i>Aotus</i> aff. <i>procumbens</i> Meissner (MAB 1865)	A2	B2	..		4	
DS	<i>Aotus</i> sp. A (KRN 9833)	PA.	IEC	A3		1	
SS	<i>Bossiaea</i> <i>dentata</i> (R. Br.)Benth.	B2		2,3	
DS	<i>Bossiaea</i> <i>leptacantha</i> E. Pritzel	APP	C4	B3	..		4,5	
DS	<i>Bossiaea</i> <i>preissii</i> Meissner	B2	..		2,3	
DS	<i>Brachysema</i> <i>daviesioides</i> (Turcz.)Benth.	P.P	C2		4,5	
MP	<i>Brachysema</i> <i>latifolium</i> R. Br.	..P	A2		2,3	
DS	<i>Burtonia</i> <i>conferta</i> DC.	B2		4	
DS	<i>Burtonia</i> <i>hendersonii</i> (Paxton)Benth.	A2		4,5	
SS	<i>Burtonia</i> <i>scabra</i> (Smith)R. Br.	B3		2,3	
DS	<i>Burtonia</i> aff. <i>viscida</i> E. Pritzel (MAB 2478)	A1		4	
DS	<i>Chorizema</i> <i>aciculare</i> (DC.)C. Gardner	P.A	C3	C2	..		4	
DS	<i>Chorizema</i> <i>cytisoides</i> Turcz.	A1	..		2,3	
DS	<i>Chorizema</i> <i>nervosum</i> T. Moore	..P	B2		3	
DS	<i>Chorizema</i> <i>uncinatum</i> C.R.P. Andrews	..P	C2	C2		3	
DS	<i>Chorizema</i> sp. A (MAB 2135)	...	3V	A1		2	
MS	<i>Daviesia</i> <i>benthamii</i> Meissner ssp. <i>benthamii</i>	AAP	D4	B3		6	
DS	<i>Daviesia</i> <i>benthamii</i> Meissner ssp. A (MAB 2215)	C2		1	
SS	<i>Daviesia</i> <i>gracilis</i> M.D. Crisp	A1		4	
SS	<i>Daviesia</i> <i>incrassata</i> Smith	D3		4	
DS	<i>Daviesia</i> <i>lancifolia</i> Turcz.	P.P	B2	B2	B2		4	
SS	<i>Daviesia</i> <i>nematophylla</i> F. Muell. ex Benth.	C3		2,5	
DS	<i>Daviesia</i> <i>nudiflora</i> Meissner	..P	D3		4	
SS	<i>Daviesia</i> <i>pachyphylla</i> F. Muell.	A3		3	
DS	<i>Daviesia</i> <i>rhombifolia</i> Meissner	..P	B3		4	
DS	<i>Daviesia</i> <i>teretifolia</i> R. Br. ex Benth.	..P	B2	B2	2	
DS	<i>Daviesia</i> sp. A (MAB 1552)	B2	..	2	
DS	<i>Daviesia</i> sp. B (MAB 4623)	..P	3KC	B1	..	4	
DS	<i>Daviesia</i> sp. C (KRN 8162)	..P	IEC	A4	..	1	
DS	<i>Daviesia</i> sp. D (MAB 2364)	B2	..	2	
DS	<i>Daviesia</i> sp. E (MAB 4525)	..P	3VC	B3	..	1	
SS	<i>Daviesia</i> sp. F (KRN 6800)	C2	..	3	
DS	<i>Dillwynia</i> <i>acerosa</i> S. Moore	...	3K	B3	..	2,5	
DS	<i>Dillwynia</i> <i>divaricata</i> (Turcz.)Benth.	...	3K	B3	..	2,3	
SS	<i>Dillwynia</i> <i>uncinata</i> (Turcz.)J. Black	P.P	B2	A1	A1	B2		4	
DS	<i>Eutaxia</i> <i>cuneata</i> Meissner	A2	..	2,3	
DS	<i>Eutaxia</i> <i>densifolia</i> Turcz.	A2	..	4	
DS	<i>Eutaxia</i> <i>microphylla</i> (R. Br.)J. Black var. <i>microphylla</i>	..P	A1	..	6	
DS	<i>Eutaxia</i> <i>parvifolia</i> Benth.	..P	A2	..	4	
DS	<i>Gastrolobium</i> <i>bilobum</i> R. Br.	4	
SS	<i>Gastrolobium</i> <i>reticulatum</i> (Meissner)Benth.	P.P	B2	..	2,3	
MS	<i>Gastrolobium</i> <i>spinosum</i> Benth. var. <i>spinosum</i>	..AP	C3	..	4	
DS	<i>Gompholobium</i> <i>baxteri</i> Benth.	..P	B2	..	2,3	
DS	<i>Gompholobium</i> <i>knightianum</i> Lindley	..P	C2	..	4	
DS	<i>Gompholobium</i> <i>marginatum</i> R. Br.	B1	A1	4	
DS	<i>Gompholobium</i> <i>viscidulum</i> Meissner	B1	..	4	
DS	<i>Hovea</i> <i>pungens</i> Benth.	..P	C3	..	4	
DS	<i>Hovea</i> <i>trisperma</i> Benth.	A2	4	
SS	<i>Indigofera</i> <i>australis</i> Willd. var. <i>australis</i>	..PP	C2	..	6	
DS	<i>Isotropis</i> <i>drummondii</i> Meissner	A1	4	
DS	<i>Jacksonia</i> <i>aphylla</i> (Turcz.)Druce	..P	C3	..	2,3	
DS	<i>Jacksonia</i> <i>capitata</i> Benth.	..P	B2	C2	4	
SS	<i>Jacksonia</i> <i>lehmannii</i> Meissner	A1	4	
DS	<i>Jacksonia</i> <i>racemosa</i> Meissner	A1	4	
DS	<i>Jacksonia</i> aff. <i>racemosa</i> Meissner (KRN 9789)	C2	..	2,3	
MP	<i>Kennedia</i> <i>beckxiana</i> (F. Muell.)F. Muell.	...	2R	B2	..	3	
MP	<i>Kennedia</i> <i>eximia</i> Lindley	..P	B1	..	2,3	
DS	<i>Latrobea</i> sp. A (MAB 3862)	A2	..	1	
AS	* <i>Medicago</i> <i>polymorpha</i> L.	A2	..	7	
AS	* <i>Medicago</i> <i>truncatula</i> Gaertner ssp. <i>longispinea</i> Urb.	A2	..	7	
SS	<i>Mirbelia</i> <i>multicaulis</i> (Turcz.)Benth.	B1	..	4	
SS	<i>Mirbelia</i> sp. A (KRN 7372)	B1	..	4	
DS	<i>Oxylobium</i> <i>microphyllum</i> Benth.	P..	C3	B2	2,3	
SS	<i>Oxylobium</i> <i>obovatum</i> Benth.	A2	..	4	
SS	<i>Oxylobium</i> <i>parviflorum</i> Benth. var. <i>parviflorum</i>	..P	D2	B2	4	
SS	<i>Oxylobium</i> <i>parviflorum</i> Benth. var. <i>revolutum</i> C. Gardner	A1	4	
SS	<i>Oxylobium</i> <i>parviflorum</i> Benth. var. <i>stenocarpum</i> C. Gardner	A2	4	
SS	<i>Phyllota</i> sp. A (MAB 3726)	A2	..	1	
SS	<i>Pultenaea</i> <i>adunca</i> Turcz.	P.P	C3	..	2,3	
DS	<i>Pultenaea</i> <i>arida</i> E. Pritzel	..AP	3WC	C3	..	2,3	
DS	<i>Pultenaea</i> <i>barbata</i> C.R.P. Andrews	A2	..	2,3	
DS	<i>Pultenaea</i> <i>capitata</i> (Turcz.)Druce	A2	4	
DS	<i>Pultenaea</i> <i>conferta</i> Benth.	A.P	B3	..	2,3	
DS	<i>Pultenaea</i> <i>elastica</i> (F. Muell.)M.D. Crisp	A.P	B3	..	6	
DS	<i>Pultenaea</i> <i>ericifolia</i> Benth.	..P	A1	..	2,3	

Appendix 1 (continued). Flora List

LF		SEB		LANDFORM									DIS	
		FPS	CS	B	D	F1	F2	G	L	M	S			
DS	<i>Pultenaea neurocalyx</i> Turcz. var. <i>neurocalyx</i>	..P	C2	..	2,3	
DS	<i>Pultenaea spinulosa</i> (Turcz.)Benth.	A2	..	4,5	
DS	<i>Pultenaea verruculosa</i> Turcz. var. <i>verruculosa</i>	A2	..	2,3	
DS	<i>Pultenaea</i> sp. A (MAB 3600)	B2	..	1	
DS	<i>Pultenaea</i> sp. B (MAB 4280)	..P	2VC	B3	..	1	
DS	<i>Pultenaea</i> sp. C (MAB 2565)	...	3V	B2	..	1	
DS	<i>Pultenaea</i> sp. D (MAB 2826)	...	2V	B3	..	1	
DS	<i>Pultenaea</i> sp. E (MAB 2798)	...	1E	B2	..	1	
DS	<i>Pultenaea</i> sp. F (MAB 2835)	B2	..	1	
DS	<i>Pultenaea</i> sp. G (MAB 3600b)	A2	..	1	
DS	<i>Pultenaea</i> sp. H (KRN 7928)	...	1E	A2	..	1	
DS	<i>Pultenaea</i> sp. I (KRN 9441)	A1	..	1	
DS	<i>Sphaerolobium daviesioides</i> Turcz.	P.P	A2	2,3	
DS	<i>Sphaerolobium grandiflorum</i> (R. Br.)Benth.	A2	..	4	
DS	<i>Sphaerolobium linophyllum</i> (Huegel)Benth.	A..	A2	..	4	
DS	<i>Sphaerolobium macranthum</i> Meissner	A2	..	4	
SS	<i>Swainsona colutooides</i> F. Muell.	..P	A2	..	6	
DS	<i>Templetonia aculeata</i> (F. Muell.)Benth.	..P	A2	6	
MS	<i>Templetonia retusa</i> (Vent.)R. Br.	C3	..	6	
SS	<i>Templetonia sulcata</i> (Meissner)Benth.	..P	A2	C3	B3	6
167 GERANIACEAE														
AS	* <i>Erodium cicutarium</i> (L.)L'Her.	A2	B2	..	7
AS	<i>Erodium crinitum</i> Carolin	C3	6
HP	<i>Pelargonium australe</i> Willd.	..PP	D4	..	A1	6
HP	<i>Pelargonium drummondii</i> Turcz.	..P	C4	4
168 OXALIDACEAE														
HP	<i>Oxalis corniculata</i> L.	..P	C3	..	C3	7
170 LINACEAE														
AS	<i>Linum marginale</i> Cunn. ex Planchon	..P	A2	6
173 ZYGOPHYLLACEAE														
DS	<i>Zygophyllum apiculatum</i> F. Muell.	C2	6
DS	<i>Zygophyllum aurantiacum</i> (Lindley)F. Muell.	..P	B2	..	6
DS	<i>Zygophyllum glaucum</i> F. Muell.	PPP	B2	..	6
AS	<i>Zygophyllum ovatum</i> Ewart & J. White	..P	C2	..	B2	6
175 RUTACEAE														
SS	<i>Boronia baeckeacea</i> F. Muell.	AAA	C3	4
DS	<i>Boronia coerulescens</i> F. Muell. ssp. <i>coerulescens</i>	P..	B2	6
DS	<i>Boronia crassifolia</i> Bartling	AAP	C3	C3	..	4
DS	<i>Boronia crenulata</i> Smith var. <i>crenulata</i>	A2	4
DS	<i>Boronia crenulata</i> Smith var. <i>gracilis</i> (Benth.) Paul G. Wilson	A2	4
DS	<i>Boronia</i> aff. <i>fabianooides</i> (Diels)Paul G. Wilson (KRN 7973)	...	3V	C2	2,3
DS	<i>Boronia inconspicua</i> Benth.	A2	2,3
DS	<i>Boronia inornata</i> Turcz. ssp. <i>inornata</i>	PAP	D5	2,3
DS	<i>Boronia inornata</i> Turcz. ssp. <i>leptophylla</i> (Turcz.) Burgman	D5	6
DS	<i>Boronia oxyantha</i> Turcz. var. <i>brevicalyx</i> (Benth.) Paul G. Wilson	...	3RC	A3	3
DS	<i>Boronia ramosa</i> (Lindley)Benth.	P.P	A1	4
DS	<i>Boronia scabra</i> Lindley	A3	..	A3	4
DS	<i>Boronia spathulata</i> Lindley	C3	..	B3	4
DS	<i>Eriostemon fitzgeraldii</i> C.R.P. Andrews	..PP	3RC	C2	C2	..	2
DS	<i>Eriostemon rhomboideus</i> Paul G. Wilson	..P	A2	2
DS	<i>Eriostemon thryptomenoides</i> S. Moore	A2	..	A2	3,5
DS	<i>Eriostemon</i> aff. <i>thryptomenoides</i> S. Moore (MAB 1535)	...	2E	C2	1
TS	<i>Geijera linearifolia</i> (DC.)J. Black	..P	B2	..	6
DS	<i>Microcybe multiflora</i> Turcz. var. <i>multiflora</i>	APP	D4	6
DS	<i>Microcybe multiflora</i> Turcz. var. <i>baccharoides</i> (F. Muell.)Ewart & Tovey	P.P	D3	2
DS	<i>Microcybe pauciflora</i> Turcz.	P..	B2	6
MS	<i>Nematolepis phebaloides</i> Turcz.	..P	B2	C3	..	2,3
SS	<i>Phebalium filifolium</i> Turcz.	AAP	C3	4,5
SS	<i>Phebalium lepidotum</i> (Turcz.)Paul G. Wilson var. lepidotum	P.P	B2	C2	..	4,5
SS	<i>Phebalium microphyllum</i> Turcz.	AP.	A2	4,5
SS	<i>Phebalium</i> aff. <i>microphyllum</i> Turcz.(KRN 6109)	..P	A2	2,5
183 POLYGALACEAE														
DS	<i>Comesperma acerosum</i> Steetz	...	3V	C2	4
DS	<i>Comesperma calymera</i> Labill.	..P	A1	6
DS	<i>Comesperma ciliatum</i> Steetz	..P	A1	4,5

Appendix 1 (continued). Flora List

LF		RES		LANDFORM								DIS	
		PPF	C5	B	D	F1	F2	G	L	M	S		
DS	<i>Comesperma confertum</i> Labill.	A1	..	4
DS	<i>Comesperma drummondii</i> Steetz	PPP	B2	4,5
CL	<i>Comesperma integerrimum</i> Endl.	A1	4,5
DS	<i>Comesperma lanceolatum</i> (R. Br.) Benth.	B1	..	2,3
SS	<i>Comesperma scoparium</i> Steetz	B1	..	4
DS	<i>Comesperma spinosum</i> F. Muell.	PPA	C3	B3	B3	..	4
CL	<i>Comesperma volubile</i> Labill.	PPP	B1	B1	..	6
185	EUPHORBIACEAE												
DS	<i>Amperea ericoides</i> Adr. Juss.	P.P	A1	A1	..	4
DS	<i>Amperea</i> sp. A (KRN 5147)	..P	A2	1
SS	<i>Beyeria lechenaultii</i> (DC.) Baillon	PAP	C2	6
DS	<i>Monotaxis occidentalis</i> Endl.	A1	A1	4
DS	<i>Phyllanthus calycinus</i> Labill.	..AA	C2	..	C2	6
AS	<i>Poranthera microphylla</i> Brongn.	..PP	A1	A1	6
AS	<i>Poranthera</i> sp. A (MAB 3739)	A2	1
LS	<i>Ricinocarpus trichophorus</i> Muell. Arg.	..P	2VC	B4	2,3
DS	<i>Stachystemon brachyphyllum</i> Muell. Arg.	B2	B1	..	2,3
DS	<i>Stachystemon polyandrus</i> (F. Muell.) Benth.	B2	B2	2,3
DS	<i>Stachystemon</i> sp. A (KRN 9773)	...	1E	A2	..	1
202	STACKHOUSIACEAE												
DS	<i>Stackhousia monogyna</i> Labill.	PPP	B3	B3	4,5
DS	<i>Stackhousia muricata</i> Lindley	A2	6
DS	<i>Stackhousia scoparia</i> Benth.	..P	B1	A1	..	4,5
DS	<i>Tripterococcus brunonis</i> Endl.	P..	A2	B2	4
207	SAPINDACEAE												
MS	<i>Dodonaea amblyophylla</i> Diels	AAA	C3	4,5
DS	<i>Dodonaea bursariifolia</i> F. Muell.	PPP	C4	B3	..	6
DS	<i>Dodonaea caespitosa</i> Diels	..P	A1	4
SS	<i>Dodonaea ceratocarpa</i> Endl.	..AP	E4	4
MS	<i>Dodonaea concinna</i> Benth.	..P	B2	2,3
DS	<i>Dodonaea glandulosa</i> J.G. West	..P	2VC	B3	2
DS	<i>Dodonaea pinifolia</i> Miq.	..P	A1	..	4,5
LS	<i>Dodonaea ptarmicaefolia</i> Turcz.	..AP	C3	2,3
SS	<i>Dodonaea stenozyga</i> F. Muell.	PAP	C4	6
MS	<i>Dodonaea viscosa</i> Jacq.	..P	B2	6
TS	<i>Heterodendrum oleaeifolium</i> Desf.	B3	..	6
215	RHAMNACEAE												
DS	<i>Cryptandra glabriflora</i> Benth.	AAA	D3	C3	4,5
DS	<i>Cryptandra nutans</i> Steudel	..P	C3	4,5
DS	<i>Cryptandra parvifolia</i> Turcz.	C3	2,5
DS	<i>Cryptandra polyclada</i> Diels	P..	A2	2
SS	<i>Cryptandra pungens</i> Steudel	A.P	C3	..	C3	A1	..	4,5
DS	<i>Cryptandra</i> sp. A (MAB 3897)	C3	1
DS	<i>Cryptandra</i> sp. B (MAB 3257)	C3	1
SS	<i>Pomaderris intangenda</i> F. Muell.	...	3V	A2	2,5
MS	<i>Pomaderris myrtilloides</i> Fenzl	C3	2,5
SS	<i>Pomaderris</i> sp. A (KRN 9788)	B1	..	1
DS	<i>Siegfriedia darwinoides</i> C. Gardner	...	3VC	B2	2,3
DS	<i>Spyridium complicatum</i> F. Muell.	P.P	B2	D2	..	B2	..	2,5
DS	<i>Spyridium cordatum</i> (Turcz.) Benth.	P..	C3	2,3
SS	<i>Spyridium oligocephalum</i> (Turcz.) Benth.	P.P	3RC	C3	4
DS	<i>Spyridium rotundifolium</i> F. Muell.	..P	B3	2
DS	<i>Spyridium</i> sp. A (KRN 6108)	...	3K	B3	2
DS	<i>Spyridium</i> sp. B (KRN 7931)	...	3KC	A2	2
DS	<i>Spyridium</i> sp. C (KRN 8212)	A2	2,5
MS	<i>Trymalium</i> aff. <i>ledifolium</i> Fenzl (KRN 5606)	..A.	A2	2,5
DS	<i>Trymalium</i> sp. A (MAB 2070)	C2	2
221	MALVACEAE												
HP	<i>Alyogyne hakeifolia</i> (Giord.) Alef.	..P	A2	6
HP	<i>Alyogyne huegelii</i> (Endl.) Fryx. var. <i>grossulariaeifolius</i> (Miq.) A.S. Mitchell	A2	6
DS	<i>Lawrenzia berthae</i> (F. Muell.) Melville	..P	C3	6
DS	<i>Lawrenzia densiflora</i> (E.G. Barker) Melville	C3	..	5
HP	<i>Lawrenzia diffusa</i> (Benth.) Melville	..P	3VC	B2	A2	..	4,5
HP	<i>Lawrenzia spicata</i> Hook.	A3	6
SS	<i>Lawrenzia squamata</i> Nees ex Miq.	..A.	B5	..	6
223	STERCULIACEAE												
SS	<i>Guichenotia ledifolia</i> Gay	..A.	A2	B2	4
DS	<i>Lasiopetalum compactum</i> S. Paust	A2	..	2,3
DS	<i>Lasiopetalum indutum</i> Steudel	..P	A2	..	2,3
DS	<i>Lasiopetalum quinquevium</i> Turcz.	A2	2,3
DS	<i>Lasiopetalum rosmarinifolium</i> (Turcz.) Benth.	P.P	C3	4

Appendix 1 (continued). Flora List

LF	REF	LANDFORM										D16		
		FPS	CB	B	D	F1	F2	G	L	M	S			
SS	Rulingia cygnorum (Steudel)C. Gardner var. cygnorum	C1	4
SS	Rulingia aff. platycalyx Benth. (MAB 2039)	B2	1
SS	Rulingia rotundifolia Turcz.	A2	2,5
DS	Thomasia angustifolia Steudel	..P	B2	4
DS	Thomasia foliosa Gay	B2	4
DS	Thomasia grandiflora Lindley	C1	4
DS	Thomasia microphylla S. Paust	A2	..	2,3
DS	Thomasia petalocalyx F. Muell.	..P	A2	6
DS	Thomasia purpurea (Aiton)Gay	A2	4
226	DILLENACEAE													
DS	Hibbertia acerosa (R. Br. ex DC.)Benth.	..P	C3	C3	4
DS	Hibbertia eatoniae Diels	P..	A2	..	4
DS	Hibbertia andrewsiana Diels	...	3RC	B3	..	3
DS	Hibbertia exasperata (Steudel)Briq.	P..P	C3	4
DS	Hibbertia gracilipes Benth.	A..P	C3	C3	..	4
DS	Hibbertia aff. gracilipes Benth. (MAB 1048)	..P	A2	1
DS	Hibbertia inclusa Benth.	P..P	C3	4
DS	Hibbertia lineata Steudel	C3	4
DS	Hibbertia pungens Benth.	PAA	C3	B1	..	4,5
DS	Hibbertia racemosa (Endl.)Gilg	B1	B1	..	4
DS	Hibbertia recurvifolia (Steudel)Benth.	..P	C3	4
DS	Hibbertia aff. recurvifolia (Steudel)Benth. (MAB 1583)	..P	B1	1
DS	Hibbertia rupicola (S. Moore)C. Gardner	APP	C3	B3	4
DS	Hibbertia sp. A (MAB 1055)	A3	2
DS	Hibbertia sp. B (MAB 1107)	..P	A1	2
DS	Hibbertia sp. C (MAB 1822)	B1	2
DS	Hibbertia sp. D (MAB 2193)	P..	A2	2
DS	Hibbertia sp. E (MAB 3823)	A1	2,5
DS	Hibbertia sp. F (MAB 3771)	A1	2
DS	Hibbertia sp. G (MAB 1450)	A2	2,5
DS	Hibbertia sp. H (MAB 4028)	..P	2,5
236	FRANKENIACEAE													
MP	Frankenia brachyphylla Summerh.	..P	B4	C3	..	5
DS	Frankenia cinerea A. DC.	..AP	C3	D4	..	6
DS	Frankenia densa Summerh.	B3	..	6
DS	Frankenia desertorum Summerh.	..A	B4	..	6
DS	Frankenia pauciflora DC.	..A	B3	..	6
243	VIOLACEAE													
DS	Hybanthus floribundus (Lindley)F. Muell. ssp. floribundus	A3	..	6
263	THYMELAEACEAE													
DS	Pimelea angustifolia R. Br.	..P	C3	C1	..	4
SS	Pimelea argentea R. Br.	A2	4
DS	Pimelea brachyphylla Benth.	..P	B2	..	4
DS	Pimelea brevifolia R. Br.	P..P	A2	A2	..	4,5
DS	Pimelea suaveolens Meissner	..P	B3	..	4,5
SS	Pimelea aff. suaveolens Meissner (KRN 25)	A1	..	2,3
DS	Pimelea sulphurea Meissner	B2	..	4
DS	Pimelea tinctoria Meissner	A..	B3	..	2,3
SS	Pimelea sp. A (KRN 70)	A1	..	2,3
DS	Pimelea sp. B (MAB 2846)	A1	..	2
MS	Pimelea sp. C (KRN 9769)	A1	..	3
273	MYRTACEAE													
SS	Agonis obtusissima F. Muell.	A2	..	2,3
SS	Agonis spathulata Schauer	..P	C3	..	2,3
DS	Angasomyrtus salina M.E. Trudgen & G.J. Keighery	..P	2VC	C4	1
SS	Astartea ambigua F. Muell.	AFA	C2	C3	C4	B2	D3	..	4
SS	Astartea clavulata Turcz.	A2	..	4
DS	Astartea heteranthera C. Gardner	..P	A2	2,5
SS	Baeckea blackettii F. Muell.	..P	2,3
SS	Baeckea corynophylla F. Muell.	B2	..	4
SS	Baeckea crassifolia Lindley var. icosandra F. Muell. ex Benth.	...	3RC	A2	2
SS	Baeckea aff. crassifolia Lindley (MAB 1666)	A1	2
SS	Baeckea crispiflora F. Muell.	PPP	4
MS	Baeckea latens C.R.P. Andrews	..P	C3	..	C3	4,5
SS	Baeckea aff. latens C.R.P. Andrews A (MAB 4271)	P..	B2	..	A2	B2	2
SS	Baeckea aff. latens C.R.P. Andrews B (MAB 4618)	..P	A2	2
SS	Baeckea polyandra F. Muell.	A2	..	B2	..	3
SS	Baeckea preissiana (Schauer)Domin	PAP	C3	4,5
DS	Beaufortia elegans Schauer	A1	4
SS	Beaufortia empetrifolia (Reichb.)Schauer	..A	D5	C3	2,3

Appendix I (continued). Flora List

LF	RES	LANDFORM								DIS		
		TPB	CS	B	D	F1	F2	G	L		M	S
SS	Beaufortia aff. empetrifolia (Reichb.)Schauer (KRN 7970)	...	IV	A3	2
DS	Beaufortia micrantha Schauer var. micrantha	AAA	D4	C4	4
DS	Beaufortia aff. micrantha Schauer (MAB 1712)	B3	A3	2
SS	Beaufortia schaueri Preiss ex Schauer	P2A	C4	...	D5	...	B2	B3	4
SS	Beaufortia aff. schaueri Preiss ex Schauer (MAB 2737)	..P	B3	1
DS	Calothamnus gibbosus Benth.	..P	C3	2,3
DS	Calothamnus gracilis R. Br.	AAA	C3	C3	...	E3	C3	...	2,3
SS	Calothamnus lateralis Lindley	..P	A1	4
MS	Calothamnus quadrifidus R. Br.	AAA	C3	...	E4	...	A1	B2	4
MS	Calothamnus villosus R. Br.	A2	4
DS	Calytrix decandra DC.	P..	C3	...	3
DS	Calytrix leschenaultii (Schauer)Benth.	APA	D4	...	C3	...	C2	C3	4,5
DS	Calytrix aff. stipulosa W. Fitzg.	PAA	A2	C4	A3	2,5
DS	Calytrix tenuiramea (Turcz.)Benth.	A2	4
SS	Calytrix tetragona Labill.	..P	A2	A3	6
DS	Calytrix sp. (MAB 3267)	A2	1
MS	Chamelaucium axillare F. Muell. ex Benth.	A2	2,3
MS	Chamelaucium ciliatum Desf.	P.P	C3	C4	2,3
DS	Chamelaucium drummondii Meissner	A1	4
MS	Chamelaucium megalopetalum F. Muell. ex Benth.	..P	C4	B2	...	4
SS	Chamelaucium sp. A (KRN 7954)	...	2V	A1	2,5
SS	Chamelaucium sp. B (MAB 4350)	..P	A1	2
DS	Conothamnus aureus (Turcz.)Domin	D3	...	4
DS	Darwinia diosmoides (DC.)Benth.	.PA	C4	B3	A3	...	4,5
DS	Darwinia aff. luehmannii F.Muell. & Tate (MAB 3241)	A2	4
DS	Darwinia vestita (Endl.)Benth.	C2	...	4
DS	Darwinia sp. A (MAB 1110)	..P	C3	1
DS	Darwinia sp. B (MAB 1839)	..P	C3	1
DS	Darwinia sp. C (MAB 1850)	..P	B2	1
DS	Darwinia sp. D (MAB 1274)	A2	1
DS	Darwinia sp. E (KRN 2426)	..P	H2	...	2,3
DS	Darwinia sp. F (MAB 4517)	..P	1EC	A4	1
DS	Darwinia sp. G (KRN 7975)	...	1E	A2	1
DS	Darwinia sp. H (KRN 9775)	A1	...	1
MAT	Eucalyptus anceps (R. Br. ex Maiden)Blakely	C4	6
MAT	Eucalyptus angulosa Schauer	B4	...	6
MAS	Eucalyptus angustissima F. Muell.	...	2RC	A3	B5	2
MAS	Eucalyptus aff. angustissima F. Muell. (MAB 3767)	...	3E	A2	1
MAT	Eucalyptus annulata Benth.	C6	4,5
MAS	Eucalyptus calycogona Turcz.	.AP	D5	...	B3	6
MAS	Eucalyptus celastroides Turcz. ssp. virella Brooker	AA.	B4	4
MAS	Eucalyptus conglobata (R. Br. ex Benth.)Maiden	..P	C4	C4	6
MAS	Eucalyptus cylindriflora Maiden & Blakely	AP.	C5	A3	...	4,5
MAS	Eucalyptus aff. cylindriflora Maiden & Blakely (MAB 1840)	A2	1
MAS	Eucalyptus decipiens Endl.	B4	4
MAS	Eucalyptus deflexa Brooker	P..	3VC	B4	2
ST	Eucalyptus dielsii C. Gardner	..P	3VC	C5	2
ST	Eucalyptus diptera C.R.P. Andrews	AA.	C5	B5	...	2
ST	Eucalyptus aff. diptera C.R.P. Andrews A (MAB 2980)	C5	1
ST	Eucalyptus aff. diptera C.R.P. Andrews B (MAB 2317)	...	2R	C4	5
MAS	Eucalyptus discreta Brooker	...	3RC	B5	2,5
MAS	Eucalyptus aff. discreta Brooker (MAB 4472)	..P	A3	1
ST	Eucalyptus eremophila (Diels)Maiden	AAA	C5	D5	E6	...	C4	...	4,5
MAS	Eucalyptus falcata Turcz.	C4	4
ST	Eucalyptus flocktoniae (Maiden)Maiden	AAA	C4	D5	6
MAS	Eucalyptus foecunda Schauer	A.P	C3	C5	C5	6
MAS	Eucalyptus foecunda Schauer ssp. A (MAB 2650)	..P	A3	2
MAS	Eucalyptus forrestiana Diels ssp. forrestiana	..A	3VC	C5	B2	...	2
MAS	Eucalyptus forrestiana Diels ssp. dolichorrhyncha Brooker	..P	2EC	B4	1
MAS	Eucalyptus forrestiana Diels ssp. stoatei C.J. Robinson	...	2E	B5	3
MAS	Eucalyptus gardneri Maiden	B3	A5	...	4
DT	Eucalyptus keesellii Maiden & Blakely	..A	D5	2,3
MAS	Eucalyptus gracilis F. Muell.	..P	C5	6
MAS	Eucalyptus aff. gracilis F. Muell. A (MAB 2564)	C5	1
MAS	Eucalyptus aff. gracilis F. Muell. B (MAB 4021)	..P	B2	1
MAS	Eucalyptus grossa F. Muell. ex Benth.	.AP	C5	2,5
MAS	Eucalyptus halophila D.J. Carr & S.G.M. Carr	..P	3VC	B5	B5	...	2
MAS	Eucalyptus incrassata Labill.	PAA	C4	D5	...	C4	...	6

Appendix 1 (continued). Flora List

LF		HERS		LANDFORM								Dist	
		FPS	CS	B	D	F1	F2	G	L	M	S		
SS	Melaleuca glaberrima F. Muell.	.PA	C3	C4	B3	4
TS	Melaleuca hamulosa Turcz.	P.P	C3	4
MS	Melaleuca holosericea Schauer var. holosericea	AA.	B2	..	D4	2,3
MS	Melaleuca aff. holosericea Schauer (MAB 3060)	A2	2
MS	Melaleuca aff. lanceolata Otto (MAB 2546)	A4	2,5
MS	Melaleuca lateralis Turcz.	PP.	D4	B2	4
MS	Melaleuca lateriflora Benth.	AAA	D4	4,5
SS	Melaleuca leptospermoides Schauer	P.P	C3 B3	4
SS	Melaleuca aff. leptospermoides Schauer (MAB 2820)	A2	2
LS	Melaleuca aff. nesophila F. Muell. (KRN 8284)	..A	3RC	C3	1
TS	Melaleuca pauperiflora F. Muell.	APA	D4	6
TS	Melaleuca aff. pauperiflora F. Muell. (KRN 7694)	D4	2
DS	Melaleuca pentagona Labill. var. pentagona	A..	C3	B5	4
MS	Melaleuca pentagona Labill. var. subulifolia Schauer	D4	3
SS	Melaleuca pulchella R. Br.	..P	B3	D3	2,3
MS	Melaleuca pungens Schauer	..P	4
MS	Melaleuca aff. pungens Schauer (MAB 3842)	AAP	C3	4
TS	Melaleuca quadrifaria F. Muell.	P.P	D4	2
DS	Melaleuca scabra R. Br.	AAA	D4	D4	C3	..	A3 C4	2,5
DS	Melaleuca aff. scabra R. Br. (MAB 2552)	..AP	4
MS	Melaleuca sparsiflora Turcz.	B4	4
MS	Melaleuca striata Labill.	..	2K	C3	4
MS	Melaleuca suberosa (Schauer)C. Gardner	..P	C4	2,3
MS	Melaleuca subfalcata Turcz.	..P	B2	2,3
SS	Melaleuca subtrigona Schauer	..P	B3	C3	..	2,3
MS	Melaleuca tenella Benth.	..P	C3	2,5
MS	Melaleuca thymoides Labill.	..P	A2	2,3
MS	Melaleuca thyooides Turcz.	..P	C3	B3	2,3
MS	Melaleuca aff. thyooides Turcz. (MAB 1836)	..P	C3	D4	4,5
MS	Melaleuca uncinata R. Br.	PP.	B2	2
MS	Melaleuca undulata Benth.	AAA	C3	D4	D4	C4	C3	B4	C3	6
MS	Melaleuca aff. undulata Benth. (KRN 6432)	P..	A3	B3	A2	4
MS	Melaleuca urceolaris F. Muell. ex Benth.	PAA	D4	2
MS	Melaleuca aff. urceolaris F. Muell. ex Benth. (MAB 4496)	..P	D4	4
MS	Melaleuca sp. A (MAB 4516)	..P	D4	1
MS	Melaleuca sp. B (MAB 2795)	..P	D4	1
MS	Melaleuca sp. C (MAB 4581)	C4	1
MS	Melaleuca sp. D (MAB 1467)	..P	3KC	C3	1
SS	Melaleuca sp. E (MAB 1863)	B2	1
MS	Melaleuca sp. F (MAB 2583)	..P	C3	2
MS	Melaleuca sp. G (MAB 4551)	B2	2
MS	Melaleuca sp. H (MAB 3921)	..P	C4	2
MS	Melaleuca sp. I (MAB 4317)	B3	2
MS	Melaleuca sp. J (MAB 4499)	..P	C4	2
MS	Melaleuca sp. K (KRN 7962)	C4	2
MS	Melaleuca sp. L (KRN 9774)	B2	2
MS	Melaleuca sp. M (KRN 9783)	A3	3
SS	Melaleuca sp. N (KRN 8186)	A4	3
SS	Micromyrtus elobata (F. Muell.)Benth.	..	1E	A3	1
SS	Micromyrtus imbricata Benth.	..P	C3	C3	2,3
SS	Micromyrtus obovata (Turcz.)J.W. Green	..P	C3	2
DS	Micromyrtus racemosa Benth.	AAA	C2	4,5
SS	Pericalymma ellipticum (Endl.)Schauer	..P	C3	2,5
MS	Phymatocarpus maxwellii F. Muell.	..P	D4	C4	4
SS	Rinzia communis M.E. Trudgen	PAA	C4	E5	2,3
SS	Rinzia sp. A (MAB 4433)	..P	B2	B2	2
MS	Thryptomene appressa C.R.P. Andrews	..P	A2	2
MS	Thryptomene australis Endl.	P..	C3	5
SS	Thryptomene saxicola (Cunn. ex Hook.)Schauer	PAP	C3	D4	2,5
DS	Verticordia acerosa Lindley	B5	4
DS	Verticordia brownii (Desf.)DC.	..P	C3	4
DS	Verticordia chrysantha Endl.	..A	A3	B4	4,5
DS	Verticordia densiflora Lindley	A.A	C3	C3	4
DS	Verticordia aff. drummondii Schauer (MAB 2868)	A.P	C3	D3	4
DS	Verticordia endlicheriana Schauer	..P	3RC	A2	2
DS	Verticordia grandiflora Endl.	AA.	B3	A3 C3	4
DS	Verticordia humilis Benth.	..P	B2	4
DS	Verticordia insignis Endl.	P.P	A2	A2	2,3
DS	Verticordia mitchelliana C. Gardner	PPA	C3	4,5
DS	Verticordia oxylepis Turcz.	..P	B2	2,5
DS	Verticordia picta Endl.	C3	2,3
DS	Verticordia plumosa (Desf.)Druce	PA.	A2	B3	4
DS	Verticordia roei Endl.	..AA	B3	B3	B2	4
DS	Verticordia roei Endl.	AAA	C3	B4	4,5
275	ONAGRACEAE												
AB	*Oenothera stricta Lebed. ex Link	A3	7

Appendix 1 (continued). Flora List

LF	RES	PPS	CS	LANDFORM								DIE
				B	D	F1	F2	G	L	M	S	
276 HALORAGACEAE												
HP	Glischrocaryon aureum (Lindley)Orch. var. angustifolium (Nees)Orch.	P.P	B1	D3	B2	..	4,5
HP	Glischrocaryon flavescens (J. Drumm. ex Hook.) Orch.	B2	B2	6
AS	Gonocarpus nodulosus Nees	C3	4,5
DS	Gonocarpus trichostachyus (Benth.)Orch.	..	3K	B3	4
DS	Gonocarpus sp. A (KRN 9759)	A2	1
DS	Haloragis sp. A (KRN 8269)	..	1E	A2	1
DS	Haloragodendron glandulosum Orch.	A2	2,3
HY	Myriophyllum petraeum Orch.	..P	A4	2
281 APIACEAE												
AS	Daucus glochidiatus (Labill.)Fischer, C. Meyer & Ave-Lall.	PP.	B2	6
AS	Hydrocotyle alata A. Rich.	A2	4
AS	Hydrocotyle diantha DC.	B3	6
AS	Hydrocotyle medicaginoides Turcz.	..P	3RC	B3	..	4
AS	Hydrocotyle sp. A (MAB 4419)	..P	A2	1
DS	Platysace compressa (Labill.)Norman	..P	A2	4
DS	Platysace effusa (Turcz.)Norman	AAA	C3	C2	..	A2	D2	..	4,5
DS	Platysace maxwellii (F. Muell.)Norman	A.P	C3	2
AS	Trachymene anisocarpa (Turcz.)B.L. Burttt	..P	A2	6
AS	Trachymene croniniana (F. Muell.)T. Durand	..	3V	A2	3
AS	Trachymene cyanopetala (F. Muell.)Benth.	..AP	A2	6
AS	Trachymene ornata (Endl.)Druce var. ornata	..AA	B3	..	A2	B3	6
AS	Trachymene pilosa Smith	..PA	B2	6
DS	Xanthosia huegelii (Benth.)Stuedel	B1	..	4
DS	Xanthosia pusilla Bunge	B2	6
288 EPACRIDACEAE												
DS	Acrotriche cordata (Labill.)R. Br.	B1	..	6
DS	Andersonia macranthera F. Muell.	..P	2EC	B3	B2	..	3
DS	Andersonia micrantha R. Br.	..P	A1	..	3
DS	Andersonia parvifolia R. Br.	..P	C3	C3	..	2,3
DS	Astroloma drummondii Sonder	A1	..	3
DS	Astroloma epacridis (DC.)Druce	PP.	A1	4
DS	Astroloma microphyllum Stscheegl.	..P	A1	3
DS	Astroloma pallidum R. Br.	C3	4
DS	Astroloma aff. prostratum R. Br. (MAB 1176)	A1	2
DS	Astroloma tectum R. Br.	A1	..	3
DS	Brachyloma concolor (F. Muell.)C. Gardner	..P	C3	2,3
DS	Brachyloma sp. A (KRN 5521)	..	3K	A1	A1	2
SS	Coleanthera myrtoides Stscheegl.	PP.	B1	4,5
SS	Conostephium drummondii (Stschegl.)C. Gardner	PPP	3RC	B2	D4	A1	..	4
SS	Conostephium minus Lindley	..P	A2	..	4
SS	Conostephium roei Benth. (MAB 3492)	PPP	B2	..	C4	4
SS	Conostephium sp. A (MAB 3063)	..	3K	B2	1
SS	Conostephium sp. B (MAB 1656)	B2	1
DS	Leucopogon aff. allittii F. Muell. (MAB 1482)	..	2E	B2	1
DS	Leucopogon assimilis R. Br.	A2	4
DS	Leucopogon bossiaea F. Muell.	..P	3KC	B2	2,3
DS	Leucopogon brevicuspis Benth.	..P	3KC	C3	4
DS	Leucopogon aff. brevislorus F. Muell. (MAB 1207)	A2	2,3
DS	Leucopogon concinnus Benth.	C2	B2	..	2,3
DS	Leucopogon conostephioides DC.	..P.P	C4	4
DS	Leucopogon aff. conostephioides DC. (MAB 1085)	A2	2,3
DS	Leucopogon corynocarpus Sonder	..P.P	B2	2,3
DS	Leucopogon crassifolius Sonder	A2	2,3
SS	Leucopogon cuneifolius Stschegl.	..PP	B2	..	B1	..	2,3
DS	Leucopogon dielsianus E. Pritzel	..P.P	B2	2,3
DS	Leucopogon fimbriatus Stschegl.	..AA	B2	4
SS	Leucopogon flavescens Sonder var. brevifolius Benth.	D3	B2	2,5
DS	Leucopogon aff. hamulosus E. Pritzel (MAB 1211)	..A.	B2	C2	3
DS	Leucopogon minutifolius W. Fitzg.	..A	D4	..	C3	2,5
MS	Leucopogon obovatus (Labill.)R. Br.	A2	4
DS	Leucopogon aff. obovatus (Labill.)R. Br. (MAB 1125)	A2	1
DS	Leucopogon opponens F. Muell.	..	3H	A2	3
DS	Leucopogon aff. ovalifolius Sonder (MAB 1482)	A2	1
DS	Leucopogon aff. pulchellus Sonder (MAB 2738)	A2	1
SS	Leucopogon rubicundus F. Muell. ex Benth.	..A	B4	2
SS	Leucopogon woodsii F. Muell.	..	3RC	B2	2,3
DS	Leucopogon sp. A (MAB 3962)	B2	1
DS	Leucopogon sp. B (MAB 3936)	A2	1
DS	Leucopogon sp. C (MAB 3920)	B2	1
DS	Leucopogon sp. D (MAB 3447)	C3	1
DS	Leucopogon sp. E (MAB 3708)	..	1E	A2	1

Appendix 1 (continued). Flora List

LF		RES		LANDFORM								DIS
		FFS	CS	B	D	F1	F2	G	L	M	S	
DS	Leucopogon sp. F (KRN 8045)	B2	2
DS	Leucopogon sp. G (KRN 8123)	...	1E	A2	1
DS	Leucopogon sp. H (KRN 8148)	A2	2
DS	Leucopogon sp. I (KRN 8213)	...	1E	A2	1
SS	Leucopogon sp. J (KRN 5885)	C1	..	4,5
DS	Leucopogon sp. K (KRN 9791)	A1	..	1
DS	Leucopogon sp. L (KRN 9794)	A1	..	1
DS	Leucopogon sp. M (KRN 8173)	...	1V	..	B2	1
DS	Leucopogon sp. N (KRN 8283)	...	3V	A3	4,5
SS	Lysinema ciliatum R. Br.	AAA	E4	C3	C4	..	D2	D3	..	4
SS	Monotoca leucantha E. Pritzel	..P	C3	4
DS	Monotoca oligarrhenoides F. Muell.	B2	..	2
DS	Monotoca tamariscina F. Muell.	A2	3
DS	Oligarrhena micrantha R. Br.	C3	B1	..	4
DS	Styphelia intertexta A.S. George	P.P	C3	B2	4,5
SS	Styphelia pulchella (Stschegl.)Druce	A2	2,3
293	PRIMULACEAE											
AS	*Anagallis arvensis L.	..AP	D2	7
DS	Samolus repens (Forster & G. Forster)Pers.	..P	B3	6
302	LOGANIACEAE											
SS	Logania buxifolia F. Muell.	P.P	D4	2,3
DS	Logania campanulata R. Br.	A1	..	4
DS	Logania micrantha Benth.	PP.	C2	A1	..	4
SS	Logania stenophylla F. Muell.	..P	D3	3
SS	Logania aff. stenophylla F. Muell. (MAB 2174)	A2	1
AS	Mitrasacme paradoxa R. Br.	..P.	C2	..	A3	6
303	GENTIANACEAE											
AS	*Centaurium spicatum (L.)Fritsch ex Janchen	..P.	B2	7
AS	Sebaea ovata (Labill.)R. Br.	A2	..	A2	6
303A	MENYANTHACEAE											
HP	Villarsia parnassifolia (Labill.)R. Br.	..P	B2	4
304	APOCYNACEAE											
LS	Alyxia buxifolia R. Br.	..PAP	D4	6
307	CONVOLVULACEAE											
CL	Convolvulus erubescens Sims	A1	6
MP	Wilsonia humilis R. Br.	PAA	B3	..	D4	9
MP	Wilsonia rotundifolia Hook.	P.P	3RC	B2	..	6
310	BORAGINACEAE											
SS	Halgania andromedifolia Behr & F. Muell.	PAP	D4	4,5
SS	Halgania aff. andromedifolia Behr & F. Muell. (MAB 1876)	B2	2
DS	Halgania aff. preissiana Lehm. (MAB 1513)	A2	2
DS	Halgania sp. A (MAB 3547b)	...	2V	A2	2
DS	Halgania sp. B (MAB 2825)	B2	2
AS	Heliotropium sp. A (KRN 6982)	...	2R	B2	2,5
311A	CHLOANTHACEAE											
DS	Dicrastylis parvifolia F. Muell.	A2	5
SS	Lachnostachys ferruginea Hook.	C3	4
SS	Pityrodia axillaris (Endl.)Druce	A3	4,5
313	LAMIACEAE											
SS	Hemigenia eutaxioides C.R.P. Andrews	..P	D3	2,5
SS	Hemigenia aff. eutaxioides C.R.P. Andrews (MAB 1755)	...	3K	B3	..	A2	2
SS	Hemigenia teretiuscula F. Muell.	A2	2,5
SS	Hemigenia sp. A (MAB 1633)	A2	..	1
SS	Microcorys barbata R. Br.	..P	B3	4
DS	Microcorys glabra (Bartling)Benth.	P..	A2	..	B2	4
SS	Microcorys virgata R. Br.	A2	2,3
DS	Prostanthera baxteri Cunn. ex Benth.	A2	..	6
DS	Prostanthera serpyllifolia (R. Br.)Briq. ssp. microphylla (R. Br.) B.J. Conn	PPP	D3	..	6
DS	Prostanthera sp. A (MAB 1193)	...	1E	A2	..	1
DS	Teucrium myriocladum Diels	A2	..	2
AS	Teucrium sessiliflorum Benth.	A3	..	6
SS	Westringia cephalantha F. Muell.	P..	A3	..	4,5
DS	Westringia dampieri R. Br.	D4	..	6
DS	Westringia aff. dampieri R. Br. (MAB 4476)	..P	A2	..	1
DS	Westringia rigida R. Br.	APP	B4	..	6

Appendix 1 (continued). Flora List

LF	RES	LANDFORM										DIS				
		PPS	CS	B	D	F1	F2	G	L	M	S					
315	SOLANACEAE															
SS	Anthocercis genistoides Miers	.P	A1	2,5
SS	Anthocercis viscosa R. Br. ssp. caudata Haegi	D3	2,3
SS	Cyphanthera microphylla Miers	A.P	D3	2,5
HP	Solanum capsiciforme (Domin) Baylis	C3	6
HP	Solanum hoplopetalum Bitter & Summerh.	P..	C3	6
DS	*Solanum nigrum L.	C3	7
SS	Solanum simile F. Muell.	C3	6
HP	Solanum symonii H. Eichler	..P	C3	6
316	SCROPHULARIACEAE															
AS	Glossostigma diandrum (L.) Kuntze	A2	6
323	LENTIBULARIACEAE															
AS	Polypompholyx tenella (R. Br.) Lehm.	..P	C3	B2	6
326	MYOPORACEAE															
MP	Eremophila biserrata Chinn.	A1	1
LS	Eremophila calorhados Diels	P..	B2	2,5
SS	Eremophila decipiens Ostenf.	AAP	D3	6
DS	Eremophila densifolia F. Muell.	B3	2,3
MS	Eremophila denticulata F. Muell.	..P	3RC	A1	2,3
MS	Eremophila dichroantha Diels	PAP	3RC	C3	2
MS	Eremophila glabra (R. Br.) Ostenf.	P.P	D4	6
LS	Eremophila pachyphylla Diels	APP	B4	2,5
MP	Eremophila serpens Chinn.	..	3V	A2	2,3
MP	Eremophila aff. serpens Chinn. (KRN 8180)	..	3E	A1	1
SS	Eremophila sp. A (MAB 2587)	..P	A3	1
MS	Myoporum beckeri F. Muell. ex Benth.	PP.	2KC	B3	2,3
DS	Genus indet. (MAB 2851)	..	2E	B3	1
329	PLANTAGINACEAE															
AS	Plantago hispida R. Br.	.AP	C3	4
331	RUBIACEAE															
DS	Opercularia vaginata Labill.	PPP	D3	4,5
339	CAMPANULACEAE															
AS	*Wahlenbergia capensis (L.) A. DC.	..P	B2	7
AS	Wahlenbergia gracilentia Loch.	.A.	C3	..	B2	6
AS	Wahlenbergia gracilis A. DC.	B2	6
340	LOBELIACEAE															
AS	Isotoma hypocrateriformis (R. Br.) Druce	.PP	C3	..	B1	4
AS	Isotoma scapigera (R. Br.) Don	..P	B3	2,5
AS	Lobelia rarifolia F. Wimmer	.P.	B1	4
341	GOODENIACEAE															
HP	Anthotium rubriflorum F. Muell. ex Benth.	P..	C2	2,3
DS	Coopernookia polygalacea (Vriese) Carolin	..P	B3	4,5
DS	Coopernookia strophiolata (F. Muell.) Carolin	PPP	B2	D3	B2	B3	6
DS	Dampiera carinata Benth.	..P	3RC	C3	4
DS	Dampiera cuneata R. Br.	B2	2,5
DS	Dampiera fasciculata R. Br.	A1	2,3
DS	Dampiera lavandulacea Lindley	A.A	D3	B2	..	6
DS	Dampiera oligophylla Benth. ssp. juncea (Benth.) Rajput & Carolin	C1	B1	G1	4
DS	Dampiera parvifolia R. Br.	C2	2,3
DS	Dampiera sacculata F. Muell. ex Benth.	P..	A2	B2	4
DS	Dampiera tenuicaulis E. Pritzel var. tenuicaulis	B1	4,5
DS	Dampiera trigona Vriese	..	3H	B2	4
DS	Dampiera sp. A (MAB 4624)	..P	B2	1
DS	Dampiera sp. B (KRN 9800)	A1	1
RP	Goodenia affinis Vriese	P.P	A1	6
AS	Goodenia berardiana (Gaudich.) Carolin	PP.	C3	6
DS	Goodenia caerulea R. Br.	..P	C2	4
DS	Goodenia concinna Benth.	..P	B2	..	C2	4
SS	Goodenia decursiva W. Fitzg.	B2	2,3
AS	Goodenia filiformis R. Br. var. filiformis	C3	4
DS	Goodenia incana R. Br.	C2	4,5
DS	Goodenia laevis Benth.	P.P	A2	4
RP	Goodenia pinnatifida Schidl.	B2	6
DS	Goodenia pterygosperma R. Br.	A2	A2	2
DS	Goodenia scapigera R. Br.	.AP	C3	..	C2	..	B1	..	2
DS	Goodenia trichophylla (Vriese) Benth.	..P	3RC	B2	4
RP	Goodenia watsonii F. Muell. & Tate	P..	A2	2
DS	Goodenia sp. A (MAB 4334)	..P	B2	1
DS	Lechenaultia aff. floribunda Benth. (KRN 6523)	P.P	C3	2
DS	Lechenaultia formosa R. Br.	C3	B2	..	4

Appendix 1 (continued). Flora List

LF		HSH		LANDFORM								DIB
		PPS	CS	B	D	F1	F2	G	L	M	S	
MP	Lechenaultia tubiflora R. Br.	A1	..	4
DS	Lechenaultia sp. A (KRN 6470)	...	3V	A3	2
SS	Scaevola bursariifolia J. Black	.PP	C3	6
SS	Scaevola myrtifolia (Vriese)Krause	D3	6
DS	Scaevola thesioides Benth. var. filifolia	4
	E. Pritzel	B2	4
DS	Scaevola thesioides Benth. var. thesioides	B1	..	A1	4
MP	Scaevola sp. A (KRN 9677)	...	IE	A3	1
AS	Velleia cynopotamica F. Muell.	B2	6
RP	Velleia trinervis Labill.	B2	..	B2	4
343 STYLIDIACEAE												
AS	Levenhookia dubia Sonder	.P	A1	6
AS	Levenhookia pauciflora Benth.	B3	4
AS	Levenhookia pusilla R. Br.	.PP	C3	..	B2	6
AS	Levenhookia stipitata (Sonder)F. Muell.	PPP	B1	4
AS	Levenhookia sp. A (MAB 4388)	.P	A2	1
RP	Stylidium adpressum Benth.	4
RP	Stylidium assimile R. Br.	.A	C3	3
DS	Stylidium breviscapum R. Br.	PP	C3	4,5
DS	Stylidium breviscapum R. Br. var. A (KRN 9763)	A3	3
AS	Stylidium calcaratum R. Br.	.PP	C3	6
RP	Stylidium caricifolium Lindley ssp. caricifolium	B1	3
DS	Stylidium dielsianum E. Pritzel	.P	C3	2,5
RP	Stylidium ecorne (F. Muell. ex R. Erickson & J.H. Willis)P.G. Farrel & S.H. James	A2	4
RP	Stylidium hirsutum R. Br.	A2	3
RP	Stylidium macranthum Carlg.	.P	C3	C3	3
RP	Stylidium piliferum R. Br. ssp. piliferum	.P	C3	B2	4
DS	Stylidium repens R. Br. var. repens	A1	4
RP	Stylidium schoenoides DC.	.P	B1	4
RP	Stylidium sp. A (MAB 4311)	.P	A1	1
345 ASTERACEAE												
AS	Actinobole uliginosum (A. Gray)H. Eichler	.AP	C2	..	B2	D3	6
AS	Angianthus conocephalus (J. Black)P.S. Short	.PP	B2	C3	6
AS	Angianthus preissianus (Steetz)Benth.	.P	A4	6
AS	Angianthus tomentosus Wendl.	B3	6
AS	*Arctotheca calendula (L.)Levyns	.P	A1	..	B1	..	A2	7
AS	Asteridea asteroides (Turcz.)G. Kroner	A1	4
AS	Asteridea atrixioides (Sonder & F. Muell.) G. Kroner	.A	C3	6
AS	Blennospora drummondii A. Gray	PP	B2	6
AS	Brachycome ciliaris (Labill.)Less. var. ciliaris	P.P	D3	6
DS	Brachycome ciliaris (Labill.)Less. var. lanuginosa (Steetz)Benth.	A1	6
AS	Brachycome exilis Sonder	.P	D3	6
AS	Brachycome goniocarpa Sonder & F. Muell. ex Sonder	A2	..	A2	6
AS	Brachycome lineariloba (DC.)Druce	.P	B2	6
AS	Brachycome perpusilla (Steetz)J. Black var. perpusilla	.PP	C3	D3	..	6
AS	Brachycome pusilla Steetz	.PA	C3	D3	..	4,5
AS	Calotis hispidula (F. Muell.)F. Muell.	P.	B4	6
AS	Chrysocoryne pusilla (Benth.)Endl.	PAA	B2	6
AS	Chrysocoryne uniflora Turcz.	P.	A2	4
AS	*Cirsium vulgare (Savi)Ten.	A2	7
AS	*Conyza bonariensis (L.)Cronq.	.P	A3	7
AS	Cotula cotuloides (Steetz)Druce	.P	B3	4,5
SS	Cratystylis conocephala (F. Muell.)S. Moore	.A	B4	6
AS	*Crepis capillaris (L.)Wallr.	.P	A2	7
AS	Elachanthus pusillus F. Muell.	...	3K	D3	6
AS	Gnaphalium indutum J.D. Hook.	.P	3KC	B3	6
AS	*Gnaphalium pennsylvanicum Willd.	.P	A2	7
AS	Gnephosis brevifolia (A. Gray)Benth.	.P	A2	A2	..	3
AS	Gnephosis pygmaea (A. Gray)Benth.	.P	D3	2,5
DS	Helichrysum lepidophyllum (Steetz)Benth.	PP	B2	C2	..	4
AS	Helichrysum leucopsidium DC.	.P	A2	A2	6
DS	Helichrysum obtusifolium F. Muell. & Sonder ex Sonder	.P	D3	..	C2	6
DS	Helichrysum aff. obtusifolium F. Muell. & Sonder ex Sonder (MAB 4308)	B2	1
SS	Helichrysum occidentale N. Burb.	.P	A2	2
AS	Helipterum demissum (A. Gray)Druce	A2	6
AS	Helipterum manglesii (Lindley)F. Muell. ex Benth.	.A	C3	4,5
AS	Helipterum pygmaeum (DC.)Benth.	P.P	C2	..	B3	B3	..	B3	..	4,5
AS	Helipterum pyrethrum (Steetz)Benth.	...	3K	A2	4
AS	Helipterum tenellum Turcz.	P.P	B4	6
AS	Hyalochlamys globifera A. Gray	P.	B3	2,5

Appendix I (continued). Flora List

LF		RES		LANDFORM								DIS
		FPS	CS	B	D	F1	F2	G	L	M	S	
AS	*Hypochaeris glabra L.	..A	D2	D2	C2	D2	E2	7
AS	Millotia myosotidifolia (Benth.)Steetz	B2	6
AS	Millotia tenuifolia Cass. var. tenuifolia	PAP	C3	..	B2	6
AS	Myriocephalus rhizocephalus (DC.)Benth.	A2	6
DS	Olearia ciliata (Benth.)F. Muell. ex Benth. var. ciliata	..P	C2	..	B1	6
DS	Olearia exiguifolia (F. Muell.)F. Muell. ex Benth.	C2	6
DS	Olearia homolepis (F. Muell.)F. Muell. ex Benth.	B1	2,5
DS	Olearia muelleri (Sonder)Benth.	ANP	D4	6
DS	Olearia muricata (Steetz)Benth.	B2	4
DS	Olearia aff. muricata (Steetz)Benth. (MAB 1422)	A2	1
DS	Olearia ramosissima (DC.)Benth.	P.	C3	2,5
SS	Olearia revoluta F. Muell. ex Benth.	PA.	B2	B2	..	A2	4,5
DS	Olearia sp. A (MAB 1792)	..	3E	A2	2
AS	Podolepis auriculata DC.	A2	4
HP	Podolepis capillaris (Steetz)Diels	PAP	C2	..	C3	6
AS	Podolepis gracilis (Lehm.)R.A. Graham	A2	4
AS	Podolepis lessonii (Cass.)Benth.	P.P	C1	6
HP	Podolepis microcephala Benth.	..	3R	A2	5
AS	Podotheca angustifolia (Labill.)Less.	PP.	C2	6
AS	Podotheca gnaphalioides R.A. Graham	A2	4,5
AS	Pogonolepis stricta Steetz	..P	B3	B3	5
AS	*Pseudognaphalium luteo-album (L.)Hilliard & B.L. Burtt	B1	..	C1	7
AS	Quinetia urvillei Cass.	P.	D3	6
AS	Rutidosia multiflora (Nees)Robinson	B4	B3	6
AS	Scyphocoronis major (Turcz.)Druce	P.P	B3	O3	6
AS	Senecio glossanthus (Sonder)Belcher	PAP	C2	..	C2	A2	6
AS	Senecio lautus G. Forster ex Willd. ssp. dissectifolius Ali	C3	6
AS	Senecio quadridentatus Labill.	PP.	B1	B1	6
AS	Siloxerus pygmaeus (A. Gray)P.S. Short	..P	B2	D3	4,5
AS	*Sonchus oleraceus L.	..P	C1	..	G1	B1	7
AS	*Ursinea anthemoides (L.)Poiret	..P.	D3	7
DS	Vittadinia australasica (Turcz.)N. Burb. var. australasica	B2	6
DS	Vittadinia dissecta (Benth.)N. Burb. var. hirta N. Burb.	B2	4,5
AS	Waitzia aurea (Benth.)Steetz	PPP	D3	D3	4,5
AS	Waitzia citrina (Benth.)Steetz	..P	D3	B2	6
AS	Waitzia paniculata (Steetz)F. Muell. ex Benth.	B2	B3	4
AS	Genus indet. A (KRN 8559)	..PP	3KC	B3	2,5

Appendix 2. Taxa with important conservation values

This list highlights those taxa in Appendix 1 requiring implementation of conservation measures to ensure their survival. Some taxa listed require additional surveys to better understand their distribution and biology. Conservation status codes follow Leigh *et al.* (1981) - see Appendix 1.

Taxon	Endangered	Vulnerable	Rare	Poorly known
Acacia aff. bidentata B	.	2VCP	.	.
Acacia aff. cochlearis	.	.	.	2K
Acacia crassuloides	.	3VCA	.	.
Acacia excentrica	.	.	.	3K
Acacia pritzeliana	.	.	3RCP	.
Acacia sorophylla	.	3VCP	.	.
Acacia sp. B	.	2V	.	.
Acacia sp. E	.	3VCP	.	.
Acacia sp. I	.	2VCP	.	.
Acacia sp. K	.	.	3R	.
Acacia sp. N	.	1V	.	.
Acacia sp. O	2E	.	.	.
Acacia sp. P	3E	.	.	.
Acacia sp. R	1E	.	.	.
Acacia sp. U	.	.	.	2KCP
Acacia sp. V	.	.	.	3KCP
Acacia sp. W	2E	.	.	.
Acacia sp. X	1E	.	.	.
Adenanthos ileticus	2E	.	.	.
Andersonia macranthera	2EC	.	.	.
Angasomyrtus salina	.	2VCP	.	.
Aotus sp. A	1EC	.	.	.
Asteraceae genus indet. A	.	.	.	3KCP
Baeckea crassifolia var. isosandra	.	.	3RC	.
Banksia blechnifolia	.	3VCP	.	.
Banksia pilostylis	.	3VCP	.	.
Beaufortia aff. empetrifolia	.	3V	.	.
Billardiera mollis	.	2V	.	.
Boronia aff. fabianoides	.	3V	.	.
Boronia oxyantha var. brevicalyx	.	.	3RC	.
Brachyloma sp. A	.	.	.	3K
Cassytha micrantha	.	3VCP	.	.
Chamelaucium sp. A	.	2V	.	.
Chorizema sp. A	.	2V	.	.
Comesperma acerosum	.	3V	.	.
Conostephium drummondii	.	.	3RCP	.
Conostephium sp. A	.	.	.	3K
Conostylis phathyrantha	.	.	3R	.
Conostylis sp. A	2E	.	.	.
Cypselocarpus haloragoides	.	3VC	.	.
Dampiera carinata	.	.	3RCP	.
Dampiera trigona	.	.	3R	.
Darwinia sp. F	1EC	.	.	.
Darwinia sp. G	1E	.	.	.
Daviesia sp. B	.	.	.	3KCP
Daviesia sp. C	1EC	.	.	.
Daviesia sp. E	.	3VCP	.	.
Dillwynia acerosa	.	.	.	3K
Dillwynia divaricata	.	.	.	3K
Dodonaea glandulosa	.	2VCP	.	.
Drosera sp. A	.	.	3R	.
Dryandra aff. pteridifolia	.	3VCP	.	.
Elachanthus pusillus	.	.	.	3K

Appendix 2 (continued). Taxa with important conservation values

Taxon	Endangered	Vulnerable	Rare	Poorly known
<i>Eremophila denticulata</i>	.	.	3RCP	.
<i>Eremophila dichroantha</i>	.	.	3RCA	.
<i>Eremophila serpens</i>	.	2V	.	.
<i>Eremophila aff. serpens</i>	3E	.	.	.
<i>Eriostemon fitzgeraldii</i>	.	.	3RCP	.
<i>Eriostemon aff. thryptomenoides</i>	2E	.	.	.
<i>Eucalyptus angustissima</i>	.	.	2RC	.
<i>Eucalyptus aff. angustissima</i>	2E	.	.	.
<i>Eucalyptus deflexa</i>	.	3VCP	.	.
<i>Eucalyptus dielsii</i>	.	3VCP	.	.
<i>Eucalyptus aff. diptera</i> A	.	.	2R	.
<i>Eucalyptus discreta</i>	.	.	3RC	.
<i>Eucalyptus forrestiana</i> ssp. <i>forrestiana</i>	.	3VCA	.	.
<i>Eucalyptus forrestiana</i> ssp. <i>dolichorhyncha</i>	2ECP	.	.	.
<i>Eucalyptus forrestiana</i> ssp. <i>stoatei</i>	2E	.	.	.
<i>Eucalyptus halophila</i>	.	2VCP	.	.
<i>Eucalyptus merrickiae</i>	.	3V	.	.
<i>Eucalyptus aff. micranthera</i>	.	.	.	2K
<i>Eucalyptus nutans</i>	.	.	2RC	.
<i>Eucalyptus ovularis</i>	.	3VCP	.	.
<i>Eucalyptus</i> sp. A	.	2V	.	.
<i>Eucalyptus</i> sp. B	1E	.	.	.
<i>Eucalyptus</i> sp. H	2E	.	.	.
<i>Eucalyptus</i> sp. I	1E	.	.	.
<i>Eucalyptus</i> sp. J	1E	.	.	.
<i>Gahnia</i> sp. A	.	.	.	2K
<i>Glyceria fluitans</i>	.	2V	.	.
<i>Gnaphalium indutum</i>	.	.	.	3KCP
<i>Gonocarpus trichostachyus</i>	.	.	.	3K
<i>Goodenia trichophylla</i>	.	.	3RCP	.
<i>Grevillea aneura</i>	.	3VCP	.	.
<i>Gyrostemon ditrigynus</i>	2E	.	.	.
<i>Gyrostemon prostratus</i>	3E	.	.	.
<i>Hakea</i> sp. A	1E	.	.	.
<i>Halgania</i> sp. A	.	2V	.	.
<i>Haloragis</i> sp. A	1E	.	.	.
<i>Heliotropium</i> sp. A	.	.	2R	.
<i>Helipterum pyrethrum</i>	.	.	.	3K
<i>Hemigenia aff. eutaxioides</i>	.	.	.	3K
<i>Hibbertia andrewsiana</i>	.	.	3RC	.
<i>Hydrocotyle medicaginoidea</i>	.	.	3RCP	.
<i>Hypoxis</i> sp. A	.	2V	.	.
<i>Isolepis</i> sp. A	2E	.	.	.
<i>Isopogon tridens</i>	.	3V	.	.
<i>Kennedia beckxiana</i>	.	.	2R	.
<i>Kunzea baxteri</i>	.	.	3RCP	.
<i>Lawrencia diffusa</i>	.	3VCP	.	.
<i>Lechenaultia</i> sp. A	.	3V	.	.
<i>Lepidosperma</i> sp. B	.	.	.	3K
<i>Leptospermum maxwellii</i>	.	.	3R	.
<i>Leucopogon aff. allittii</i>	2E	.	.	.
<i>Leucopogon bossiaea</i>	.	.	.	3KCP
<i>Leucopogon brevicuspis</i>	.	.	.	3KCP
<i>Leucopogon opposens</i>	.	.	3R	.
<i>Leucopogon woodsii</i>	.	.	3RC	.
<i>Leucopogon</i> sp. E	1E	.	.	.
<i>Leucopogon</i> sp. G	1E	.	.	.

Appendix 2 (continued). Taxa with important conservation values

Taxon	Endangered	Vulnerable	Rare	Poorly known
<i>Leucopogon</i> sp. I	1E	.	.	.
<i>Leucopogon</i> sp. M	.	1V	.	.
<i>Leucopogon</i> sp. N	.	3V	.	.
<i>Melaleuca cliffortioides</i>	.	.	3RCP	.
<i>Melaleuca aff. nesophila</i>	.	.	3RCA	.
<i>Melaleuca sparsiflora</i>	.	.	.	2K
<i>Melaleuca</i> sp. C	.	.	.	3KCP
<i>Melaleuca</i> sp. N	1E	.	.	.
<i>Microcorys virgata</i>	.	.	.	3K
<i>Myoporum beckeri</i>	.	.	.	2KCP
Myoporaceae genus indet.	2E	.	.	.
<i>Olearia</i> sp. A	3E	.	.	.
<i>Persoonia tortifolia</i>	.	.	.	2K
<i>Persoonia</i> sp. A	.	.	3RCP	.
<i>Podolepis microcephala</i>	.	.	3R	.
<i>Pomaderris intangenda</i>	.	3V	.	.
<i>Prostanthera</i> sp. A	1E	.	.	.
<i>Pultenaea arida</i>	.	.	3RCA	.
<i>Pultenaea</i> sp. B	.	2VCP	.	.
<i>Pultenaea</i> sp. C	.	3V	.	.
<i>Pultenaea</i> sp. D	.	2V	.	.
<i>Pultenaea</i> sp. E	1E	.	.	.
<i>Pultenaea</i> sp. H	1E	.	.	.
<i>Ricinocarpos trichophorus</i>	.	.	2VCP	.
<i>Scaevola</i> sp. A	1E	.	.	.
<i>Schoenus caespitius</i>	.	.	3RCP	.
<i>Schoenus</i> sp. A	.	.	.	3K
<i>Siegfriedia darwinioides</i>	.	3VC	.	.
<i>Spyridium oligocephalum</i>	.	.	3RCP	.
<i>Spyridium</i> sp. A	.	.	.	3K
<i>Spyridium</i> sp. B	.	.	.	3KC
<i>Stachystemon</i> sp. A	1E	.	.	.
<i>Thysanotus parviflorus</i>	.	3VCP	.	.
<i>Trachymene croniniana</i>	.	3V	.	.
<i>Triglochin muelleri</i>	.	.	.	3K
<i>Triodia concinna</i>	.	.	.	3K
<i>Verticordia aff. drummondii</i>	.	.	3RCP	.
<i>Wilsonia rotundifolia</i>	.	.	3RCP	.
<i>Wurmbea sinora</i>	.	.	.	3K