

Floristic communities of the Ravensthorpe Range, Western Australia

ADRIENNE MARKEY¹, STEPHEN KERN^{2,3} AND NEIL GIBSON¹

¹ Science Division, Department of Environment and Conservation, Locked Bag 104, Bentley Delivery Centre, Western Australia 6983. Email: Adrienne.Markey@dec.wa.gov.au; Neil.Gibson@dec.wa.gov.au

² Western Botanical, PO Box 3393, Bassendean, WA, 6842.

³ current contact email: stephenkern@hotmail.com

ABSTRACT

The Ravensthorpe Range consists of a series of ridges and peaks located in the greater Fitzgerald Biosphere reserve, in south-west Western Australia near the township of Ravensthorpe. A floristic classification of the vegetation of the range is described and compared to recent vegetation mapping undertaken in the area. The floristic classification was derived from an analysis of 580 taxa from 265 plots that were located to cover the topographic and geographic extent of the range. Twenty-one communities are described, and these show broad agreement with the recent detailed mapping, except for one widespread upland unit that was floristically heterogeneous. The floristic communities correlated with site physical attributes, notably topographic position, substrate and altitude. In addition to the high beta diversity, the Ravensthorpe Range has a large number of geographically restricted species, species listed as threatened and species being considered for listing. Despite the high conservation values of the range, reservation is limited to two small A-class nature reserves off the main range. These reserves are not representative of the full diversity of flora and communities on the range. Mineral exploration and mining are currently active or proposed for parts of the range, and mining tenements cover most of the area of the range. The impacts of exploration, mining and *Phytophthora* present significant challenges to managing this major biodiversity hotspot.

Keywords: endemic, threatened taxa, beta diversity, classification

INTRODUCTION

The Yilgarn Craton of Western Australia is composed of ancient granitoid gneisses, within which are inter-bedded extensive belts of metamorphosed sedimentary and volcanic rocks. These are referred to locally as 'greenstone' belts, which manifest as hills and ranges when exposed above the surrounding plains. Recent surveys of these landforms have found them to be repositories of endemic and rare plant taxa (Gibson et al. 2007, 2010, 2011), and short range endemic invertebrate species (i.e. species with total range of less than 10,000 km²; Harvey 2002; Edward & Harvey 2010). They also support floristic communities which differ in composition to both the surrounding lowlands and adjacent ranges (Gibson et al. 2007, 2010, 2011).

The Ravensthorpe Range is one such greenstone landform located on the Yilgarn Craton, in the far south-western corner of Western Australia (Fig. 1) and within the South West Australian Floristic Region (SWAFR, sensu Hopper & Gioia 2004). This region is known for its exceptional biological diversity, and is considered to be one of 34 global biodiversity hotspots because of a

combination of high species richness, endemism and habitat loss (Myers et al. 2000; Myers 2003). The Ravensthorpe Range occurs within a node of high species diversity and endemism within the SWAFR (Hopper & Gioia 2004), and is part of the United Nations Educational, Scientific and Cultural Organisation's Fitzgerald River Biosphere Reserve (Chapman & Newbey 1995). Because of its location, the Ravensthorpe Range is recognised as a significant biological corridor between the Fitzgerald River National Park and the Great Western Woodlands (Chapman & Newbey 1995; Watson 1991; Watson & Wilkins 1999). The range is noted for its high biodiversity values (Craig et al. 2008; Moir et al. 2009; Edward & Harvey 2010; McDonald 2010), including being important habitat for several threatened species of vertebrates (Chapman & Newbey 1995; Harris et al. 2008). The most recent census of vascular plants recorded 1324 native species (Craig 2008) from Beard's Ravensthorpe vegetation system (Beard 1972, 1973), which covers both the range and areas of mafic geology immediately to the south-west. At least 58 taxa are endemic to, or largely restricted to, this vegetation system (Craig 2008; Kern et al. 2008; the Western Australian Herbarium 1998–). Some 76 taxa in the Ravensthorpe vegetation system are listed as being of conservation significance

(Craig 2008; Smith 2010), and 60 of these are known to occur on the range (Harris et al. 2008; Western Australian Herbarium 1998–). Five ecological communities of conservation significance are listed for the range (Craig et al. 2008; Kern et al. 2008; Department of Environment and Conservation 2010). Like many greenstone ranges on the Yilgarn Craton, the Ravensthorpe Range is rich in mineral resources and of interest to the resources sector (Witt 1998).

STUDY SITE

The Ravensthorpe Range is an extensive outcropping of greenstone bedrock located 550 km south-east of Perth, Western Australia. The main range is a narrow, linear feature c. 50 km in length trending north-west to south-east, and up to 7 km wide. The centre of the range is 7 km east of the town of Ravensthorpe and the southern edge 25 km from the coastal township of Hopetoun. There are six peaks named on the main range; Mt Short is the highest peak (453 m), followed by Mt Benson (404 m),

Mt Desmond (340 m), Mt Chester (300m), Mt McMahon (274 m) and Overshot Hill (377 m; Fig. 1). Immediately east of the main ridges are a series of low, undulating pediments and hills. Isolated from the main range, Bandalup Hill (228 m) is a prominent feature 34 km east of Ravensthorpe township and on the eastern margin of the survey area (Fig. 1). The main range is relatively low in elevation, being only, at the most, c. 150 m above the surrounding plains. Numerous drainage lines and several ephemeral creeks drain from the ranges into the Steere and Jerdacuttup Rivers, which incise the central and southern portion of the range. These two main rivers are saline and flow only after winter rainfall, contracting to a series of permanent pools during the summer months.

Climate

The Ravensthorpe Range is located near the edge of the transitional rainfall zone of the SWAFR, and the area has a warm Mediterranean climate, with cool and wet winters and warm to hot summers (Harris et al. 2008; Beard 1990). The climate is moderated by proximity to the




Figure 1. Ravensthorpe Range survey area showing major landmarks and the locations of the 266 permanent plots. Darker shading indicates upland areas.

Southern Ocean, and rainfall is predominately derived from southerly fronts that occur during the winter months (Craig et al. 2008; Chapman & Newbey 1995). However, summer rainfall can originate from the north as remnants of tropical cyclones. On very rare occasions snowfalls have been reported on the range (Chapman & Newbey 1995). The nearest meteorological station is Ravensthorpe township (1901–2007), which has an annual average rainfall of 426 mm (Bureau of Meteorology 1908–). Data from average annual rainfall grids indicate that rainfall across the range declines from 523 to 391 mm with distance from the coastline (Harris et al. 2008).

Geology

The Ravensthorpe Range is essentially a linear outcropping of a complex of Archean (c. 3 Ga) metavolcanic (mafic and ultramafic) and metasedimentary bedrock, which is inter-bedded within the granitoid gneiss of the southern Yilgarn Craton (Witt 1997, 1998). These metasediments and metavolcanics are faulted and folded to produce a sharp ridge and peaks, the uplands of which are covered by an undulating upland of Quaternary laterites or silcretes, produced from *in situ* weathering. The eroding margins of this caprock give way to colluvial slopes with some outcropping bedrock and deep valleys incised into steep slopes. Surrounding the Ravensthorpe Range are Tertiary sediments, colluvium and sand that overlie the Archean granitoid gneiss, forming extensive, low-lying plains and alluvial flats interrupted by outcrops and hills. The nearest significant granite-quartzite hills are associated with the Barren and Eyre Ranges, the closest peaks being 25–40 km south of Ravensthorpe.

The geology of the Ravensthorpe greenstone belt and associated Ravensthorpe Range has been described and mapped at a scale of 1:100,000 (Witt 1996). Witt (1996, 1997) described three geological terranes that cover the Ravensthorpe Range survey area, of which the Carlingup Terrane is the main tectonic unit. The Ravensthorpe Terrane abuts onto the western margin of the Carlingup Terrane, and the Cocanarup greenstones underlie part of Mt Short. The Carlingup Terrane is composed of four main stratigraphic units: the Bandalup Ultramafics, the Chester Formation, Maydon Basalt and the Hatfield Formation. The Bandalup Ultramafics (metamorphosed undivided ultramafic rocks) occur on lower slopes and undulating low hills, while the Maydon Basalt (metamorphosed undivided mafic rocks) dominates low hills between the main range and Bandalup Hill. The Hatfield Formation (pelite, psammite, metamorphic and felsic volcanic rocks) lies on the eastern margin of main ridge. It is the durable, erosion-resistant Chester Formation (pelite, psammite and metamorphosed sedimentary rocks) which forms the main ridges of the range (Witt 1996, 1997).

Soils of the Ravensthorpe Range reflect underlying lithology and topography, with skeletal to shallow soils being formed *in situ* on upland laterites and bedrock, colluvium being deposited on hill slopes and foot slopes, and alluvial deposits accumulating in drainage features and

alluvial flats at the base of the range. Associated with the mafic and ultramafic lithologies are mineral-rich clays (Beard 1973; Bennett 1987; Chapman & Newbey 1995).

Vegetation

The Ravensthorpe Range is situated within the Esperance Plains Bioregion (Department of Environment and Water Resources 2007). The vegetation was first described by Beard (1972, 1973) as part of a wider survey of the Ravensthorpe region. According to these publications, three vegetation systems occur on the range (Ravensthorpe, Oldfield and Esperance Systems). The Ravensthorpe System covers most of the range and wider surrounds to the south-west and east. Beard recognised eight physiognomic communities on the range. These communities are associated with the topographic and soil catenary sequence on the range and fall into three categories: 'Thickets' on summit ridges; 'Mallee' on pediments and low hills; and 'Sclerophyll Woodland' in deep, broad valleys. Bennett (1987) and Chapman and Newbey (1995) subsequently focussed on sections of the range, providing greater floristic and structural detail for these sections and describing their associations with edaphic and topographic factors. Targeted surveys on other parts of the range have been undertaken for environment impact assessments (e.g. Cockerton & Craig 2000; Craig 1999, 2004, 2005). More recently, detailed vegetation mapping of the main range between Mt Short and Kundip has described 70 vegetation units based on structure and composition (Craig et al. 2008).

Land use

Over the past century the Ravensthorpe region has been an agricultural and mining centre, and these activities have had impacts upon the vegetation of the range. The region lies in the south-east of the wheatbelt agricultural zone, and a significant proportion of the woodlands around the range have been cleared or cut-over. Wheatfields abut the base of the northern and western foothills, taking advantage of the heavy, rich soils derived from the mafic bedrock (Beard 1973).

The Ravensthorpe greenstone belt is highly prospective for minerals, and has a long history of exploration and mining for base and precious minerals, notably gold, copper, nickel and silver (Witt 1998). The original mining centre was established in the 1890s at the now abandoned township of Kundip, and several open pits and tailings dumps remain in the southern section of the range from these and more recent mining activities (Harris et al. 2008). Major operations to exploit significant nickel laterite and magnesite deposits at Bandalup Hill commenced in January 2008, and a lithium/tantalum mine immediately north of Ravensthorpe township (Mt Cattlin) was approved in 2009. Exploration activities are in progress at various locations over the range, and active or pending mining tenements cover almost the entire extent (Harris et al. 2008). Although not in the tenure of the Department of Environment and Conservation (DEC),

the Department manages these areas and conducts prescribed burns and feral animal baiting programs. Most of the range is Unallocated Crown Land, with the Overshot Hill and Kundip Nature Reserves being the only parts of the range within the conservation estate (both A-class nature reserves; Fig. 1).

Objectives

The aim of this study was to produce a classification of the vegetation communities on the Ravensthorpe Range based on floristic composition, in order to allow the regional context of proposed developments to be determined in a consistent and repeatable manner. These data were also used to examine compositional heterogeneity in some of the 70 mapping units that cover the northern sections of the range (Craig et al. 2008) and to assess the relationship between the vegetation mapping and the floristic classification.

METHODS

Two hundred permanently marked, georeferenced plots were established by a team of consultants (Western Botanical Pty. Ltd.) and rescored over a period of several months in 2007. A further 66 plots were established by the DEC in the following late spring and summer of 2008–09 (Fig. 1). Plots were placed to cover the geographical and topographical extent of the range in order to capture the widest range of floristic variation. No recently burnt vegetation was sampled in order to minimise the inclusion of fire ephemerals in the dataset. For each plot, location (to within an accuracy of 4 m) and altitude were determined using a GPS (Garmin, GPS76). The occurrence of all vascular species was recorded in each 10 × 10 m plot. Each of these plots was nested within a larger 20 × 20 m plot that was used to record canopy species (trees, mallee shrubs >3 m). Plot size was based on species-area curves from Bennett (1987) and sampling methodology follows Gibson et al. (2004). Taxa occurring within and adjacent to plots were collected for identification at the Western Australian Herbarium (PERTH), where over 1500 voucher specimens have been lodged. Further information on taxa was obtained from online records of the Western Australian Herbarium (1998–).

Data analyses were performed using routines available in PRIMER-E and PERMANOVA+ (PRIMER version 6, Clarke & Gorley 2006). Presence/absence data were analysed using Bray & Curtis association measures. This measure has proved extremely robust in the analysis of ecological data (Faith et al. 1987). Classifications of both the plots and species were performed using UPGMA clustering (Sneath & Sokal 1973). Similarity profiles (SIMPROF) and indicator species analysis (INDVAL) were used to examine the classification structure (Dufrene & Legendre 1997; Clarke et al. 2008). Floristic nomenclature follows that of the Western Australian Herbarium (1998–).

A preliminary analysis was undertaken on 593 taxa

identified from within the 266 plots. The subsequent analysis was performed on 580 taxa from 265 plots, after one outlying plot and its singleton taxa were omitted. Hybrid taxa were also omitted and several closely-related or intergrading pairs of taxa were amalgamated at a higher taxonomic level, mostly from lower ranks to species level (*Hybanthus floribundus*, *Hakea pandanicarpa*, *Eucalyptus astringens*, *Melaleuca pentagona*, *Acacia sulcata*, *Pultenaea indira* and *Schoenus sublaxus*) or a species complex later resolved to be separate taxa (*Leucopogon* sp. Coujinup [MA Burgman 1085] and *Leucopogon* sp. Newdegate [M Hislop 3585], and *Lysinema ciliatum* and *Lysinema pentapetalum*). Preliminary analyses showed that these amalgamations had little effect on the classifications.

For each community a list of diagnostic, common, short-range endemic and conservation taxa was compiled and a photograph of a plot representative of each community is provided. These photos were taken along plot diagonals. Diagnostic taxa indicative of a community type were identified using indicator species analysis (INDVAL), implemented in PC-ORD (Dufrêne & Legendre 1997; McCune & Mefford 1999). Indicator values were derived from species fidelity and constancy within each community, where a value of 100 indicated that a species was present in all plots of a community and absent from all other plots. Diagnostic species were defined as those with INDVAL values 14 or greater. Common taxa were those that occurred at frequencies of ≥25% of plots in a community. Short-range endemic (SRE) taxa were defined as those that had a total range of less than 10,000 km². Four categories are listed: 1, taxa with ranges <10 km²; 2, taxa with ranges 10–100 km²; 3, taxa with ranges 100–1,000 km²; 4, taxa with ranges 1,000–10,000 km². Convex hulls were calculated from distributional data downloaded from the Western Australian Herbarium, which include our vouchers (accessed 22nd November 2010). Conservation taxa listing follows Smith (2010) and recent updates as shown on Florabase (Western Australian Herbarium 1998–): R = threatened taxa formally listed under Western Australian legislation; P1–P4 = taxa being considered for listing (codes follow Smith 2010).

Typical soils (McDonald et al. 1998) and the number of plots in each of the vegetation map units of Craig et al. (2008) were also listed for each community. Vegetation map units were determined by intersecting the vegetation map with plot location. Some 166 of the 266 plots fell within the mapped area; seven of these were in mosaic units that were not included in the community descriptions. Plots were located in 40 of the 70 vegetation units recognized by Craig et al. (2008), with frequencies ranging from 1 to 44 plots per vegetation unit.

RESULTS

Flora

A total of 697 taxa (species, subspecies and varieties, including putative new taxa and nine new putative hybrid

or intergrade taxa) were recorded within and adjacent to the 266 plots (Appendix 1). Of these, six are listed as rare and 37 are on DEC's priority list (Appendix 1). Six species are naturalised herbaceous weed taxa (*Asphodelus fistulosus*, *Avena fatua*, *Lepidium africanum*, *Sonchus oleraceus*, *Lolium perenne* and *Asparagus asparagoides*). The highest species numbers were from Myrtaceae (136 taxa), Fabaceae (112; 64 Papilionoidea), Proteaceae (69), Cyperaceae (53), Ericaceae (28) and Goodeniaceae (23). The most species-rich genera were *Eucalyptus* (45 taxa and 8 hybrids), *Acacia* (46), *Melaleuca* (35), *Lepidosperma* (33), *Hakea* (19), *Grevillea* (17), *Daviesia* (14 and 1 hybrid), *Boronia* (13), *Schoenus* (14) and *Hibbertia* (11).

Craig (2008) listed 55 taxa believed to be endemic (99–100% of known populations) or almost endemic (80–99% of known populations) to Beard's (1972, 1973) Ravensthorpe vegetation system that covers most of the range as well as extensive areas to the south-west. These definitions are somewhat arbitrary and rely on boundaries defined at 1:250,000-scale mapping. In contrast, we defined short-range endemic taxa as those taxa recorded

in our survey that had a total range of less than 10,000 km², following definitions used for invertebrate taxa (Harvey 2002). Some very restricted taxa fall outside this definition, for example taxa with a small fragmented population restricted to rare habitats but with a total range of >10,000 km² (e.g. *Tetrapetra planata*—total range 10,072.5 km²). Sixty-four SREs were found, four restricted to a range of less than 10 km², and a further seven with ranges of less than 100 km². These taxa were not restricted to particular families but primarily represented shrubs, with a minor occurrence of eucalypt mallees, trees and mallets (Table 1).

Classification

All but two of the 265 plots were classified into 21 groups (subsequently referred to as communities; Fig. 2). Groups have been resolved at variable similarity levels, and decisions on selecting groups at particular levels have been based on branch support from SIMPER analysis, from patterns in the sorted data table (Appendix 2), from indicator species analysis (Appendix 3) and from field

Table 1

Short-range endemic taxa (SRE) recorded on the Ravensthorpe Range. SRE were defined as those taxa that had a total range less than 10,000 km². Four categories are listed: 1 = taxa with ranges <10 km²; 2 = taxa with ranges 10–100 km²; 3 = taxa with ranges 100–1000 km²; and 4 = taxa with ranges 1000–10,000 km².

Family	Taxon	SRE code	Family	Taxon	SRE code
Casuarinaceae	<i>Allocasuarina hystricosa</i>	4		<i>Eucalyptus gardneri</i> subsp. <i>ravensthorpensis</i>	3
Dilleniaceae	<i>Hibbertia abyssa</i>	1		<i>Eucalyptus megacornuta</i>	4
	<i>Hibbertia atrichosepala</i>	1		<i>Eucalyptus oleosa</i> subsp. <i>corvina</i>	4
Droseraceae	<i>Drosera grievei</i>	4		<i>Eucalyptus purpurata</i>	2
Ericaceae	<i>Acotricha orbicularis</i>	1		<i>Eucalyptus stoatei</i>	4
	<i>Leucopogon infuscatus</i>	3		<i>Eucalyptus x bennettiae</i>	3
	<i>Lissanthe pleurandrodes</i>	4		<i>Kunzea cincinnata</i>	3
Euphorbiaceae	<i>Beyeria cockertonii</i>	2		<i>Kunzea similis</i> subsp. <i>mediterranea</i>	1
	<i>Beyeria villosa</i>	3		<i>Leptospermum</i> sp. Bandalup Hill (G Cockerton 11001)	4
Fabaceae	<i>Acacia bifaria</i>	4		<i>Melaleuca penicula</i>	4
	<i>Acacia disticha</i>	4		<i>Melaleuca pomphostoma</i>	4
	<i>Acacia durabilis</i>	4		<i>Melaleuca stramentosa</i>	3
	<i>Acacia heterochroa</i> subsp. <i>heterochroa</i>	4		<i>Melaleuca ulicoides</i>	3
	<i>Acacia loricina</i> var. <i>crassifolia</i>	4		<i>Micromyrtus navicularis</i>	4
	<i>Acacia ophiolithica</i>	4	Pittosporaceae	<i>Marianthus mollis</i>	3
	<i>Acacia pinguiculosa</i> subsp. <i>pinguiculosa</i>	4	Proteaceae	<i>Adenanthes glabrescens</i> subsp. <i>exasperatus</i>	4
	<i>Acacia pusilla</i>	4		<i>Banksia coriijuga</i>	2
	<i>Acacia</i> sp. Ravensthorpe Range (BR Maslin 5463)	3		<i>Banksia foliosissima</i>	4
	<i>Daviesia megacalyx</i>	2		<i>Banksia laevigata</i> subsp. <i>laevigata</i>	4
	<i>Pultenaea calycina</i> subsp. <i>proxena</i>	3		<i>Grevillea fulgens</i>	4
	<i>Pultenaea craigiana</i>	3		<i>Grevillea patentiloba</i> subsp. <i>platypoda</i>	4
	<i>Pultenaea indira</i> subsp. <i>monstrosita</i>	4		<i>Grevillea punctata</i>	3
	<i>Pultenaea wudjariensis</i>	2		<i>Grevillea rigida</i> subsp. <i>distans</i>	4
				<i>Grevillea sulcata</i>	3
Goodeniaceae	<i>Dampiera deltoidea</i>	4		<i>Isopogon</i> sp. Ravensthorpe (DB Foreman 1207)	3
	<i>Goodenia phillipsiae</i>	3	Rhamnaceae	<i>Spyridium glaucum</i>	3
	<i>Goodenia stenophylla</i>	3	Rutaceae	<i>Boronia oxyantha</i> var. <i>brevicalyx</i>	4
Lamiaceae	<i>Microcorys pimeleoides</i>	3		<i>Boronia ternata</i> var. <i>elongata</i>	4
Malvaceae	<i>Guichenotia anota</i>	3		<i>Microcybe pauciflora</i> subsp. <i>grandis</i>	2
	<i>Guichenotia apetala</i>	3	Sapindaceae	<i>Dodonaea trifida</i>	4
Myrtaceae	<i>Calothamnus roseus</i>	2	Scrophulariaceae	<i>Eremophila densifolia</i> subsp. <i>erecta</i>	4
	<i>Eucalyptus cernua</i>	4		<i>Thymelaeaceae</i>	
	<i>Eucalyptus desmondensis</i>	3		<i>Pimelea physodes</i>	4




Figure 2. Summary dendrogram of the 21 communities found on the Ravensthorpe Range, derived from the classification of presence/absence data of 580 taxa from 265 plots. Three major groups and the two unallocated plots are shown.

observations. At the highest level in the dendrogram, three broad groups were recognised (Fig. 2). The first group (communities 1–7) are predominately woodlands with average species richness per community of 10–27.8 taxa per plot, generally occurring lower in the landscape. The second group (communities 8–12) are mostly mallee shrublands with moderate levels of species richness (14.8–24.5 taxa per plot), found across the landscape; while the third group (communities 13–21) are species-rich mallee shrublands (16.7–42.7 taxa per plot) generally found high in the landscape. Detailed descriptions of each community are given below.

Woodlands group

Community 1: Lowland *Eucalyptus flocktoniae*/*E. phenax*/*E. calycogona* woodlands.

Map unit(s): Eflo/Ephe (4), Egar (2).

This community consisted of woodlands of mallee and mallets, typically dominated by *Eucalyptus flocktoniae* subsp. *flocktoniae*, *E. phenax* subsp. *phenax* and *E. calycogona* subsp. *calycogona*, but also with *E. gardneri* subsp. *ravensthorpensis* and *E. oleosa* subsp. *corvina* as co-dominant canopy species at some sites. The understorey was usually a sparse assemblage of shrubs, grasses and herbs. This community was found around Overshot Hill and in the central part of the range, usually on the gently- to moderately-inclined lower hillslopes of the range.

Soils: dark red-brown clay loam, no exposed bedrock.

No. plots 8; Taxa/plot (SD) 20.4 ± 2.9 ; Altitude 202–284 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus phenax* subsp. *phenax* (19). Shrubs *Acacia* sp. Ravensthorpe Range [BR Maslin

5463] (22), *Enchylaena tomentosa* var. *tomentosa* (16), *Grevillea patentiloba* subsp. *platypoda* (22). Herbs *Austrodanthonia setacea* (17), *Austrostipa acrociliata* (21), *A. hemipogon* (22), *A. sp.* Carlingup Rd [S Kern & R Jasper 184959] (43), *Daucus glochidiatus* (22), *Oxalis perennans* (50), *Ptilotus holosericeus* (63), *Sonchus oleraceus* (20), *Vittadinia gracilis* (25).

Most common taxa (% frequency): Trees *Eucalyptus flocktoniae* subsp. *flocktoniae* (100), *E. phenax* subsp. *phenax* (100), *E. calycogona* subsp. *calycogona* (50), *E. gardneri* subsp. *ravensthorpensis* (25), *E. oleosa* subsp. *corvina* (25), Shrubs *Scenna artemisioides* subsp. *x artemisioides* (75), *Acacia glaucoptera* (63), *A. crinacea* (50), *Gastrolobium parviflorum* (50), *Grevillea patentiloba* subsp. *platypoda* (50), *Enchylaena tomentosa* var. *tomentosa* (38), *Exocarpos aphyllus* (38), *Grevillea huegelii* (38), *Lasiopetalum compactum* (38), *Olearia muelleri* (38), *Acacia* sp. Ravensthorpe Range [BR Maslin 5463] (25), *Acrotriche ramiflora* (25), *Boronia inornata* subsp. *inornata* (25), *Dodonaea pinifolia* (25), *Hakea commutata* (25), *Hibbertia psilocarpa* (25), *Melaleuca hamata* (25), *Santalum acuminatum* (25), *Sclerolaena diacantha* (25). Herbs *Austrodanthonia setacea* (63), *Austrostipa hemipogon* (63), *A. sp.* Carlingup Rd [S Kern & R Jasper 184959] (63), *Ptilotus holosericeus* (63), *Oxalis perennans* (50), *Austrostipa acrociliata* (38), *A. elegantissima* (38), *Billardiera fusiformis* (25), *Daucus glochidiatus* (25), *Lepidosperma* sp. Saltbush Hill [KR Newbey 4118] (25), *Platysace maxwellii* (25), *Sonchus oleraceus* (25), *Vittadinia gracilis* (25).

Short range endemics

100–1000 km²: *Acacia* sp. Ravensthorpe Range (BR Maslin 5463), *Eucalyptus gardneri* subsp. *ravensthorpensis*, *Melaleuca ulicoides*.

1000–10,000 km²: *Acacia bifaria*, *A. pusilla*, *Eucalyptus oleosa* subsp. *corvina*, *Grevillea patentiloba* subsp. *platypoda*.

Conservation taxa: *Acacia bifaria* (P3), *A. sp.* Ravensthorpe Range [BR Maslin 5463] (P1).



Figure 3. Community 1, mallee woodland with sparse understorey (survey plot R216).

Community 2: *Eucalyptus salmonophloia* woodlands on lower hillslopes.

Map unit(s): Esal (3), Eflo/Esgu (1), Egar (1).

Community 2 typically consisted of *Eucalyptus salmonophloia* tall open forest and woodlands, sometimes co-occurring with *E. gardneri* subsp. *ravensthorpensis* or other eucalypt taxa (*E. phenax*, *E. flocktoniae*) and overtops an open to sparse shrubland. This community type was found on Ravensthorpe Range, near the confluence of the Jerdacutup River and Cordingup Creek where a fault bisects the Range.

Soils: generally red-brown sandy loams of varying depth, with limited exposure of outcropping bedrock.

No. plots 5; Taxa/plot (SD) 21.2 ± 2.4 ; Altitude 157–204 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus salmonophloia* (80). Shrubs *Olearia muelleri* (53), *Acacia erinacea* (41), *Rhagodia crassifolia* (31), *Acacia glaucoptera* (24), *Wilsonia humilis* (19). Herbs *Lepidosperma* sp. Mt Chester [S. Kern et al. LCH 16596] (58), *Dianella brevicaulis* (50), *Austrostipa pycnostachya* (54), *A. elegantissima* (33), *A. hemipogon* (20).

Most common taxa (% frequency): Trees *Eucalyptus salmonophloia* (80), *E. phenax* subsp. *phenax* (40). Shrubs *Acacia erinacea* (100), *A. glaucoptera* (100), *Olearia muelleri* (100), *Rhagodia crassifolia* (60), *Wilsonia humilis* (60), *Daviesia nematophylla* (40), *Grevillea pectinata* (40), *Melaleuca hamata* (40), *Santalum acuminatum* (40), *Scilla artemisioides* subsp. x *artemisioides* (40). Herbs *Austrostipa elegantissima* (80),

Lepidosperma sp. Mt Chester [S. Kern et al. LCH 16596] (80), *Austrostipa hemipogon* (60), *A. pycnostachya* (60), *Dianella brevicaulis* (60), *Austrodanthonia setacea* (40), *Cassytha mclantha* (40), *Dianella revoluta* var. *revoluta* (40), *Gahnia ancistrophylla* (40), *Neurachne alopecuroides* (40).

Short range endemics

100–1000 km²: *Eucalyptus gardneri* subsp. *ravensthorpensis*.

Conservation taxa: *Cassinia arcuata* (P2).

Community 3: *Eucalyptus flocktoniae*/*E. phenax*/*E. calycogona*/*E. cernua* woodlands over *Melaleuca* spp.

Map unit(s): Eflo/Ephe (5), Eflo/Mgor (3), Edie (1), Eext (1), Eind/Mpau (1), Epil/Edep (1), Mosaic Mallec/Mund & Mspp (1).

This community type was widespread and typically consisted of tall mallee shrublands and mallet woodlands dominated by mixed eucalypts, notably *Eucalyptus flocktoniae* subsp. *flocktoniae*, *E. phenax* subsp. *phenax* and *E. calycogona* subsp. *calycogona*, also with *E. pileata*, *E. cernua* and *E. platypus* as less common canopy eucalypt species. It had a rich assemblage of *Melaleuca* species (*M. cucullata*, *M. pauperiflora*, *M. cliffortioides*, *M. lateriflora* subsp. *lateriflora*) associated with it, and often one or several species would form a dense shrub mid-stratum. There was some suggestion of internal structure within this community and further sampling may resolve subgroups. Community 3 was found on gently- to moderately-inclined hillslopes, from lower to upper slopes but predominantly on midslopes across the range. There was a tendency for the *M. hamata* dominated sites to occur on mid to upper slopes.



Figure 4. Community 2, tall salmon gum woodland (survey plot R 196).

Soils: predominantly red-brown to brown clays, with some clay loams; exposed shale bedrock was rare.

No. plots 34; Taxa/plot (SD) 19.0 ± 0.8 ; Altitude 75–334 m

Diagnostic taxa (INDVAL): Trees *Eucalyptus calycogona* subsp. *calycogona* (16), *E. phenax* subsp. *phenax* (14). Shrubs *Melaleuca ulicoides* (20), *Acacia binata* (18), *Pultenaea purpurea* (15). Herbs *Lepidosperma fimbriatum* (15).

Most common taxa (% frequency): Trees *Eucalyptus flocktoniae* subsp. *flocktoniae* (91), *E. phenax* subsp. *phenax* (88), *E. calycogona* subsp. *calycogona* (53), *E. cernua* (32). Shrubs *Melaleuca ulicoides* (68), *Pultenaea purpurea* (53), *Acacia glaucoptera* (50), *Hakea commutata* (44), *Acacia binata* (41), *Exocarpos aphyllus* (41), *Dodonaea concinna* (38), *Acacia ingrata* (35), *Daviesia nematophylla* (35), *Melaleuca cincinnata* (35), *Boronia inornata* subsp. *leptophylla* (29), *Hibbertia psilocarpa* (26), *Melaleuca torquata* (26). Herbs *Cassytha melantha* (65), *Lepidosperma fimbriatum* (50), *Lepidosperma gahnioides* (26).

Short range endemics

10–100 km²: *Eucalyptus purpurata*.

100–1000 km²: *Acacia* sp. Ravensthorpe Range (B.R. Maslin 5463), *Eucalyptus gardneri* subsp. *ravensthorpensis*, *Grevillea punctata*, *G. sulcata*, *Melaleuca ulicoides*, *Spyridium glaucum*.

1000–10,000 km²: *Acacia ophiolithica*, *A. pusilla*, *Dodonaea trifida*, *Eucalyptus cernua*, *E. oleosa* subsp. *corvina*, *Melaleuca pomphostoma*.

Conservation taxa: *Acacia* sp. Ravensthorpe Range [BR Maslin 5463] (P1), *Eucalyptus purpurata* (R), *Grevillea fastigiata* (P4), *G. punctata* (P3), *G. sulcata* (P1).

Community 4: *Eucalyptus tenera*/*E. flocktoniae*/*E. pileata* tall mallee shrublands over *Melaleuca* spp.

Map unit(s): Epil/Edep/Mspp (2), Epil/Edep (1), Eunc/Essp (1).

Note: Since vegetation mapping was undertaken many Ravensthorpe Range collections of *Eucalyptus depauperata* have been redetermined as *Eucalyptus tenera*.

Community 4 was described from five plots, and consisted of tall mallee shrublands comprised of a number of eucalypt species, most commonly *E. tenera*, *E. flocktoniae*, *E. pileata* and *E. phaenophylla*. Unlike community 3, *E. phenax* was notably absent from this community. A variety of species of *Melaleuca* was found in the tall shrub stratum, many of these common to, or characteristic of, the community type. All sites were located in the northern section of the range, on the gently inclined, lower slopes around Mt Short and Overshot Hill.

Soils: red-brown to brown coloured, light to heavy clays and clay loams, relatively deep with no exposed bedrock.

No. plots 5; Taxa/plot (SD) 25.2 ± 2.4 ; Altitude 285–308 m

Diagnostic taxa (INDVAL): Trees *Eucalyptus tenera* (67), *E. pileata*



Figure 5. Community 3, typical mallee woodland over dense *Melaleuca* shrubland (survey plot R 140).



Figure 6. Community 4, tall mallee shrubland (survey plot R 107) on north end of the range.

(29), *E. flocktoniae* subsp. *flocktoniae* (16). **Shrubs** *Melaleuca societatis* (60), *M. undulata* (57), *M. teuthidoides* (45), *M. johnsonii* (40), *M. coronicarpa* (30), *Daviesia benthamii* (24), *Melaleuca torquata* (24), *Microcybe albiflora* (24), *Pultenaea purpurea* (19), *Grevillea pectinata* (18), *Melaleuca acuminata* subsp. *acuminata* (18), *Thomasia microphylla* (17). **Herbs** *Lepidosperma* sp. [GF Craig 8243] (23), *L. sp.* Maydon [S Kern et al. LCH 17844] (22), *Cassytha melantha* (17).

Most common taxa (% frequency): **Trees** *Eucalyptus flocktoniae* subsp. *flocktoniae* (100), *E. tenera* (80), *E. pileata* (80), *E. phaenophylla* (60), *E. calycogona* subsp. *calycogona* (40). **Shrubs** *Exocarpos aphyllus* (60), *Grevillea pectinata* (60), *Hibbertia psilocarpa* (60), *Melaleuca societatis* (60), *M. teuthidoides* (60), *M. torquata* (60), *M. undulata* (60), *Pultenaea purpurea* (60), *Spyridium cordatum* (60), *Boronia inconspicua* (40), *B. inornata* subsp. *leptophylla* (40), *Coopernochla strophiolata* (40), *Daviesia benthamii* (40), *Gastrolobium parviflorum* (40), *Hakea commutata* (40), *Hibbertia pungens* (40), *Melaleuca acuminata* subsp. *acuminata* (40), *M. coronicarpa* (40), *M. cucullata* (40), *M. glaberrima* (40), *M. hamata* (40), *M. johnsonii* (40), *M. lateriflora* subsp. *lateriflora* (40), *M. pauperiflora* subsp. *pauperiflora* (40), *Microcybe albiflora* (40), *Thomasia microphylla* (40), *Wilsonia humilis* (40). **Herbs** *Cassytha melantha* (80), *Gahnia ancistrophylla* (40), *Lepidosperma* sp. Bandalup Scabrid [N Eveleigh 10798] (40), *L. sp.* [GF Craig 8243] (40), *L. sp.* Maydon [S Kern et al. LCH 17844] (40).

Short range endemics

1000–10,000 km²: *Acacia pusilla*.

Conservation taxa: none recorded.

Community 5: *Eucalyptus clivicola/E. extensa/E. platypus* woodlands and open forest.

Map unit(s): Ecli (2), Ecer (1), Eext (1), Epla (1), Epla/Mcuc (1), Esab (1).

Community 5 consisted of woodlands and open forests most commonly dominated by mixtures of *Eucalyptus clivicola*, *E. extensa*, *E. platypus*, *E. flocktoniae* and/or *E. pileata*. Many of the dominant tree species were mallets. The understorey shrub stratum was typically sparse and the diversity of taxa in the understorey was low. Community 5 was present across most of the range except for the area around Bandalup Hill on gently to moderately inclined, lower to middle hill slopes.

Soils: clay and clay loams to sandy clays with no exposed bedrock.

No. plots 12; Taxa/plot (SD) 11.8 ± 1.5; Altitude 102–369 m

Diagnostic taxa (INDVAL): **Trees** *Eucalyptus platypus* (33), *E. extensa* (31), *E. clivicola* (21).

Most common taxa (% frequency): **Trees** *Eucalyptus clivicola* (67), *E. extensa* (58), *E. flocktoniae* subsp. *flocktoniae* (58), *E. pileata* (50), *E. platypus* (50). **Shrubs** *Exocarpos aphyllus* (50), *Acacia glaucoptera* (42), *Melaleuca torquata* (42), *Lasiopetalum compactum* (33), *Melaleuca cucullata* (33), *Boronia inornata* subsp. *leptophylla* (25), *Grevillea patentiloba* subsp. *patentiloba* (25), *G. pectinata* (25), *Hibbertia psilocarpa* (25), *Melaleuca hamata* (25), *Microcybe albiflora* (25), *Olearia muelleri* (25). **Herbs** *Lepidosperma* sp. Bandalup Scabrid [N Eveleigh 10798] (25), *L. sp.* Ravensthorpe [GF Craig 5188] (25).



Figure 7. Community 5, often a mallet woodland over a sparse understorey (survey plot R198).

Short range endemics

100–1000 km²: *Melaleuca ulicoides*, *Pultenaea craigiana*.

1000–10,000 km²: *Eucalyptus cernua*, *E. oleosa* subsp. *corvina*.

Conservation taxa: *Pultenaea craigiana* (P3).

Community 6: *Eucalyptus gardneri* subsp. *ravensthorpensis* woodlands and tall open forest.

Map unit(s): Egar (2), Emeg (1).

This community consisted of woodlands and tall open forests of *Eucalyptus gardneri* subsp. *ravensthorpensis*, often with the co-occurrence of both *E. cernua* and *E. phenax*, and *E. megacornuta* on occasion. The associated understorey was usually sparse; it occurred on moderately to steeply inclined upper hill slopes. Plots of this community type were located along the main ridges in the central region of the main range and also on Overshot Hill.

Soils: red-brown clay loam soils with minimal or no exposure of bedrock.

No. plots 5; Taxa/plot (SD) 9.8 ± 1.3; Altitude 178–321 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus gardneri* subsp. *ravensthorpensis* (53), *E. cernua* (25). Shrubs *Boronia oxyantha* var. *brevicalyx* (26), *Spyridium glaucum* (24), *Gastrolobium parviflorum* (14).

Most common taxa (% frequency): Trees *Eucalyptus gardneri* subsp. *ravensthorpensis* (100), *E. cernua* (60), *E. phenax* subsp. *phenax* (60). Shrubs *Boronia oxyantha* var. *brevicalyx* (80), *Gastrolobium parviflorum* (80), *Spyridium glaucum* (60), *Acacia erinacea* (40), *Beyeria sulcata* var. *brevipes* (40), *Exocarpos aphyllus* (40), *Persoonia tereticolia* (40).

Short range endemics

100–1000 km²: *Eucalyptus gardneri* subsp. *ravensthorpensis*, *Spyridium glaucum*.

1000–10,000 km²: *Boronia oxyantha* var. *brevicalyx*, *Eucalyptus cernua*, *E. megacornuta*.

Conservation taxa: none recorded.

Community 7: *Eucalyptus astringens*/*E. clivicola* woodlands.

Map unit(s): East (1), Epla (1).

Community 7 was variable, typically dominated by woodlands of *Eucalyptus astringens* and *E. clivicola*. However, one site was an *E. leptocalyx*/*E. flocktoniae*/*E. phaeophylla* open mallee shrubland over dense *Melaleuca stramentosa* shrubland. These woodlands occurred both in the north and south of the range, were typically species poor and usually recorded on gently to moderately inclined midslopes.

Soils: grey to grey-brown sandy clay loams with no exposed bedrock.



Figure 8. Community 6, forming open woodlands along the range crest (survey plot R095).



Figure 9. Community 7, typically open woodlands (survey plot R171), but mallee shrublands were also encountered.

No. plots 4; Taxa/plot (SD) 12.8 ± 1.9 ; Altitude 129–335 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus astringens* (75), *E. clivicola* (26). Shrubs *Melaleuca stramentosa* (48), *Dodonaea trifida* (47), *Boronia oxyantha* var. *brevicalyx* (23). Herbs *Lepidosperma* sp. [GF Craig 8249] (22).

Most common taxa (% frequency): Trees *Eucalyptus astringens* (75), *E. clivicola* (75), *E. flocktoniae* subsp. *flocktoniae* (50), *E. falcata* subsp. *falcata* (25), *E. incrassata* (25), *E. leptocalyx* (25), *E. phaenophylla* (25), *E. suggrandis* subsp. *suggrandis* (25). Shrubs *Boronia oxyantha* var. *brevicalyx* (75), *Melaleuca stramentosa* (75), *Baeckea corynophylla* (50), *Dodonaea trifida* (50), *Exocarpos aphyllus* (50), *Melaleuca hamata* (50), *Acacia gonophylla* (25), *A. patagiata* (25), *Acrotriche parviflora* (25), *Boronia inconspicua* (25), *Chorizema trigonum* (25), *Daviesia articulata* (25), *D. lancifolia* (25), *Dodonaea amblyophylla* (25), *Gastrolobium parviflorum* (25), *Gompholobium confertum* (25), *Grevillea patentiloba* subsp. *patentiloba* (25), *Hakea laurina* (25), *Hibbertia psilocarpa* (25), *Lasiopetalum compactum* (25), *Melaleuca bracteosa* (25), *Nematolepis phebaloides* (25), *Rinzia communis* (25), *Siegfriedia darwinicoides* (25), *Spyridium glaucum* (25), *Tempeltonia retusa* (25), *Thomasia microphylla* (25). Herbs *Lepidosperma* sp. [GF Craig 8249] (25), *L. sp.* Mt Chester [S Kern et al. LCH 16596] (25), *L. sp.* Ravensthorpe [GF Craig 5188] (25).

Short range endemics

100–1000 km²: *Melaleuca stramentosa*, *Spyridium glaucum*.

1000–10,000 km²: *Boronia oxyantha* var. *brevicalyx*, *Dodonaea trifida*.

Conservation taxa: *Acrotriche parviflora* (P4).

Mallee shrubland group

Community 8: *Eucalyptus* sp. Ravensthorpe (AS George 616) tall mallee shrubland or *Allocasuarina*/ *Melaleuca* shrublands.

Map unit(s): Edes/Alca (1), Epro/Mspp (1), EspR/Mcli (1).

This community was structurally heterogeneous, typically being *Eucalyptus* sp. Ravensthorpe (AS George 616) tall mallee shrubland. Less frequent canopy eucalypt species included *E. flocktoniae*, *E. phenax*, *E. proxima* and *E. brachycalyx*, while *Allocasuarina* (*A. campestris* or *A. hystericosa*) and *Melaleuca* formed the dominant canopy stratum at some sites. All plots classified as this community were located in the south-eastern region of the Ravensthorpe Range, on gently undulating, hilly terrain east of the main ridges. Sites were generally gently to moderately inclined hillslopes, occurring from lower slopes to hillcrests.

Soils: typically red-brown clays with occasional metavolcanic outcropping.

No. plots 12; Taxa/plot (SD) 23.0 ± 2.0 ; Altitude 84–190 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus* sp. Ravensthorpe [AS George 616] (20). Shrubs *Calytrix tetragona* (48), *Grevillea fastigiata* (39), *Cryptandra myriantha* (33), *Styphelia pulchella* (25), *Dodonaea pinifolia* (22), *Hibbertia rostellata* (19), *Calothamnus quadrifidus* (18), *Hybanthus floribundus* subsp. *adpressus* (17). Herbs *Stackhousia monogyna* (34), *Lepidosperma diurnum* (32), *Neurachne alopecuroidea* (16), *Wurmbea cernua* (15).



Figure 10. Community 8, forming dense tall shrublands (survey plot R047) on the south-eastern section of the range.

Most common taxa (% frequency): Trees *Eucalyptus* sp. Ravensthorpe [AS George 616] (58). Shrubs *Dodonaca pinifolia* (92), *Hakea verrucosa* (75), *Hibbertia pungens* (75), *Calothamnus quadrifidus* (67), *Calytrix tetragona* (67), *Melaleuca hamata* (67), *Hybanthus floribundus* subsp. *adpressus* (58), *Acacia ophiolithica* (50), *Daviesia anceps* (50), *Astrolobma epacridis* (42), *Grevillea fastigiata* (42), *Melaleuca cliffortioides* (42), *Styphelia pulchella* (42), *Astrolobma serratifolium* (33), *Cryptandra myriantha* (33), *Hibbertia rostellata* (33), *Acacia glaucoptera* (25), *Allocasuarina campestris* (25), *A. hystricosa* (25), *Grevillea oligantha* (25), *Halgnania andromedifolia* (25), *Petrophile fastigiata* (25), *Verticordia acerosa* var. *preissii* (25). Herbs *Neurachne alopecuroides* (83), *Lepidosperma diurnum* (75), *Stackhousia monogyna* (50), *Cassytha glabella* (42), *Dianella revoluta* var. *revoluta* (42), *Dampiera lavandulacea* (33), *Wurmbea cernua* (33), *Gahnia ancistrophylla* (25), *Spartochloa scirpoidea* (25).

Short range endemics

100–1000 km²: *Kunzea cincinnata*, *Pultenaea calycina* subsp. *proxena*.

1000–10,000 km²: *Acacia ophiolithica*, *Allocasuarina hystricosa*.

Conservation taxa: *Allocasuarina hystricosa* (P4), *Grevillea fastigiata* (P4), *Pultenaea calycina* subsp. *proxena* (P4).

Community 9: *Eucalyptus phenax*/*E. cernua* over *Melaleuca cliffortioides* and *Acacia ophiolithica*.

Map unit(s): EspR/Mcli (1).

Community 9 generally consisted of tall mallee shrubland dominated by *Eucalyptus phenax* and/or *E. cernua*, also with *E. leptocalyx* or *E. calycogona* as less common canopy species. The shrub stratum ranged from mid-dense to dense in structure. Significant indicator species included the vulnerable, DRF-listed *Beyeria cockertonii*. This community generally occurred on gently to moderately inclined hill slopes, on the lower to mid slopes of low, undulating hills on the eastern side of the main range and on Bandalup Hill.

Soils: rocky grey-brown medium to heavy clays, with no exposed bedrock.

No. plots 5; Taxa/plot (SD) 22.8 ± 4.6; Altitude 123–231 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus leptocalyx* (19). Shrubs *Beyeria cockertonii* (60), *Grevillea oligantha* (28), *Melaleuca pomphostoma* (24), *Hakea verrucosa* (23), *Pultenaea rotundifolia* (23), *Melaleuca cliffortioides* (21), *Acacia ophiolithica* (17), *Dodonaca pinifolia* (16), *Melaleuca ulicoides* (16), *Acacia glaucoptera* (15). Herbs *Wurmbea cernua* (22).

Most common taxa (% frequency): Trees *Eucalyptus phenax* subsp. *phenax* (80), *E. leptocalyx* (60), *E. cernua* (40), *E. uncinata* (40). Shrubs *Hakea verrucosa* (100), *Acacia glaucoptera* (80), *Dodonaca pinifolia* (80), *Grevillea oligantha* (80), *Acacia ophiolithica* (60), *Beyeria cockertonii* (60), *Daviesia anceps* (60), *Hibbertia psilocarpa* (60), *Melaleuca cliffortioides* (60), *Melaleuca ulicoides* (60), *Acrotriche cordata* (40), *Cryptandra wilsonii* (40), *Hakea commutata* (40), *Halgnania andromedifolia* (40), *Hybanthus floribundus* subsp. *adpressus* (40), *Logania buxifolia* (40), *Melaleuca lateriflora* subsp. *lateriflora* (40), *M. pomphostoma* (40), *Nematolepis phebaliooides*



Figure 11. Community 9 was found scattered along the range and most often formed tall open mallees over a dense understorey (survey plot R092).



Figure 12. Community 10, dense *Melaleuca pomphostoma* shrubland on skeletal soils, apparently restricted to Overshot Hill (survey plot R205).

(40), *Pomaderis brevifolia* (40), *Pultenaea rotundifolia* (40), *Santalum acuminatum* (40). **Herbs** *Gahnia aristata* (40), *Lepidosperma diurnum* (40), *Wurmbea cernua* (40).

Short range endemics

10–100 km²: *Beyeria cockertonii*, *Pultenaea wudjariensis*.

100–1000 km²: *Leucopogon infuscatus*, *McIalcea ulicoides*, *Pultenaea calycina* subsp. *proxena*.

1000–10,000 km²: *Acacia ophiolithica*, *Eucalyptus cernua*, *McIalcea pomphostoma*.

Conservation taxa: *Beyeria cockertonii* (R), *Pultenaea calycina* subsp. *proxena* (P4), *P. wudjariensis* (P1)

Community 10: *McIalcea pomphostoma* heaths on skeletal soils on outcrops.

Map unit(s): not mapped.

This community was found at only two sites, both of which occurred on Overshot Hill Nature Reserve. Structurally, it formed a dense shrubland dominated by a variety of taxa, notably *McIalcea pomphostoma* and *Hakea verrucosa*. Over this dense shrubland were emergent tall shrubs and mallets of *Eucalyptus cernua* that appear to be outliers from the adjacent mallet woodlands. *Allocasuarina huilliana* was an additional emergent tree in the site near the crest of Mt Short. This community was found on both crests and lower down the slope in gently inclined, shallow valleys eroded into the hillside.

Soils: skeletal to shallow red-brown sandy clay loams with significant exposure of the metavolcanic bedrock.

No. plots 2; Taxa/plot (SD) 21.0 ± 2.0; Altitude 326–372 m.

Diagnostic taxa: insufficient plots.

Most common taxa: insufficient plots.

Short range endemics

1000–10,000 km²: *Eucalyptus cernua*, *McIalcea pomphostoma*.

Conservation taxa: none recorded.

Community 11: *Eucalyptus brachycalyx*/*E. indurata* tall, open mallee shrublands.

Map unit(s): Ebra (1), Eind/Mpau (1), Eole/Mpau (1).

Community 11 typically consisted of tall, open mallee shrublands typically dominated by *Eucalyptus brachycalyx* and *E. indurata*, but also having *E. purpurea* or *E. sp.* Ravensthorpe (AS George 616) as dominant species at some sites. The shrub understorey tended to be structurally dominated by a few main taxa, a number of which (*Leptomeria pachyclada*, *Pultenaea calycina* subsp. *proxena*), are restricted to the south-eastern part of the range, and some species (*E. purpurea* and *Acrotriche orbicularis*) are restricted to Bandalup Hill. Community 11 occurred on the gently inclined mid to upper slopes of low, undulating hills on the eastern side of the main range and on Bandalup Hill.



Figure 13. Community 11 was structurally variable, here forming a low open woodland (survey plot R086) on the south-eastern side of the main range and Bandalup Hill.

Soils: generally rocky brown silty loams – silty clay loams with some magnesite from weathered ultramafics. There was limited magnesite outcrop.

No. plots 15; Taxa/plot (SD) 16.4 ± 2.1 ; Altitude 106–244 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus indurata* (56), *E. brachycalyx* (38). Shrubs *Beyeria villosa* (53), *Pultenaea calycina* subsp. *proxima* (42), *Acacia pauperiflora* subsp. *pauperiflora* (35), *Leptomeria pachyclada* (20), *Acacia brevifolia* (20), *Acacia ingrica* (19), *A. mutabilis* subsp. *mutabilis* (19), *Boronia inornata* subsp. *inornata* (17), *Halgania andromedifolia* (14).

Most common taxa (% frequency): Trees *Eucalyptus brachycalyx* (87), *E. indurata* (80). Shrubs *Acacia ingrica* (80), *Acacia pauperiflora* subsp. *pauperiflora* (80), *Boronia inornata* subsp. *inornata* (67), *Pultenaea calycina* subsp. *proxima* (67), *Halgania andromedifolia* (60), *Beyeria villosa* (53), *Hakea verrucosa* (47), *Pomaderris brevifolia* (47), *Dodonaea concinna* (40), *Leptomeria pachyclada* (40), *Westringia dampieri* (40), *Daviesia anceps* (33), *Nematolepis phebaloides* (33), *Acacia mutabilis* subsp. *mutabilis* (27), *Exocarpos sparteus* (27), *Hibbertia rostellata* (27), *Acacia cliffortioides* (27). Herbs *Cassytha glabella* (33), *Tetraria* sp. Mt Madden [CD Turley 40 BP/897] (27).

Short range endemics

0–10 km²: *Acrotriche orbicularis*.

10–100 km²: *Eucalyptus purpurata*.

100–1000 km²: *Beyeria villosa*, *Acacia ulicoides*, *Pultenaea calycina* subsp. *proxima*.

1000–10,000 km²: *Acacia ophiolithica*, *Eucalyptus oleosa* subsp. *corvina*.

Conservation taxa: *Acrotriche orbicularis*, *Beyeria villosa* (P4), *Eucalyptus purpurata* (R), *Pultenaea calycina* subsp. *proxima* (P4).

Community 12: *Eucalyptus oleosa* subsp. *corvina* tall, open mallee shrubland and open forest.

Map unit(s): Eole (2), EspR/Mcli (1).

A small, structurally heterogeneous group closely allied to community 11, community 12 consisted of tall, open mallee shrubland and open forest dominated by *Eucalyptus oleosa* subsp. *corvina*, co-occurring with other eucalypt species (*E. brachycalyx*, *E. calycogona*, *E. myriadena* subsp. *myriadena*). The shrub understorey was structurally sparse. Sites of this community were located on the on slopes east of the main range in the central to south-east part of the range on lower to upper slopes. The sites varied from gentle to steep.

Soils: typically brown to grey clay loams and clays, sometimes with komatiite and magnesite outcropping. Soils and surface shale, komatiite and magnesite were derived from underlying mafic and ultramafic substrates.

No. plots 4; Taxa/plot (SD) 13.8 ± 0.8 ; Altitude 102–214 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus oleosa* subsp. *corvina* (65), *E. celastroides* subsp. *virella* (22). Shrubs *Boronia inornata*



Figure 14. Community 12 was also structurally variable here forming a tall mallee shrubland (survey plot R028) was also found on the eastern side of main range.

subsp. *inornata* (38), *Rhagodia crassifolia* (21), *Senna artemisioides* subsp. x *artemisioides* (21).

Most common taxa (% frequency): Trees *Eucalyptus oleosa* subsp. *corvina* (100), *E. brachycalyx* (50), *E. celsastroides* subsp. *virella* (25), *E. clivicola* (25), *E. extensa* (25), *E. indurata* (25), *E. myriadena* subsp. *myriadena* (25), *E. phenax* subsp. *phenax* (25), *E. pileata* (25). Shrubs *Boronia inornata* subsp. *inornata* (100), *Senna artemisioides* subsp. x *artemisioides* (75), *Acacia glaucoptera* (50), *A. ingrata* (50), *Exocarpos aphyllus* (50), *Hovea acanthoclada* (50), *Lasiopetalum compactum* (50), *Rhagodia crassifolia* (50), *Santalum acuminatum* (50), *Acacia cyclops* (25), *A. erinacea* (25), *Atriplex semibaccata* (25), *Dodonaea concinna* (25), *Enchytraea tomentosa* var. *tomentosa* (25), *Exocarpos sparteus* (25), *Halgnia andromedifolia* (25), *Hibbertia psilocarpa* (25), *Melaleuca acuminata* subsp. *acuminata* (25), *M. pauperiflora* subsp. *pauperiflora* (25), *Nematolepis phlebalooides* (25), *Olearia muelleri* (25), *Phyllanthus scaber* (25), *Pultenaea purpurea* (25), *Westringia dampieri* (25). Herbs *Austrostipa elegantissima* (50), *A. exilis* (25), *Lepidosperma fimbriatum* (25), L. sp. Saltbush Hill [KR Newbey 4118] (25), *Platysace maxwellii* (25).

Short range endemics

1000–10,000 km²: *Eucalyptus oleosa* subsp. *corvina*.

Conservation taxa: none recorded.

Species-rich mallee shrubland group

Community 13: Upper slope *Eucalyptus gardneri* subsp. *ravensthorpensis*/*E. clivicola*/*E. falcata* woodlands/mallee shrublands.

Map unit(s): Ecli (5), Egard (5), Efal (3), Efal/Eple (2), Mham (2), Efal/Alca (1), Emeg (1).

Community 13 usually consisted of woodlands and mallee shrublands dominated by *Eucalyptus gardneri* subsp. *ravensthorpensis* and *E. falcata* subsp. *falcata*, but *E. clivicola* was also a frequent canopy species. The understorey was typically an open shrubland composed commonly of the taller shrubland species. *Melaleuca thapsina* is a distinctive tall shrub that was found at some sites as dense stands within a distinctive belt on the upper slopes of the range. At some sites, it formed thickets under very sparse stands of *E. pleurocarpa* and/or *E. falcata* mallee shrubs. As with other woodland communities, species richness was relatively low. This community was mostly found in the central part of the range typically on mid to upper hillslopes, where the slope varies from gentle to steep.

Soils: rocky red-brown to dark brown clay loams and clay loam sands. Exposed bedrock included massive gossan.

No. plots 23; Taxa/plot (SD) 16.5 ± 1.1; Altitude 119–339 m.



Figure 15. Community 13 was one of the most structurally variable communities, here forming an open woodland (survey plot R261).

Diagnostic taxa (INDVAL): Shrubs *Phebalium tuberculosum* (48), *Trymalium elachophyllum* (16).

Most common taxa (% frequency): Trees *Eucalyptus falcata* subsp. *falcata* (74), *E. clivicola* (39), *E. gardneri* subsp. *ravensthorpensis* (39). **Shrubs** *Gastrolobium parviflorum* (74), *Melaleuca hamata* (65), *Phebalium tuberculosum* (65), *Dodonaea pinifolia* (43), *Hibbertia pungens* (43), *Lasiopetalum compactum* (43), *Santalum acuminatum* (43), *Beyeria sulcata* var. *brevipes* (39), *Labichea lanccolata* subsp. *brevifolia* (30), *Acacia* sp. Ravensthorpe [RS Cowan & BR Maslin RSC A-760] (26), *Hakea laurina* (26), *Philotheca gardneri* subsp. *gardneri* (26), *Trymalium elachophyllum* (26). **Herbs** *Platysace maxwellii* (61), *Austrodanthonia setacea* (39), *Austrostipa hemipogon* (30), *Stylium albomontis* (30), *Lepidosperma* sp. Cordingup [GF Craig 6138] (26), *Thysanotus patersonii* (26).

Short range endemics

100–1000 km²: *Eucalyptus gardneri* subsp. *ravensthorpensis*, *Isopogon* sp. Ravensthorpe (DB Foreman 1207), *Kunzea cincinnata*, *Spiridium glaucum*.

1000–10,000 km²: *Acacia durabilis*, *A. pusilla*, *Boronia oxyantha* var. *brevicalyx*, *B. ternata* var. *elongata*, *Eucalyptus megacornuta*, *Grevillea patentiloba* subsp. *platypoda*.

Community 14: Upland *Eucalyptus falcata*, *E. pleurocarpa*, *Beaufortia orbifolia*, *Banksia* spp. mallee shrublands.

Map unit(s): Efal/Eple (13), Blac/Borb (3), Ecli (1), Efal (1), Efal/Alca (1), Eunc/Essp (1).

Community 14 consisted of upland *Eucalyptus falcata*/*E. pleurocarpa* mallee shrublands that occur on the lateritic duricrust that covers the ridges of the main range. This community was structurally heterogeneous. These shrub mallees overtopped a tall, mid-dense to dense shrub stratum. On occasion, this upper shrub stratum was dominated by stands of *Allocasuarina acutivalvis* subsp. *acutivalvis* or *A. campestris*. Most collections of the relatively uncommon and endemic species, *Hibbertia atrichosepala* were from this community type. Community 14 was located across the north-south extent of the range, mostly on the gently to moderately inclined mid to upper hillslopes and extending up onto crests. These are areas of the range where laterite has formed from in situ weathering.

Soils: stony brown sandy loams, with infrequent outcropping of bedrock—usually laterite and occasionally some metavolcanics/gossan.

No. plots 26; Taxa/plot (SD) 24.9 ± 1.7 ; Altitude 119–447 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus falcata* subsp. *falcata* (17), *E. pleurocarpa* (14). **Shrubs** *Philotheca gardneri* subsp. *gardneri* (34), *Brachyloma geissoloma* subsp. *geissoloma* (29), *Beaufortia orbifolia* (27), *Acacia* sp. Ravensthorpe [RS Cowan & BR Maslin RSC A-760] (18), *Banksia laevigata* subsp. *laevigata* (18), *Boronia ternata* var. *elongata* (18), *Hakea multilineata* (17), *Isopogon* sp. Ravensthorpe [DB Foreman 1207] (15), *Hakea obtusa* (14).

Most common taxa (% frequency): Trees *Eucalyptus falcata* subsp. *falcata* (92), *E. pleurocarpa* (85), *E. uncinata* (31). **Shrubs** *Beaufortia orbifolia* (69), *Hakea obtusa* (65), *Melaleuca hamata* (65), *Gastrolobium parviflorum* (62), *Philotheca gardneri* subsp. *gardneri* (62), *Acacia* sp. Ravensthorpe [RS Cowan & BR Maslin RSC A-



Figure 16. Community 14, open mallee shrubland on the lateritic ridges along main range (survey plot R250).

760] (54), *Santalum acuminatum* (50), *Hibbertia gracilipes* (46), *Isopogon* sp. Ravensthorpe [DB Foreman 1207] (46), *Brachyloma geissoloma* subsp. *geissoloma* (38), *Lasicopetalum compactum* (38), *Acrotriche ramiflora* (35), *Beyeria sulcata* var. *brevipes* (35), *Goodenia pinifolia* (35), *Leucopogon cuneifolius* (35), *Banksia laevigata* subsp. *laevigata* (31), *Calothamnus quadrifidus* (31), *Hakea verrucosa* (31), *Rhadinothamnus rufus* subsp. *amblycarpus* (31), *Boronia ternata* var. *elongata* (27), *Persoonia teretifolia* (27). **Herbs** *Stylium albomontis* (65), *Lepidosperma* sp. [RL Barrett 3553] (46), *L.* sp. Cordingup [GF Craig 6138] (42), *Platysace maxwellii* (31), *Dampiera angulata* subsp. *angulata* (27).

Short range endemics

0–10 km²: *Hibbertia atrichosepala*.

100–1000 km²: *Guichenotia anota*, *G. apetala*, *Isopogon* sp. Ravensthorpe (DB Foreman 1207), *Kunzea cincinnata*, *Marianthus mollis*, *Micromyrtus navicularis*, *Spyridium glaucum*

1000–10,000 km²: *Acacia heterochroa* subsp. *heterochroa*, *A. ophiolitica*, *A. pinguiculosa* subsp. *pinguiculosa*, *Banksia laevigata* subsp. *laevigata*, *Boronia ternata* var. *elongata*, *Grevillea patentiloba* subsp. *platypoda*, *Leptospermum* sp. Bandalup Hill (G Cockerton 11001), *Melaleuca pomphostoma*.

Conservation taxa: *Banksia laevigata* subsp. *laevigata* (P4), *Guichenotia anota* (P1), *G. apetala* (P1), *Hibbertia atrichosepala* (P1), *Marianthus mollis* (R), *Micromyrtus navicularis* (P3), *Pterostylis* sp. Ongerup [KR Newbey 4874] (P4).

Community 15: *Eucalyptus* spp/*Banksia lehmanniana*/*B. heliantha* mallee shrublands.

Map unit(s): Dque (1), Efal/Eple (1), Mstr (1).

Community 15 has close floristic affinities to community 14. This community consisted of mixed *Eucalyptus* mallee shrublands, where

a variety of combinations of *Eucalyptus* species were co-dominant, including *E. pleurocarpa*, *E. falacata* subsp. *falcata*, *E. uncinata*, *E. preissiana* subsp. *preissiana* and *E. lehmannii* subsp. *parallela* over a tall shrubland. *Lepidosperma* sp. (AS George 9935) was a common understorey sedge on the slopes of Bandalup Hill. Community 15 was located on the crest of Bandalup Hill, and around Mt Desmond and Mt Chester on the southern parts of the main range ridge. This community was typically associated with moderately-inclined hill slopes, generally on upper slopes but ranging from lower slopes to crests.

Soils: variable in texture from brown to grey sandy loams, to clays and clay sands.

No. plots 6; Taxa/plot (SD) 32.5 ± 1.2 ; Altitude 165–315 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus preissiana* subsp. *preissiana* (34), *E. lehmannii* subsp. *parallela* (32). **Shrubs** *Banksia heliantha* (36), *Grevillea concinna* subsp. *lehmanniana* (33), *Hakea obtusa* (33), *Andersonia* aff. *lehmanniana* [S Kern & R Jasper LCH 17769] (31), *Beaufortia orbifolia* (25), *Boronia subsessilis* (24), *Daviscia mollis* (23), *Grevillea nudiflora* (19), *Leucopogon carinatus* (19), *L. infuscatus* (19), *Boronia oxyantha* var. *brevicalyx* (18), *Spiridium majoranifolium* (18), *Chorizema trigonum* (17), *Taxandria spathulata* (17), *Hovea acanthoclada* (16). **Herbs** *Stylium albomontis* (19), *Lepidosperma* sp. Cordingup [GF Craig 6138] (18).

Most common taxa (% frequency): Trees *Eucalyptus pleurocarpa* (67), *E. uncinata* (67), *E. preissiana* subsp. *preissiana* (50), *E. lehmannii* subsp. *parallela* (50), *E. falcata* subsp. *falcata* (33), *E. incrassata* (33), *E. pluricaulis* subsp. *pluricaulis* (33). **Shrubs** *Hakea obtusa* (100), *Banksia heliantha* (83), *Beaufortia orbifolia* (67), *Boronia oxyantha* var. *brevicalyx* (67), *Gastrolobium parviflorum* (67), *Hibbertia pungens* (67), *Hovea acanthoclada* (67), *Taxandria spathulata* (67), *Acrotriche cordata* (50), *A. ramiflora* (50), *Banksia*




Figure 17. Community 15, species-rich shrublands on the lateritic ridges in the south of the study area (survey plot R002).

Icmanniana (50), *Beaufortia schaueri* (50), *Boronia inconspicua* (50), *Chorizema trigonum* (50), *Leucopogon carinatus* (50), *L. infuscatus* (50), *Melaleuca hamata* (50), *Acacia heterochroa* subsp. *heterochroa* (33), *Andersonia* aff. *Icmanniana* [S Kern & R Jasper LCH 17769] (33), *Banksia cirsoides* (33), *Boronia crassifolia* (33), *B. subsessilis* (33), *Daviesia mollis* (33), *Grevillea concinna* subsp. *Icmanniana* (33), *G. nudiflora* (33), *Isopogon* sp. Ravensthorpe [DB Foreman 1207] (33), *Lasiopetalum compactum* (33), *Leptospermum* sp. Bandalup Hill [G Cockerton 11001] (33), *Melaleuca pentagona* (33), *M. rigidifolia* (33), *M. stramentosa* (33), *Petrophile glauca* (33), *Rhadinothamnus rufus* subsp. *amblycarpus* (33), *Siegfriedia darwinioidea* (33), *Spyridium majoranifolium* (33). **Herbs** *Stylium albomontis* (83), *Lepidosperma* sp. Cordingup [GF Craig 6138] (67), *L. sp.* Saltbush Hill [KR Newbey 4118] (50), *Lomandra mucronata* (50), *Tetraria* sp. Mt Madden [CD Turley 40 BP/897] (50), *Casytha glabella* (33), *C. melantha* (33), *Lepidosperma* sp. [AS George 9935] (33), *L. sp.* Mt Short [S Kern et al. LCH 17510] (33), *Lomandra micrantha* subsp. *teretifolia* (33).

Short range endemics

0–10 km²: *Hibbertia abyssa*.

1000–10,000 km²: *Guichenotia apetala*, *Isopogon* sp. Ravensthorpe (DB Foreman 1207), *Leucopogon infuscatus*, *Marianthus mollis*, *Melaleuca stramentosa*, *Spyridium glaucum*.

10000–10,000 km²: *Acacia disticha*, *A. durabilis*, *A. heterochroa* subsp. *heterochroa*, *A. laricina* var. *crassifolia*, *Boronia oxyantha* var. *brevicalyx*, *Grevillea patentiloba* subsp. *platypoda*, *Leptospermum* sp. Bandalup Hill (G Cockerton 11001).

Conservation taxa: *Guichenotia apetala* (P1), *Hibbertia abyssa* (R), *Marianthus mollis* (R).

Community 16: *Eucalyptus falcata* subsp. *falcata*/E. spp. mallee shrublands.

Map unit(s): Ecli (2), East (1), Efal/Eple (1).

Common canopy species of community 16 included *Eucalyptus falcata* subsp. *falcata* and *E. pleurocarpa*, although *E. megacornuta* occurred as the dominant stratum in open forest at one site with exposed gossameriferous bedrock. This community had only half the species richness of related communities (17–21). Community 16 occurred on the steep mid to upper slopes and gently inclined hillcrests of Mt Desmond and Mt Chester, on the southern main ridges of the range.

Soils: typically dark loamy sands or sandy loams, some gossamer outcrop.

No. plots 4; Taxa/plot (SD) 18.0 ± 2.4; Altitude 152–318 m.

Diagnostic taxa (INDVAL): Shrubs *Labichea lanceolata* subsp. *brevifolia* (48), *Hibbertia mucronata* (45), *Hovea acanthoclada* (35), *Lasiopetalum rosmarinifolium* (30), *Melaleuca pentagona* (26). **Herbs** *Lomandra mucronata* (16).

Most common taxa (% frequency): Trees *Eucalyptus falcata* subsp. *falcata* (75), *E. pleurocarpa* (75), *E. incrassata* (25), *E. megacornuta* (25), *E. thamnoides* subsp. *megista* (25). **Shrubs** *Hovea acanthoclada* (100), *Labichea lanceolata* subsp. *brevifolia* (75), *Lasiopetalum rosmarinifolium* (75), *Calothamnus quadrifidus* (50), *Hibbertia mucronata* (50), *Melaleuca pentagona* (50), *Acacia gonophylla* (25), *A. subcaerulea* (25), *Acrotricha ramiflora* (25), *Allocasuarina acutivalvis* subsp. *acutivalvis* (25), *A. humilis* (25),



Figure 18. *Eucalyptus megacornuta* open forest was an unusual structural unit in Community 16 (survey plot R045).

Astroloba serratifolium (25), *Banksia heliantha* (25), *B. lemanniana* (25), *Beyeria brevifolia* (25), *Calothamnus pinifolius* (25), *C. roseus* (25), *Chorizema trigonum* (25), *Grevillea patentiloba* subsp. *platypoda* (25), *Hakea laurina* (25), *H. lissocarpa* (25), *H. obtusa* (25), *H. verrucosa* (25), *Hibbertia atrichosepala* (25), *Leucopogon carinatus* (25), *L. sp.* *Coujinup*/sp. *Newdegate* (25), *Spyridium majoranifolium* (25), *Taxandria spathulata* (25). **Herbs** *Lomandra mucronata* (100), *Lepidosperma* sp. Saltbush Hill [KR Newbey 4118] (50), *Platysace maxwellii* (50), *Tetraria* sp. Mt Madden [CD Turley 40 BP/897] (50), *Amphipogon turbinatus* (25), *Austrodanthonia setacea* (25), *Billardiera coriacea* (25), *Cassytha glabella* (25), *Conostylis argentea* (25), *Dampiera lavandulacea* (25), *Dianella revoluta* var. *revoluta* (25), *Lepidosperma* sp. Bandalup Scabrid [N Eveleigh 10798] (25), *L. sp.* Cordingup [GF Craig 6138] (25), *L. sp.* [AS George 9935] (25), *L. sp.* [RL Barrett 3553] (25), *Lomandra micrantha* subsp. *teretifolia* (25), *Marianthus bicolor* (25), *Neurachne alopecuroidea* (25), *Stylium albomontis* (25).

Short range endemics

0–10 km²: *Hibbertia atrichosepala*.

10–100 km²: *Calothamnus roseus*.

1000–10,000 km²: *Eucalyptus megacornuta*, *Grevillea patentiloba* subsp. *platypoda*.

Conservation taxa: *Calothamnus roseus* (P1), *Hibbertia atrichosepala* (P1).

Community 17: Northern lateritic *Eucalyptus pleurocarpa*/*E. falcata* subsp. *falcata*/*E. spp.* mallee shrublands.

Map unit(s): Efal/Eple (17), Dcir (8), Dfol (2), Mosaic Efal/Eple & Eunc/Einc (2), Blac/Borb (1), Eunc/Esp (2), Mosaic Blac/Borb & Efal/Eple (1).

This community typically consisted of an overstorey of sparse mallees

dominated by *Eucalyptus pleurocarpa* and by other eucalypt species, including *E. falcata* subsp. *falcata*, *E. uncinata* and, less frequently, *E. incrassata*. The mid stratum of mid-dense to dense shrubland was very species rich and was dominated by a diverse array of Proteaceous and Myrtaceous taxa, and some sites were dominated by *Banksia cirsoides* and *B. pallida*, while a few others were dominated by dense stands of *B. foliosissima*. Community 17 was found on gently to moderately inclined hillslopes, generally ranging from the midslopes to ridge crests, occurring from the central to the northern extent of the main ridges of the Ravensthorpe Range.

Soils: rocky, sandy loams with an abundance of surface ferruginous pea gravel with limited exposure of lateritic bedrock.

No. plots 33; Taxa/plot (SD) 41.8 ± 1.5; Altitude 237–438 m.

Diagnostic taxa (INDVAL): Trees *Eucalyptus pleurocarpa* (17). **Shrubs** *Banksia pallida* (61), *Mcleuca villosiscapala* (54), *Banksia cirsoides* (42), *Leptospermum* sp. Bandalup Hill [G Cockerton 11001] (35), *Adenanthes flavidiflora* (33), *Beaufortia micrantha* var. *micrantha* (33), *Daviesia corymbosa* (30), *Hakea cygna* subsp. *cygna* (30), *Petrophile glauca* (30), *Jacksonia viscosa* (29), *Boronia crassifolia* (28), *Astroloba prostratum* (27), *Guichenotia anota* (27), *Hemigenia teretiuscula* (27), *Lysinema pentapetalum* (27), *Allocasuarina humilis* (26), *Persoonia striata* (26), *Goodenia pinifolia* (25), *Stachystemon brachyphyllus* (24), *Beaufortia schaueri* (22), *Banksia corviflora* (21), *Leucopogon cuneifolius* (21), *Hibbertia gracilipes* (20), *Banksia foliosissima* (18), *Hakea subsulcata* (18), *Isopogon* sp. Ravensthorpe [DB Foreman 1207] (16), *Petrophile seminuda* (15). **Herbs** *Dampiera juncea* (60), *Lepidosperma* sp. [RL Barrett 3570] (41), *Schoenus brevisetis* (36), *Drosera grievei* (33), *Cassytha pomiformis* (26), *Mesomelacina stygia* subsp. *stygia* (20), *Lepidobolus chactocephalus* (18), *Amphipogon turbinatus* (14), *Conostylis argentea* (14).

Most common taxa (% frequency): Trees *Eucalyptus pleurocarpa* (94), *E. falcata* subsp. *falcata* (70), *E. uncinata* (48), *E. incrassata*



Figure 19. Community 17 typically occurred as species-rich mallee shrubland (survey plot R242) and was largely restricted to laterite ridges in the north of the study area.

(39). **Shrubs** *Hibbertia gracilipes* (85), *Melaleuca villosiscapa* (85), *Leptospermum* sp. Bandalup Hill [G Cockerton 11001] (79), *Beaufortia schaueri* (76), *Allocasuarina humilis* (73), *Banksia cirsoides* (73), *Boronia crassifolia* (70), *Banksia pallida* (61), *Lysinema pentapetalum* (61), *Banksia lemanniana* (58), *Beaufortia micrantha* var. *micrantha* (58), *Goodenia pinifolia* (55), *Petrophile glauca* (55), *Leucopogon cuneifolius* (52), *Isopogon* sp. Ravensthorpe [D.B. Foreman 1207] (48), *Jacksonia viscosa* (48), *Hakea obtusa* (45), *Taxandria spathulata* (45), *Persoonia striata* (42), *Petrophile seminuda* (39), *Leucopogon* sp. Coujinup/sp. Newdegate (36), *Acacia* sp. Ravensthorpe [RS Cowan & BR Maslin RSC A-760] (33), *Adenanthes flavidiflorus* (33), *Leucopogon concinnus* (33), *Persoonia helix* (33), *Acacia heterochroa* subsp. *heterochroa* (30), *Daviesia coryloba* (30), *Guichenotia anota* (30), *Hakea cygna* subsp. *cygna* (30), *Hemigenia teretiuscula* (30), *Leptospermum spinescens* (30), *Spyridium cordatum* (30), *Astroloba prostratum* (27), *Gompholobium knightianum* (27). **Herbs** *Lomandra mucronata* (70), *Dampiera juncea* (67), *Mesomelena stygia* subsp. *stygia* (64), *Neurachne alopecuroidea* (64), *Amphipogon turbinatus* (61), *Lepidosperma* sp. [RL Barrett 3570] (61), *Conostylis argentea* (58), *Gahnia ancistrophylla* (42), *Cassytha glabella* (39), *Stylium albomontis* (39), *Tetraria* sp. Mt Madden [CD Turley 40 BP/897] (39), *Cassytha pomiformis* (36), *Schoenus brevisetis* (36), *Billardiera venusta* (33), *Drosera grievei* (33), *Lepidosperma* sp. Mt Burdett [MA Burgman & C Layman MAB 3287] (27), *Lomandra micrantha* subsp. *teretifolia* (27).

Short range endemics

10–100 km²: *Banksia corvijuga*, *Daviesia megacalyx*.

100–1000 km²: *Guichenotia anota*, *Isopogon* sp. Ravensthorpe (DB Foreman 1207), *Kunzea cincinnata*, *Leucopogon infuscatus*, *Micromyrtus navicularis*, *Spyridium glaucum*.

1000–10,000 km²: *Acacia heterochroa* subsp. *heterochroa*, *A. laricina* var. *crassifolia*, *Banksia foliosissima*, *B. laevigata* subsp. *laevigata*, *Boronia oxyantha* var. *brevicalyx*, *B. ternata* var. *elongata*, *Drosera grievei*, *Leptospermum* sp. Bandalup Hill (G Cockerton 11001), *Microcorys pimeleoides*.

Conservation taxa: *Banksia corvijuga* (P3), *B. foliosissima* (P4), *B. laevigata* subsp. *laevigata* (P4), *Chorizema ulotropis* (P4), *Daviesia megacalyx* (R), *Drosera grievei* (P1), *Guichenotia anota* (P1), *Micromyrtus navicularis* (P3), *Synaphea drummondii* (P3)

Community 18: Southern lateritic *Eucalyptus pleurocarpa*/*E. falcata* subsp. *falcata* mallee shrublands.

Map unit(s): Efal/Eple (4), Dque (1), Eple/Bmed (1).

Community 18 appeared to be a southern variant of community 17. This community typically consisted of tall, open mallee shrublands dominated by *Eucalyptus pleurocarpa*, *E. falcata* subsp. *falcata* and *E. incrassata*, over a mid-dense to dense, species-rich shrubland of *Banksia lemanniana*, *Taxandria spathulata*, *Banksia heliantha*, *Allocasuarina humilis* and *Leucopogon* sp. Coujinup/sp. Newdegate. Sites of this community were located on the south-eastern half of the Ravensthorpe Range, notably around Kundip and on Bandalup Hill. Sites generally occurred on gently to moderately inclined hillslopes, ranging from lower slopes to hillcrests.

Soils: grey-brown sandy loams, clay sands and loamy sands. Outcropping bedrock was infrequent, and varied.

No. plots 17; Taxa/plot (SD) 38.6 ± 1.7; Altitude 110–280 m.

Diagnostic taxa (INDVAL): **Trees** *Eucalyptus pleurocarpa* (19), *E. incrassata* (16), *E. falcata* subsp. *falcata* (14). **Shrubs** *Xanthorrhoea platyphylla* (53), *Petrophile squamata* subsp. northern [J Monks 40] (38), *Banksia heliantha* (35), *Allocasuarina humilis* (33), *Lysinema pentapetalum* (31), *Taxandria spathulata* (31), *Pultenaea indira* (30), *Jacksonia elongata* (28), *Calothamnus gracilis* (27), *Isopogon trilobus* (27), *Leucopogon* sp. Coujinup/sp. Newdegate (26), *Chorizema*

uncinatum (24), *Gompholobium viscidulum* (24), *Stachystemon virgatus* (23), *Adenanthes oreophilus* (21), *Calothamnus pinifolius* (21), *Banksia lemanniana* (19), *Hakea pandanicarpa* (19), *Isopogon* sp. Fitzgerald River [DB Foreman 813] (19), *Boronia crassifolia* (16), *Hibbertia gracilipes* (16), *Leptospermum* sp. Bandalup Hill [G Cockerton 11001] (16). **Herbs** *Schoenus sublaxus* (37), *Mesomelena stygia* subsp. *stygia* (33), *Amphipogon turbinatus* (22), *Conostylis bealiana* (21), *Lomandra mucronata* (20), *Argentipallium niveum* (19), *Dampiera fasciculata* (19), *Lepidosperma* sp. Mt Burdett [MA Burgman & C Layman MAB 3287] (19).

Most common taxa (% frequency): **Trees** *Eucalyptus pleurocarpa* (100), *E. falcata* subsp. *falcata* (82), *E. incrassata* (71), *E. uncinata* (29). **Shrubs** *Taxandria spathulata* (88), *Allocasuarina humilis* (82), *Banksia heliantha* (82), *Hibbertia gracilipes* (76), *Leucopogon* sp. Coujinup/sp. Newdegate (76), *Banksia lemanniana* (71), *Lysinema pentapetalum/ciliatum* (65), *Boronia crassifolia* (53), *Leptospermum* sp. Bandalup Hill [G Cockerton 11001] (53), *Petrophile squamata* subsp. northern [J. Monks 40] (53), *Xanthorrhoea platyphylla* (53), *Acacia gonophylla* (41), *Beaufortia schaueri* (41), *Chamaesciadium ciliatum* (41), *Jacksonia elongata* (41), *Leucopogon carinatus* (41), *Stachystemon virgatus* (41), *Adenanthes oreophilus* (35), *Calothamnus pinifolius* (35), *Daviesia teretifolia* (35), *Hakea pandanicarpa* (35), *Melaleuca rigidifolia* (35), *Petrophile seminuda* (35), *Pultenaea indira* (35), *Acacia heterochroa* subsp. *heterochroa* (29), *Beaufortia micrantha* var. *micrantha* (29), *Beyeria brevifolia* (29), *Calothamnus gracilis* (29), *Chorizema trigonum* (29), *Hakea laurina* (29), *Isopogon trilobus* (29), *Lasiopetalum rosmarinifolium* (29), *Leptospermum spinescens* (29). **Herbs** *Lomandra mucronata* (88), *Mesomelena stygia* subsp. *stygia* (82), *Amphipogon turbinatus* (76), *Cassytha glabella* (59), *Lepidosperma* sp. Mt Burdett [MA Burgman & C Layman MAB 3287] (53), *Schoenus sublaxus* (53), *Dampiera lavandulacea* (41), *Lomandra micrantha* subsp. *teretifolia* (41), *Conostylis bealiana* (29), *Dampiera fasciculata* (29), *Gahnia ancistrophylla* (29), *Pterostylis vittata* (29).

Short range endemics

0–10 km²: *Kunzea similis* subsp. *mediterranea*.

10–100 km²: *Calothamnus roseus*.

100–1000 km²: *Isopogon* sp. Ravensthorpe (DB Foreman 1207), *Spyridium glaucum*.

1000–10,000 km²: *Acacia heterochroa* subsp. *heterochroa*, *A. pinguiculosa* subsp. *pinguiculosa*, *Boronia oxyantha* var. *brevicalyx*, *Grevillea rigidula* subsp. *distans*, *Leptospermum* sp. Bandalup Hill (G Cockerton 11001).

Conservation taxa: *Calothamnus roseus* (P1), *Kunzea similis* subsp. *mediterranea* (R).

Community 19: Mixed *Eucalyptus flocktoniae*/*E. leptocalyx*/*E. spp* mallee shrubland.

Map unit(s): Eflo/Esug (1).

A range of canopy eucalypt species were recorded for this community type, notably mixed stands of tall, open mallee, including *Eucalyptus suggrandis* subsp. *suggrandis*, *E. leptocalyx*, *E. phaenophylla*, *E. uncinata* and *E. flocktoniae*. Common taxa in the taller understorey included *Hakea verrucosa*, *Banksia media*, and *Melaleuca hamata*. Community 19 was also species-rich and was located on the south-eastern half of the range, on an area of low, undulating hills east of the main ridge. It occurred on hillslopes with low gradients, from the lower to upper slopes of these hills.

Soils: rocky brown clay loams with no exposed bedrock.

No. plots 7; Taxa/plot (SD) 35.0 ± 3.0; Altitude 58–192 m.

Diagnostic taxa (INDVAL): **Trees** *Eucalyptus kessellii* subsp. *cognosta* (38), *E. leptocalyx* (26), *E. suggrandis* subsp. *suggrandis* (22). **Shrubs** *Coopernookia polygalacea* (41), *Styphelia intertexta*




Figure 20. Related to community 17, community 18 was generally species-rich mallee shrublands (survey plot R050) found on laterite ridges in the south of the study area.



Figure 21. Community 19, a species-rich mallee shrubland (survey plot R053) restricted to the south-eastern end of the range.

(40), *Hibbertia exasperata* (39), *Comesperma spinosum* (37), *Boronia inornata* subsp. *leptophylla* (32), *Acacia ingrata* (30), *Banksia media* (29), *Spyridium cordatum* (26), *Pomaderris brevifolia* (24), *Daviesia anceps* (21), *D. articulata* (20), *Melaleuca subfalcata* (20), *Acrotriche cordata* (18), *Grevillea pectinata* (17), *Acacia ophiolithica* (16), *Grevillea oligantha* (15), *Leucopogon infuscatus* (14), *Melaleuca ulicoides* (14). **Herbs** *Cassytha melantha* (14).

Most common taxa (% frequency): **Trees** *Eucalyptus flocktoniae* subsp. *flocktoniae* (86), *E. leptocalyx* (71), *E. suggrandis* subsp. *suggrandis* (71), *E. uncinata* (71), *E. phaenophylla* (57), *E. kessellii* subsp. *cognosta* (43), *E. phenax* subsp. *phenax* (43), *E. pleurocarpa* (29), *E. sp.* Ravensthorpe [AS George 616] (29). **Shrubs** *Acacia ingrata* (100), *Boronia inornata* subsp. *leptophylla* (86), *Daviesia anceps* (86), *Melaleuca hamata* (86), *Spyridium cordatum* (86), *Acrotriche cordata* (71), *Hakea verrucosa* (71), *Lasiopetalum compactum* (71), *Pomaderris brevifolia* (71), *Acacia ophiolithica* (57), *Coopernochla polygalacea* (57), *Daviesia articulata* (57), *Grevillea oligantha* (57), *G. pectinata* (57), *Hakea laurina* (57), *Hibbertia psilocarpa* (57), *Melaleuca ulicoides* (57), *Banksia media* (43), *Comesperma spinosum* (43), *Halgania andromedifolia* (43), *Hibbertia exasperata* (43), *H. pungens* (43), *Leucopogon infuscatus* (43), *Styphelia intertexta* (43), *Boronia inconspicua* (29), *Dodonaea pinifolia* (29), *Exocarpos aphyllus* (29), *Grevillea huegelii* (29), *G. patentiloba* subsp. *patentiloba* (29), *Logania buxifolia* (29), *Melaleuca lateriflora* subsp. *lateriflora* (29), *M. subfalcata* (29), *Microcorys glabra* (29), *Pultenaea purpurea* (29), *Siegfriedia darwinioides* (29), *Templetonia retusa* (29), *Wilsonia humilis* (29). **Herbs** *Cassytha melantha* (71), *Gahnia ancistrophylla* (57), *Lomandra micrantha* subsp. *teretifolia* (57), *Tetraria* sp. Mt Madden [CD Turley 40 BP/897] (57), *Dianella revoluta* var. *revoluta* (43), *Lepidosperma gahnioides* (43), *Cassytha glabella* (29), *Gahnia aristata* (29), *Marianthus bicolor* (29).

Short range endemics

100–1000 km²: *Leucopogon infuscatus*, *Melaleuca ulicoides*.

1000–10,000 km²: *Acacia durabilis*, *Acacia ophiolithica*, *Melaleuca pomphostoma*.

Conservation taxa: none recorded.

Community 20: *Eucalyptus uncinata*/E. *incrassata*/E. spp. mallee shrublands.

Map unit(s): Eunc/Espp (6), Efal/Eplc (5), Mallee/Mspp (3), Mosaic Efal/Eple & Eunc/Einc (3), Esug/Mspp (2), Eunc/Einc (2), Efal/Alca (1), Epro/Mspp (1).

This was a widespread, structurally heterogeneous unit that was closely allied to community 19. It typically consisted of species-rich mallee shrublands dominated by mixtures of *Eucalyptus* species, most commonly including *E. uncinata*, *E. incrassata*, *E. pleurocarpa*, *E. flocktoniae* subsp. *flocktoniae*, *E. phenax* subsp. *phenax* and *E. phaenophylla*. Other eucalypt taxa that were common in the south-eastern sites included *E. pluricaulis* subsp. *pluricaulis*, *E. proxima* and *E. suggrandis* subsp. *suggrandis*. *Hakea laurina* was a common tall shrub and significant indicator species of this community. There was some suggestion of further internal structure associated with a gradation in species composition from southern to northern sites, which could be resolved with further sampling. Community 20 generally occurred on gently inclined slopes, sometimes in minor gullies or minor dry creek lines ranging from lower to upper hillslopes, and was distributed across the north-south extent of the main range and on Bandalup Hill.

Soils: rocky grey-brown soils of a variety of soil textures, from sandy loams, clay sands, to clay loams and clays, with limited outcropping of laterite or metasedimentary bedrock.

No. plots 32; Taxa/plot (SD) 39.1 ± 1.1; Altitude 91–383 m.

Diagnostic taxa (INDVAL): **Trees** *Eucalyptus incrassata* (18), *E. uncinata* (16). **Shrubs** *Eutaxia cuneata* (25), *Hakea lissocarpa* (19), *Melaleuca rigidifolia* (16), *Hakea laurina* (15). **Herbs** *Harperia lateriflora* (19), *Tetraria* sp. Mt Madden [CD Turley 40 BP/897] (17), *Gahnia ancistrophylla* (14).

Most common taxa (% frequency): **Trees** *Eucalyptus uncinata* (84), *E. incrassata* (75), *E. phaenophylla* (59), *E. pleurocarpa* (50), *E. suggrandis* subsp. *suggrandis* (41), *E. falcata* subsp. *falcata* (34), *E. flocktoniae* subsp. *flocktoniae* (34). **Shrubs** *Melaleuca hamata* (94), *Hakea laurina* (69), *Hibbertia pungens* (69), *Hakea lissocarpa* (66), *Daviesia anceps* (56), *Melaleuca rigidifolia* (56), *Baeckea corynophylla* (50), *Eutaxia cuneata* (47), *Spyridium cordatum* (47), *Acacia ingrata* (44), *Hibbertia gracilipes* (44), *Lasiopetalum rosmarinifolium* (44), *Leucopogon* sp. Coujinup/sp. Newdegate (44), *Melaleuca glaberrima* (44), *Acacia gonophylla* (41), *Gompholobium confertum* (41), *Beaufortia schaueri* (38), *Calothamnus quadrifidus* (38), *Boronia inconspicua* (31), *Calytrix leschenaultii* (28), *Rinzia communis* (28), *Acrotriche cordata* (25), *Banksia lemanniana* (25), *Lasiopetalum compactum* (25). **Herbs** *Gahnia ancistrophylla* (81), *Tetraria* sp. Mt Madden [CD Turley 40 BP/897] (78), *Neurachne alopecuroides* (75), *Lomandra micrantha* subsp. *teretifolia* (53), *Cassytha glabella* (44), *Dianella revoluta* var. *revoluta* (44), *Lomandra mucronata* (44), *Stylium albomontis* (44), *Lepidosperma* sp. Bandalup Scabrid [N Eveleigh 10798] (41), *Platysace maxwellii* (41), *Lepidosperma* sp. Saltbush Hill [KR Newbey 4118] (38), *Marianthus bicolor* (38), *Schoenus racemosus* (38), *Amphipogon turbinatus* (31), *Conostylis argentea* (31), *Billardiera venusta* (25), *Dampiera lavandulacea* (25).

Short range endemics

10–100 km²: *Microcybe pauciflora* subsp. *grandis*.

100–1000 km²: *Goodenia phillipsiae*, *Guichenotia apetala*, *Isopogon* sp. Ravensthorpe (DB Foreman 1207), *Kunzea cincinnata*, *Leucopogon infuscatus*, *Marianthus mollis*, *Melaleuca stramentosa*, *Micromyrtus navicularis*, *Pultenaea calycina* subsp. *proxema*, *Spyridium glaucum*.

1000–10,000 km²: *Acacia heterochroa* subsp. *heterochroa*, *A. larinina* var. *crassifolia*, *A. ophiolithica*, *A. pinguisculosa* subsp. *pinguisculosa*, *A. pusilla*, *Banksia foliosissima*, *Boronia oxyantha* var. *brevicalyx*, *Eucalyptus cernua*, *Grevillea patentiloba* subsp. *platypoda*, *Leptospermum* sp. Bandalup Hill (G Cockerton 11001), *Melaleuca pomphostoma*, *Microcorys pimeleoides*.

Conservation taxa: *Banksia foliosissima* (P4), *Goodenia phillipsiae* (P4), *Guichenotia apetala* (P1), *Marianthus mollis* (R), *Microcybe pauciflora* subsp. *grandis* (P3), *Micromyrtus navicularis* (P3), *Pultenaea calycina* subsp. *proxema* (P4), *P. indira* subsp. *monstrosita* (P3), *Stachystemon vinosus* (P4).

Community 21: *Eucalyptus desmondensis*/Allocasuarina spp. tall mallee shrubland.

Map unit(s): Alac (1), Alhu (1), Epro/Mspp (1), Esug/Mspp (1).

Community 21 had some affinities to Communities 19 and 20. This community generally consisted of *Eucalyptus desmondensis* tall mallee shrubland with *Allocasuarina* spp. and *Melaleuca hamata*. This community was found west of the main ridges and with most plots (3) in the southern half of the range. Topographic position ranged from lower to upper hill slopes, on gentle to moderate gradients.

Soils: generally grey-brown to red-brown clay loams, sandy loams and clay sand, exposed bedrock rare.

No. plots 4; Taxa/plot (SD) 30.3 ± 2.5; Altitude 211–404 m.

Diagnostic taxa (INDVAL): **Trees** *Eucalyptus desmondensis* (75). **Shrubs** *Acacia mimica* var. *angusta* (72), *Grevillea dolichopoda* (70), *Gompholobium marginatum* (60), *Baeckea crassiflora* (50), *Leucopogon concinnus* (46), *Melaleuca carrii* (42), *Calytrix*



Figure 22. Community 20, a very species-rich mallee shrubland (survey plot R259) that contained large numbers of short-range endemic species. It was more widespread than the related community 19.



Figure 23. Community 21, *Eucalyptus desmondensis* – *Allocasuarina* tall mallee shrubland (survey plot R074).

leschenaultii (38), *Daviesia pachyphylla* (37), *Chorizema aciculare* subsp. *aciculare* (29), *Allocasuarina campestris* (25), *Hakea lissocarpa* (24), *Logania tortuosa* (22), *Hibbertia pungens* (20), *Astroloba serratifolium* (17). **Herbs** *Schoenus sesquispiculus* (62), *Conostylis argentea* (42), *Lepidobolus preissianus* (35), *Haemodorum discolor* (18), *Schoenus racemosus* (18).

Most common taxa (% frequency): Trees *Eucalyptus desmondensis* (75), *E. uncinata* (50), *Allocasuarina huegeliana* (25), *E. phenax* subsp. *phenax* (25), *E. pluricaulis* subsp. *pluricaulis* (25). **Shrubs** *Grevillea dolichopoda* (100), *Hibbertia pungens* (100), *Melaleuca hamata* (100), *Acacia mimicavar. angusta* (75), *Calytrix leschenaultii* (75), *Gompholobium marginatum* (75), *Hakea lissocarpa* (75), *Leucopogon concinnus* (75), *Allocasuarina campestris* (50), *Astroloba epacridis* (50), *A. serratifolium* (50), *Baeckea crispiflora* (50), *Chorizema aciculare* subsp. *aciculare* (50), *Daviesia pachyphylla* (50), *Dodonaea pinifolia* (50), *Hibbertia gracilipes* (50), *Melaleuca carrii* (50), *Acacia sulcata* (25), *Allocasuarina acutivalvis* subsp. *acutivalvis* (25), *A. hystricosa* (25), *Beaufortia schaueri* (25), *Calothamnus quadrifidus* (25), *Calytrix tetragona* (25), *Daviesia anceps* (25), *D. teretifolia* (25), *Goodenia pinifolia* (25), *Hakea incrassata* (25), *H. marginata* (25), *Halgania anagalloides* var. Southern [AE Orchard 1609] (25), *Kunzea strigosa* (25), *Leucopogon* sp. Coujinup/sp. Newdegate (25), *Logania tortuosa* (25), *Melaleuca elliptica* (25), *M. glaberrima* (25), *M. rigidifolia* (25), *M. villosisepala* (25), *Mirbelia multicaulis* (25), *Persoonia helix* (25), *Petrophile fastigiata* (25), *P. seminuda* (25), *Stachystemon virgatus* (25), *Verticordia acerosa* var. *preissii* (25). **Herbs** *Conostylis argentea* (100), *Gahnia ancistrophylla* (75), *Neurachne alopecuroides* (75), *Schoenus sesquispiculus* (75), *Amphipogon turbinatus* (50), *Lepidobolus preissianus* (50), *Lepidosperma* sp. [RL Barrett 3553] (50), *Mesomelaena stygia* subsp. *stygia* (50), *Schoenus racemosus* (50), *Agrostocrinum scabrum* subsp. *scabrum* (25), *Austrodanthonia setacea* (25), *Cassytha glabella* (25), *Dampiera lavandulacea* (25), *Dianella revoluta* var. *revoluta* (25), *Eriochilus dilatatus* subsp. *undulatus* (25), *Haemodorum discolor* (25), *Lepidosperma* sp. Archer Drive [S Kern & R Jasper LCH 18300] (25), *L. sp. Bandalup Scabrid*[N Evelegh 10798] (25), *L. sp. Cordingup* [GF Craig 6138] (25), *L. sp. Elverdton* [R Jasper et al. LCH 16844] (25), *L. sp. [RL Barrett 2770]* (25), *L. sp. Maydon* [S Kern et al. LCH 17844] (25), *Lomandra mucronata* (25), *Pterochacta paniculata* (25), *Spartochloa scirpoidea* (25), *Stylium dichotomum* (25), *S. piliferum* (25), *Stypandra glauca* (25), *Thysanotus patersonii* (25).

Short range endemics

- 100–1000 km²: *Eucalyptus desmondensis*.
- 1000–10,000 km²: *Allocasuarina hystricosa*.

Conservation taxa: *Allocasuarina hystricosa* (P4), *Eucalyptus desmondensis* (P4).

Unallocated plots

Three plots were not allocated to a community from this analysis. One plot was found to be an outlier in preliminary analyses and was omitted from subsequent analyses. That plot sampled a swamp yate (*Eucalyptus occidentalis*) tall open forest over *Acacia verricula* and *Acacia saligna* subsp. *lindleyi* tall shrubland over *Gahnia ancistrophylla*, *Lomandra effusa* and *Lepidosperma* sp. Hopetoun Road (S Kern et al. LCH 16552) open sedgeland. Seven species in this plot were not recorded in the other 265 plots. This plot appears to represent a distinct swamp yate community that occurs on the alluvial flats around Jerdacuttup Creek.

Two other plots retained in the final analyses did not cluster with other community types (Fig. 2). One plot sampled a *Eucalyptus megacornuta* woodland (plot R010) on outcropping gossan over a very sparse shrub layer

dominated by *Gastrolobium parviflorum* and *Siegfriedia darwiniooides*. This plot was species poor (9 taxa) but most closely allied to community 13. The second plot (R147) sampled an *Acacia pinguiculosa* subsp. *pinguiculosa*, *Kunzea affinis* tall shrubland over *Lepidosperma* sp. (RL Barrett 3553) open sedgeland. Floristically, this site was closest in composition to community 21, but relatively species poor (20 taxa per plot) in comparison. The low species richness found in these plots may have influenced their placement in the analysis.

Correlation with vegetation mapping

There was broad agreement between our communities defined by composition and the vegetation mapping undertaken by Craig et al. (2008; Table 2). Craig et al. (2008) arranged their vegetation mapping units under a soil–landscape classification (Department of Agriculture and Food 2006) and at this level there is good correlation between groups of our communities and the mapping units of the broad soil–landscape classification. Communities 17, 14, 18 and 15 largely correspond to seven vegetation units in the Ravensthorpe System laterite and colluvium group (Table 2). Similarly communities 6, 13, 1, 16, 3, 19, 20, 4 and 2 correspond broadly to the 12 vegetation units in the Ravensthorpe System sedimentary and colluvium group (Table 2). At a finer scale the correspondence is less clear cut.

One vegetation unit (Efal/Eple) covers 28.3% of the mapped area, while the next largest vegetation unit covers just 5% and many are less than 1% (Craig et al. 2008). The large unit was described as heterogeneous and our analysis confirmed this. Most plots in this unit belong to community 17 (18 plots) or community 14 (13 plots), with plots from five other communities being recorded from this unit. While 18 plots of community 17 were mapped as Efal/Eple, a further eight plots were mapped as the Dcir unit of the Kybulup System sandplains (Table 2), suggesting that the Dcir unit does not differ significantly (in terms of composition) from many of our plots in the Efal/Eple unit. Where sufficient replication exists, a number of the vegetation units are split between several of our communities, suggesting that structural-dominance factors contribute significantly to the identification of the mapping units (e.g. Egar, Ecil, Eflo/Ephe; Table 2).

Correlation with PEC mapping

Five Priority Ecological Communities (PECs) have been listed for the Ravensthorpe Range (Department of Environment and Conservation 2010), all of which are listed as Priority 1 PECs (Table 3). These have been described from dominant plant taxa and vegetation structure, and Craig et al. (2008) mapped four of these units in their study area. Our sampling did not include any of the '*Melaleuca* sp. Kundip Heath' (Mx map unit). Three plots from community 14 and one plot from community 17 fell within '*Banksia laevigata* – *B. lemanniana* proteaceous thicket' (Blac/Bor map unit). The

'*Eucalyptus megacornuta* mallet woodland' (Emeg map unit) was found to be floristically heterogeneous, including plots from two different community types and one of our ungrouped plots (R010, Table 3).

No plots were established in the 'Heath on Komatite at Bandalup Hill' (Acop map unit) in the northern part of the range mapped by Craig et al. (2008), but one plot occurred in the mapped area of this community at Bandalup Hill (spatial data supplied by DEC's Species and Communities Unit). This plot was classified as community 9 but the general description of this community outlined in Craig et al. (2008) suggests that a number of our plots in community 8 could also fall within that type.

Three plots that fell within the DEC mapping of the '*Eucalyptus purpurata* woodlands (Bandalup Hill)' community were classified as community 11. These plots form a small cluster within a more broadly-defined floristic unit, with *Eucalyptus purpurata* occurring in only 20% of our plots of this community type, which were more typically open mallee shrubland rather than eucalypt woodland.

DISCUSSION

Recent surveys addressing the biological diversity on the Ravensthorpe Range have demonstrated this region's significance as more than a northern buffer of the Fitzgerald River Biosphere Reserve (Deegan 2006, in Harris et al. 2008), because of the high conservation values specific to the range. It supports a large and diverse array of vascular plant taxa (Craig 2008), a high number of priority and short-range endemic plant species (Kern et al. 2008; this survey), a wide range of vegetation communities (Craig et al. 2008; this survey), and a high number of significant fauna species (Chapman & Newbey 1995; Edward & Harvey 2010). A large proportion of this diversity can be attributed to the location of this landform within the transitional rainfall zone of the SWAFR (Hopper & Gioia 2004) and at a convergence zone of different ecological regions (Beard 1990).

At a finer scale, the patterns of species and communities on the range are strongly correlated with landform, soils and edaphic variables. The Ravensthorpe Range is relatively low in altitude compared with other ranges in the wider region, such as the Stirling Ranges, where there is a clear montane vegetation belt (Pignatti et al. 1993). Nonetheless, the range shows high turnover of edaphic factors and habitats within a small area, due to the interactions of topography and a complex geology. Both this study and previous surveys (Bennett 1987; Chapman & Newbey 1995; Craig et al. 2008) have shown that the floristic communities are closely associated with site physical parameters—notably landform element, geology and soil depth and texture. This is a common pattern found among floristic communities on greenstone and BIF ranges on the Yilgarn Craton (Gibson et al. 2010, 2011), particularly the major floristic segregation between upland and lowland sites (e.g. Markey & Dillon 2008).

The closest upland shrubland communities to the Ravensthorpe Range occur on the Mt Barren quartzites, and while these share conspicuous elements (e.g. *Eucalyptus pleurocarpa*, *E. incrassata*, *E. preissiana*, *E. lehmannii*, *Banksia heliantha*, *Hakea pandanicarpa*, *Banksia lehmanniana*, *Calothamnus pinifolius*, *Allocasuarina humilis* and *Melaleuca pentagona*) there are a number of restricted taxa that clearly distinguish these mallee shrubland communities (e.g. *Banksia foliosissima*, *Acacia ophiolithica*, *Acacia* sp. Ravensthorpe Range [BR Maslin 5463], *Allocasuarina hystricosa*, *Spyridium glaucum* and *Guichenotia anota*). Beard (1972, 1973) recognized these two ranges in different vegetation systems and our data supports this interpretation.

At a more detailed scale our survey showed significant turnover in species composition within upland communities along the range, in particular resolving both a northern and southern laterite shrubland community (communities 17 and 18). There were also indications of an eastern set of floristic communities that were associated with different geological substrates, including weathered ultramafics (communities 8 and 9). This high turnover of floristic communities within and between disjunct ranges of greenstone and banded iron formation has been documented elsewhere in the Yilgarn (Gibson et al. 2007, 2010, 2011).

The unique feature of the Ravensthorpe Range is the extraordinary high number of SREs found on the range, being some six times the number of endemics recorded for the Hatter Hill – Ironcap Range (Gibson 2004) and the Bungabbin Hill area (Gibson et al. 2010). Current analysis of the SREs on the Ravensthorpe Range does not include the very complex *Lepidosperma* group (Barrett et al. 2009) and numbers could be expected to further increase when the taxonomy of this group is resolved.

Vegetation mapping

Despite the very different approaches inherent in plot-based survey and field-based vegetation mapping, a reasonable degree of concordance was seen between the two classifications. The correlation was better at a broader scale but with a few exceptions (primarily the heterogeneous Efal/Eple unit); reasonable correlation can also be seen at a finer scale. Given the complex nature of the vegetation and the small median size of the vegetation polygons (1.4 ha) some of the mismatch between the classifications may be related to boundary placement. No firm conclusion can be reached on the correlation of many of the vegetation units and our communities due to low sampling density; indeed only 40 of the 70 units were sampled. Current available mapping of the PECs suggests that they are variable in terms of floristic composition, with the exception of the '*Eucalyptus purpurata* woodlands (Bandalup Hill)', which form a subset within our community 11.

Conservation and threats

The Ravensthorpe Range has high conservation values of

Table 2

Comparison between the vegetation mapping units of Craig et al. (2008) and the communities determined in this study. In all, 166 plots fell within the area mapped by Craig et al. (2008). Seven plots that fell into mosaic units, and one plot not included in compositional analysis, were excluded from the comparison. Vegetation units are grouped by a soil-landscape classification (Department of Agriculture and Food 2006). Figures indicate the number of plots in each mapped vegetation unit – current community classification combination. A perfect match between vegetation mapping units and communities defined from our current classification would show all plots along the diagonal.

SOIL-LANDSCAPE CLASSIFICATION	Vegetation unit	CURRENT COMMUNITY CLASSIFICATION																			
		17	14	18	15	6	13	1	16	3	19	20	4	2	11	12	21	8	9	5	7
RAVENSTHORPE SYSTEM																					
<i>Laterite & Colluvium</i>																					
Crests & upper slopes	01 Efal/Eple	18	13	4	1		2		1			5									
	03 Efal		1				3														
	04 Efal/Alca		1				1					1									
	05 Dfol	2																			
	06 Dque			1	1																
	07 Blae/Borb	1	3																		
	10 Alac																	1			
<i>Metasedimentary rock & colluvium</i>																					
Crests, upper- & mid-slopes	12 Emeg					1	1														
	13 Egar					2	5	2										1			
	14 Ecli		1				5	2	2											2	
Mid- & lower slopes	15 Eflo/Ephe						4		5												
	16 Eflo/Esgug								1												
	18 Eunc/Einc									2											
	19 Eunc/Essp	1	1								7										
	20 Mallee/Mspp										3										
	22 Mham					2															
Lower slopes	24 Epil/Edep							1									1				
	25 Epil/Edep/Mspp																2				
Valleys	26 Esal																3				

Table 3

Comparison of the potentially threatened Priority Ecological Communities (DEC 2010) with the vegetation mapping unit (Craig et al. 2008) and the community determined by current classification. For each plot occurring in the PEC the community type is indicated.

Priority Ecological Community	Craig et al. 2008 map unit	Current classification
<i>Melaleuca</i> sp. Kundip Heath ¹	Mx	Not sampled
<i>Banksia laevigata</i> – <i>Banksia lemanniana</i> proteaceous thicket	Blae/Bor	3 plots community 14 1 plot community 17
<i>Eucalyptus megacornuta</i> mallet woodland	Emeg	1 plot community 6 1 plot community 13 1 plot unplaced (R010)
Heath on Komatiite at Bandalup Hill ²	Acop	1 plot community 9
<i>Eucalyptus purpurea</i> woodlands (Bandalup Hill) ²	Not mapped	3 plots community 11

1: *Melaleuca* sp. Kundip (GF Craig 6020) is now *Melaleuca sophisma* (Western Australian Herbarium 1998–).

2: Map polygons sourced DEC Species and Communities Unit, Dec 2010.

the same order as the Fitzgerald River National Park. At present, the vegetation on the Ravensthorpe Range is relatively weed-free and appears to be in excellent condition. Bridle creeper (*Asparagus asparagoides*) was the most extensive weed observed and tended to occur in riparian areas. Within intact vegetation, disturbance is localised and is the result of past mining and exploration activities and timber collection. The main biological threat to the floristic communities on the Ravensthorpe Range is *Phytophthora* infestation. Although largely absent from the range, *Phytophthora* is widespread in the adjacent Fitzgerald River National Park and at points near Kundip and south of Bandalup Hill (Harris et al. 2008). Given the number of access tracks into the range, including recreational tracks and tourist scenic drives, there is a very high risk that dieback may be introduced to the main ridges.

It is likely that some floristic communities of the range have been cleared or greatly reduced in area. Much of the lower slopes and colluvial and alluvial flats surrounding the Ravensthorpe Range possess rich clay soils and have been cleared for agriculture. Sharp boundaries occur around the edges of the range, particularly along the western and northern flanks of the range that abut directly onto pasture and wheatfields. Therefore, few colluvial outwash or lower slope sites with intact vegetation could be located in this survey, and it appears that significant expanses of lowland woodlands and forest dominated by tall trees, especially of *Eucalyptus oleosa*, *E. brachycalyx*, *E. loxophleba*, *E. occidentalis* and *E. salmonophloia*, have been lost to clearing (Beard 1972, 1973).

Inappropriate fire regimes are reported to be another threat to floristic communities on the range. Various authors (Beard 1973; Chapman & Newbey 1995; Craig et al. 2008; Harris et al. 2008) have indicated that fire has been an infrequent event on the Ravensthorpe Range during recent historical times, although Aboriginal burning regimes are unknown. This is indicated by extensive stands of long-unburnt woodlands, mallee and shrublands (Craig et al. 2008), and stands of vegetation appear to have not been burnt for at least 75 years (Chapman & Newbey 1995). There is a high number of fire sensitive, obligate seeder species in the vegetation communities, including shrub taxa in upland mallee shrubland communities. Mallets and moort taxa, such as *Eucalyptus clivicola*, *E. megacornuta*, *E. gardneri* subsp. *ravenstorpensis*, *E. platypus* and *E. cernua* are fire-sensitive species that dominate woodlands on hillslopes. Given the topography and concentration of short-range endemic taxa, the Ravensthorpe Range may have acted as a refuge for fire sensitive flora and fauna. Over the past two decades, there has been an increase in the frequency of fires on the range, resulting from lightening strikes, arson and the implementation of prescribed burning regimes (Craig et al. 2008; Harris et al. 2008). Craig et al. (2008) have reported that vegetation dominated by fire-sensitive species is recovering poorly after fire.

Not only are there management issues, but issues of land tenure and land use also impact on the conservation values of the range. Most of the range is covered by mining and exploration tenements (Harris et al. 2008), and both activities are currently underway. There are only two A-class nature reserves (Overshot Hill Nature Reserve and Kundip Nature Reserve), which only cover a small portion of the range and do not encompass the entire diversity of flora and floristic communities (McDonald 2010). A significant portion of the main range has been proposed for inclusion into the conservation estate (Department of Conservation and Land Management 1992; CTRC 1974), but no progress has been made in the decades since these proposals (Chapman & Newbey 1995; Harris et al. 2008). Mining operations are directly affecting some of the restricted SRE taxa, notably *Hibbertia abyssa*, which is restricted to the immediate area of one of the mining pits.

The Ravensthorpe Range represents one of the major biodiversity hotspots in Western Australia that will require active management to maintain these values into the future. This situation will be complicated by the high mineral prospectivity, the risk of dieback becoming established, increased fire frequency and possible climate change impacts. Given the complex and overlapping land use and tenure on the range its management will need the active participation of the local community, industry and government agencies to ensure long-term protection of these values.

ACKNOWLEDGEMENTS

Project funding was made available by the Biodiversity Conservation Initiative of the Department of Environment

and Conservation. Ryonen Butcher, Juliet Wege, Alex Williams, Russell Barrett, Paul Wilson, Bruce Maslin, Mike Hislop, Rob Davis, Malcolm French and Nathan McQuoid are all thanked for providing taxonomic advice and expertise. Karina Knight and Sue Carroll supported this work at the Western Australian Herbarium and provided access to herbarium records. Field work support and technical advice was provided by Maria Lee, Daniel Biddulph, Damien Rathbone, Frederic de Mey, Janet Newell, Mike Fitzgerald, Steven Dillon, Jessica Allen, Caroline McCormick (from DEC); Denise True, Geoff Cockerton, Rosemary Jasper, Daniel Brassington, Hayley Huges (from Western Botanical). Both Mathew Inman and Travis Inman (BHPBilliton: Ravensthorpe Nickel Operations) assisted with mine site access. Stephen van Leeuwen (DEC Woodvale), Gillian Craig and Merle Bennett (Ravensthorpe Regional Herbarium) are all acknowledged for their contributions, support and feedback on the project.

REFERENCES

- Barrett R, Barrett M, Wallace M (2009) *Preliminary Assessment of Taxonomic and Conservation Status of Lepidosperma Species (Cyperaceae) from the Greater Ravensthorpe Range. Research Report to Western Australian Department of Conservation*. Report 45 from the Genetics Laboratory, Kings Park and Botanic Garden, Botanic Gardens and Parks Authority, Perth.
- Beard JS (1972) *The Vegetation of the Newdegate and Bremer Bay Areas, Western Australia: Map and Explanatory Memoir 1:250,000 Series*. Vegmap Publications, Perth.
- Beard JS (1973) *The Vegetation of the Ravensthorpe Area, Western Australia: Map and Explanatory Memoir 1:250,000 Series*. Vegmap Publications, Perth.
- Beard JS (1990) *Plant Life of Western Australia*. Kangaroo Press, Perth.
- Bennett EM (1987) 'Ecological relationships in the vegetation Near Mt Desmond, Ravensthorpe Range, Western Australia'. PhD Thesis, University of Western Australia, Perth.
- Bureau of Meteorology (1908–) Climate Statistics for Australian Locations. Available at <http://www.bom.gov.au/climate/averages/>. [Accessed May 2008].
- Chapman A, Newbey KR (1995) A vertebrate fauna survey and some notes on the vegetation of the Ravensthorpe Range, Western Australia. *CALMscience* 1, 465–508.
- Clarke KR, RN Gorley (2006) PRIMER v6: User Manual/Tutorial. PRIMER-E. Ltd, Plymouth.
- Clarke KR, Somerfield PJ, Gorley RN (2008) Testing of null hypotheses in exploratory community analyses: Similarity profiles and biota-environment linkages. *Journal of Experimental Marine Biology and Ecology* 366, 56–69.
- Cockerton G, Craig GF (2000) Flora and vegetation surveying for Ravensthorpe Nickel Project: September–October 2000'. Unpublished report for Sinclair Knight Merz, Perth and Ravensthorpe Nickel Operations Pty Ltd, Perth.
- Craig GF (1999) 'RAV 8, Tectonic Resources NL, vegetation and flora survey'. Unpublished report to Hart, Simpson & Associates, Shenton Park.
- Craig GF (2004) 'Kundip mining leases M74/41, 51, 53 & 135 and P74/153: Vegetation and flora survey'. Unpublished report for Tectonic Resources NL, Subiaco.
- Craig GF (2005) 'Ravensthorpe iron ore project: Declared rare and priority flora survey'. Unpublished report for Resource Mining Corporation Ltd.
- Craig GF (2008) Appendix 14.5 in Harris RJ, Majer JD, Williams C, Buckley A, Stehlik D (2008). *Ravensthorpe Range and Overshot Hill – An Overview of Biodiversity Values, Threats and Conservation*. Sustaining Gondwana Working Paper Series No. 12. Curtin University of Technology, Perth.
- Craig GF, Hickman EJ, Newell J, McQuoid N, Rick AM, Sandiford EM (2008) *Vegetation of the Ravensthorpe Range, Western Australia: Mt Short to Kundip, 1:10,000 Scale*. Department of Environment and Conservation and South Coast Natural Resource Management Inc., Albany, Western Australia.
- CTRC (1974) *Conservation Reserves in Western Australia*. Report of the Conservation Through Reserves Committee to the Environmental Protection Authority, Department of Conservation and Environment, Perth.
- Department of Agriculture and Food (2006) *Soil – Landscape Systems Mapping of the South West of Western Australia*. Agriculture Western Australia, Perth. Version 4, Dec 2006.
- Department of Conservation and Land Management (1992) *South Coast Region Management Plan 1995–2002*. Department of Conservation and Land Management, Perth.
- Department of Environment and Conservation (2010) Priority Ecological Communities for Western Australia. Version 15. <http://www.dec.wa.gov.au/content/view/849/2017/> [accessed Jan 2011].
- Department of Environment and Water Resources (2007) IBRA Version 6.1. www.environment.gov.au/parks.nrs/ibra/version_6-1/. Updated 6th February [accessed June 2009].
- Dufrêne M, Legendre P (1997) Species assemblages and indicator species: The need for a flexible asymmetrical approach. *Ecological Monographs* 67, 345–366.
- Edward KL, Harvey MS (2010) A review of the Australian millipede genus Atelomastix (Diplopoda: Spirostreptida: Iulomorphidae). *Zootaxa* 2371, 1–63.

- Faith DP, Minchin PR, Belbin L (1987) Compositional dissimilarity as a robust measure of ecological distance. *Vegetatio* **69**, 57–68.
- Gibson N (2004) Flora and vegetation of the eastern goldfields ranges. Part 7, Middle and South Ironcap, Digger Rock and Hatter Hill. *Journal of the Royal Society of Western Australia* **87**, 49–62.
- Gibson N, Keighery GJ, Lyons MN, Webb A (2004) Terrestrial flora and vegetation of the Western Australian Wheatbelt. *Records of the Western Australian Museum*. Supplement No. **67**, 139–189.
- Gibson N, Coates DJ, Thiele KR (2007) Taxonomic research and the conservation status of flora in the Yilgarn Banded Iron Formation ranges. *Nuytsia* **17**, 1–12.
- Gibson N, Yates CJ, Dillon R (2010) Plant communities of the ironstone ranges of south western Australia: hotspots for plant diversity and mineral deposits. *Biodiversity and Conservation* **19**, 3951–3962.
- Gibson N, Meissner R, Markey AS, Thompson WA (2011) Patterns of plant diversity in ironstone ranges in arid south western Australia. *Journal of Arid Environments* **77**, 25–31.
- Harvey MS (2002) Short range endemism among the Australian fauna: some examples from non-marine environments. *Invertebrate Systematics* **16**, 555–570.
- Harris RJ, Majer JD, Williams C, Buckley A, Stehlík D (2008). *Ravensthorpe Range and Overshot Hill – An Overview of Biodiversity Values, Threats and Conservation*. Sustaining Gondwana Working Paper Series No. 12. Curtin University of Technology, Perth.
- Hopper SD, Gioia P (2004) The southwest Australian floristic region: evolution and conservation of a global hot spot of biodiversity. *Annual Review of Ecology, Evolution and Systematics* **35**, 623–650.
- Kern SR, Jasper R, True D (2008) 'Floristic survey of the Ravensthorpe Range, 2007'. Unpublished report prepared for the Department of Environment and Conservation by Western Botanical/Landcare Services.
- McCune B, Mefford MJ (1999) *PC-ORD. Multivariate Analysis of Ecological Data, Version 4*. MjM Software Design, Glenden Beach, Oregon.
- McDonald H (2010) 'Stability of the floristic communities of Overshot Hill Nature Reserve, Ravensthorpe, Western Australia: a comparison of multivariate techniques'. Honours thesis, Curtin University, Perth.
- McDonald RC, Isbell RF, Speight JG, Walker J, Hopkins MS (1998) *Australian Soil and Land Survey: Field Handbook*, 2nd ed. Inkata Press, Melbourne.
- Markey AS, Dillon SJ (2008) Flora and vegetation of the Yilgarn Craton: the central Tallerine Land System. *Conservation Science Western Australia* **7**, 121–149.
- Moir ML, Brennan KEC, Harvey MS (2009) Diversity, endemism and species turnover of millipedes within the south-western Australian global biodiversity hotspot. *Journal of Biogeography* **36**, 1958–1971.
- Myers NR, Mittermeier A, Mittermeier CG, da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* **403**, 853–858.
- Myers NR (2003) Biodiversity hotspots revisited. *BioScience* **53**, 916–917.
- Pignatti E, Pignatti S, Fernando (1993) Plant communities of the Stirling Range, Western Australia. *Journal of Vegetation Science* **4**, 477–488.
- Smith MG (2010) 'Declared Rare and Priority Flora List for Western Australia'. Department of Environment and Conservation, Perth.
- Sneath PHA, Sokal RR (1973) *Numerical Taxonomy: The Principles and Practice of Numerical Classification*. Freeman, San Francisco.
- Watson JR (1991) The identification of river foreshore corridors for nature conservation in the South Coast Region of Western Australia. In *Nature Conservation 2: The Role of Corridors* (eds DA Saunders, RJ Hobbs), pp 63–68. Surrey Beatty and Sons, Chipping Norton.
- Watson J, Wilkins P (1999) The Western Australian South Coast Macro Corridor Project – A bioregional strategy for nature conservation. *Parks* **9**, 7–16.
- Western Australian Herbarium (1998–) Florabase – The Western Australian Flora. Department of Environment and Conservation, <http://florabase.dec.wa.gov.au/> [last accessed January 2011].
- Witt WK (1996) *Ravensthorpe WA Sheet 2930: Western Australian Geological Survey, 1:100,000 Geological Series*. Geological Survey of Western Australia, Department of Minerals and Energy, Perth.
- Witt WK (1997) *Geology of the Ravensthorpe and Cocanarup 1:100,000 Sheets*. Geological Survey of Western Australia. Department of Minerals and Energy, Perth.
- Witt WK (1998) *Geology and Mineral Resources of the Ravensthorpe and Cocanarup 1:100,000 Sheets*. Geological Survey of Western Australia. Report 54. Department of Minerals and Energy, Perth.

APPENDIX 1

Flora list for the Ravensthorpe Range collated from material collected during the plot-based survey and arranged by family. Nomenclature follows online Census of Western Australian Flora (The Western Australian Herbarium 1998–), except for *Lepidosperma*, which follows Barrett et al. (2009). Naturalised species are denoted by an asterisk. Threatened taxa (R), taxa of conservation concern (P1–P4) as listed in Smith (2010), and recent updates as shown on Florabase (Western Australian Herbarium, 1998–).

Aizoaceae	Celastraceae
<i>Disphyma crassifolium</i>	<i>Stackhousia monogyna</i>
Amaranthaceae	<i>Stackhousia scoparia</i>
<i>Ptilotus holosericeus</i>	<i>Tripterococcus brunonis</i>
Apiaceae	Chenopodiaceae
<i>Daucus glochidiatus</i>	<i>Atriplex semibaccata</i>
<i>Platysace deflexa</i>	<i>Atriplex vesicaria</i>
<i>Platysace maxwellii</i>	<i>Chenopodium desertorum</i> subsp. <i>microphyllum</i>
Asparagaceae	<i>Enchyalaena tomentosa</i> var. <i>tomentosa</i>
<i>Asparagus asparagoides</i> *	<i>Maireana enchyalaenoides</i>
<i>Laxmannia paleacea</i>	<i>Maireana marginata</i>
<i>Lomandra collina</i>	<i>Maireana suaedifolia</i>
<i>Lomandra effusa</i>	<i>Rhagodia crassifolia</i>
<i>Lomandra hastilis</i>	<i>Rhagodia preissii</i> subsp. <i>preissii</i>
<i>Lomandra micrantha</i> subsp. <i>teretifolia</i>	<i>Sclerolaena diacantha</i>
<i>Lomandra mucronata</i>	
<i>Thysanotus dichotomus</i>	Colchicaceae
<i>Thysanotus parviflorus</i>	<i>Wurmbea cernua</i>
<i>Thysanotus patersonii</i>	
<i>Thysanotus sparteus</i>	Convolvulaceae
<i>Thysanotus thyrsoideus</i>	<i>Dichondra repens</i>
<i>Thysanotus triandrus</i>	<i>Wilsonia humilis</i>
Asphodeliaceae	
<i>Asphodelus fistulosus</i> *	Cupressaceae
Asteraceae	<i>Callitris drummondii</i>
<i>Argentipallium niveum</i>	<i>Callitris roei</i>
<i>Cassinia arcuata</i>	
<i>Lagenophora huegelii</i>	Cyperaceae
<i>Olearia ciliata</i>	<i>Caustis dioica</i>
<i>Olearia imbricata</i>	<i>Gahnia ancistrophylla</i>
<i>Olearia muelleri</i>	<i>Gahnia aristata</i>
<i>Ozothamnus lepidophyllum</i>	<i>Gahnia lanigera</i>
<i>Podolepis rugata</i>	<i>Lepidosperma diurnum</i>
<i>Pterochaeta paniculata</i>	<i>Lepidosperma carphoides</i>
<i>Senecio quadridentatus</i>	<i>Lepidosperma fimbriatum</i>
<i>Sonchus oleraceus</i> *	<i>Lepidosperma gahnioides</i>
<i>Vittadinia cervicalis</i> var. <i>cervicularis</i>	<i>Lepidosperma humile</i>
<i>Vittadinia gracilis</i>	<i>Lepidosperma aff. gracile</i> (ref: Barrett 2009 report)
Boraginaceae	<i>Lepidosperma</i> sp. (RL Barrett 4446) [pp. <i>Lepidosperma aff. pruinosa</i>]
<i>Halgania anagalloides</i> var. Southern (AE Orchard 1609)	<i>Lepidosperma</i> sp. (RL Barrett 2984) [pp. <i>Lepidosperma aff. pruinosa</i>]
<i>Halgania andromedifolia</i>	<i>Lepidosperma</i> sp. (RL Barrett 3570)
Brassicaceae	<i>Lepidosperma</i> sp. (GF Craig 8243)
<i>Lepidium africanum</i> *	<i>Lepidosperma</i> sp. (R Davis 724)
<i>Lepidium rotundum</i>	<i>Lepidosperma</i> sp. (AS George 9935)
Campanulaceae	<i>Lepidosperma</i> sp. (GF Craig 8249)
<i>Lobelia gibbosa</i>	<i>Lepidosperma</i> sp. (RLBarrett 2770)
Casuarinaceae	<i>Lepidosperma</i> sp. (RL Barrett 3553)
<i>Allocasuarina acutivalvis</i> subsp. <i>acutivalvis</i>	<i>Lepidosperma</i> sp. (RL Barrett 2766)
<i>Allocasuarina campestris</i>	<i>Lepidosperma</i> sp. (RL Barrett 3476)
<i>Allocasuarina huegeliana</i>	<i>Lepidosperma</i> sp. (RL Barrett 3522)
<i>Allocasuarina humilis</i>	<i>Lepidosperma</i> sp. A2 Inland Flat (GJ Keighery 7000)
<i>Allocasuarina hystricosa</i>	<i>Lepidosperma</i> sp. Archer Drive (S Kern & R Jasper LCH 18300)
<i>Allocasuarina microstachya</i>	<i>Lepidosperma</i> sp. Bandalup Scabrid (N Eveleigh 10798)
<i>Allocasuarina scleroclada</i>	<i>Lepidosperma</i> sp. Carracarrup Creek (S Kern, R Jasper, D Brassington LCH 16738)
<i>Allocasuarina spinosissima</i>	<i>Lepidosperma</i> sp. Cordingup (GF Craig 6138)
<i>Allocasuarina thuyoides</i>	<i>Lepidosperma</i> sp. Elverdton (R Jasper et al. LCH 16844)
	<i>Lepidosperma</i> sp. Hopetoun Road (S Kern et al. LCH 16552)

- Lepidosperma* sp. Maydon (S Kern et al. LCH 17844)
Lepidosperma sp. Mt Burdett (MA Burgman & C. Layman MAB 3287)
Lepidosperma sp. Mt Chester (S Kern et al. LCH 16596)
Lepidosperma sp. Mt Short (S Kern et al. LCH 17510)
Lepidosperma sp. Ravensthorpe (GF Craig 5188)
Lepidosperma sp. Saltbush Hill (KR Newbey 4118)
Lepidosperma sp. Shoemaker Levy (L Ang & O Davies 10815A)
Lepidosperma sp. Steere River (S Kern et al. LCH 17764)
Mesomelaena stygia subsp. *stygia*
Schoenus aff. *sesquispiculus* (S Kern & R Jasper LCH 17455)
Schoenus aff. *subbarbatus* (S Kern & R Jasper LCH 17214)
Schoenus breviculmis
Schoenus brevisetis
Schoenus obtusifolius
Schoenus pleiostemoneus
Schoenus racemosus
Schoenus sesquispiculus
Schoenus subbarbatus
Schoenus subfascicularis
Schoenus subflavus subsp. *hispid culms* (KR Newbey 8278)
Schoenus subflavus subsp. long leaves (KL Wilson 2865)
Schoenus subflavus subsp. *subflavus*
Schoenus sublaxus
Tetaria sp. Mt Madden (CD Turley 40 BP/897)
- Dilleniaceae**
- Hibbertia abyssa* (R)
Hibbertia atrichosepala (P1)
Hibbertia exasperata
Hibbertia gracilipes
Hibbertia hamulosa
Hibbertia mucronata
Hibbertia psilocarpa
Hibbertia pungens
Hibbertia rostellata
Hibbertia rupicola
Hibbertia verrucosa
- Droseraceae**
- Drosera grievei* (P1)
Drosera macrantha subsp. *macrantha*
- Elaeocarpaceae**
- Tetrahiteca appplanata* (P1)
- Ericaceae**
- Acrotriche cordata*
Acrotriche parviflora
Acrotriche ramiflora
Acrotriche orbicularis (P1)
Andersonia aff. *lehmanniana* (S Kern & R Jasper LCH 17769)
Andersonia parvifolia
Astroloma epacridis
Astroloma prostratum
Astroloma serratifolium
Brachyloma geissoloma subsp. *geissoloma*
Coleanthera myrtoides
Leucopogon aff. *opponens* (S Kern & R Jasper LCH 17971)
Leucopogon carinatus
Leucopogon concinnus
Leucopogon cuneifolius
Leucopogon denticulatus
Leucopogon fimbriatus
Leucopogon hamulosus
- Leucopogon infuscatus*
Leucopogon lloydiorum
Leucopogon sp. Coujinup (MA Burgman 1085) [pp. *Leucopogon conostephoides*]
Leucopogon sp. Newdegate (M Hislop 3585) [pp. *Leucopogon conostephoides*]
Leucopogon tamminensis var. *australis*
Lissanthe pleurandrodes
Lysinema ciliatum
Lysinema pentapetalum
Styphelia intertexta
Styphelia pulchella
- Euphorbiaceae**
- Beyeria brevifolia*
Beyeria cockertonii (R)
Beyeria lechenaultii
Beyeria sulcata var. *brevipes*
Beyeria villosa (P4)
Monotaxis grandiflora var. *grandiflora*
Monotaxis paxii
Stachystemon brachyphyllus
Stachystemon polyandrus
Stachystemon vinosus (P4)
Stachystemon virgatus
- Fabaceae**
- Acacia* aff. *binata*
Acacia aff. *ferocior* (S Kern LCH 18424)
Acacia bifaria (P3)
Acacia binata
Acacia brachyclada
Acacia chrysella
Acacia chrysocephala
Acacia crassuloides
Acacia crispula
Acacia cupularis
Acacia curvata
Acacia cyclops
Acacia deficiens
Acacia dermatophylla
Acacia diaphyllodinea
Acacia disticha
Acacia durabilis
Acacia erinacea
Acacia glaucoptera
Acacia gonophylla
Acacia grisea (P4)
Acacia harveyi
Acacia heterochroa subsp. *heterochroa*
Acacia ingrata
Acacia lachnophylla
Acacia laricina var. *crassifolia*
Acacia lasiocalyx
Acacia lasiocarpa var. *bracteolata*
Acacia mimica var. *angusta*
Acacia moirii subsp. *dasyacarpa*
Acacia mutabilis subsp. *mutabilis*
Acacia octonervia
Acacia ophiolithica
Acacia patagiata
Acacia pinguiculosa subsp. *pinguiculosa*
Acacia pravifolia
Acacia pusilla
Acacia saligna subsp. *lindleyi*
Acacia sp. Ravensthorpe (RS Cowan & BR Maslin RSC A-760)
Acacia sp. Ravensthorpe Range (BR Maslin 5463) (P1)
Acacia sphacelata subsp. *recurva*
Acacia sphacelata subsp. *sphacelata*

<i>Acacia subcaerulea</i>		<i>Templetonia retusa</i>
<i>Acacia sulcata</i> var. <i>planoconvexa</i>		<i>Templetonia sulcata</i>
<i>Acacia sulcata</i> var. <i>platyphylla</i>		
<i>Acacia verricula</i>		
<i>Bossiaeae preissii</i>		
<i>Chorizema aciculare</i> subsp. <i>aciculare</i>		
<i>Chorizema nervosum</i>		
<i>Chorizema trigonum</i>	(P4)	
<i>Chorizema ulotropis</i>		
<i>Chorizema uncinatum</i>		
<i>Daviesia anceps</i>		
<i>Daviesia articulata</i>		
<i>Daviesia benthamii</i> subsp. <i>acanthoclada</i> x subsp. <i>benthamii</i>		
<i>Daviesia benthamii</i> subsp. <i>acanthoclona</i>		
<i>Daviesia benthamii</i> subsp. <i>benthamii</i>		
<i>Daviesia dilatata</i>		
<i>Daviesia emarginata</i>	(R)	
<i>Daviesia euryloba</i>		
<i>Daviesia lancifolia</i>		
<i>Daviesia megacalyx</i>		
<i>Daviesia mollis</i>		
<i>Daviesia nematophylla</i>		
<i>Daviesia pachyphylla</i>		
<i>Daviesia retrorsa</i>		
<i>Daviesia teretifolia</i>		
<i>Dillwynia acerosa</i>		
<i>Dillwynia divaricata</i>		
<i>Eutaxia cuneata</i>		
<i>Eutaxia microphylla</i> var. <i>diffusa</i>		
<i>Gastrolobium congestum</i>		
<i>Gastrolobium crassifolium</i>		
<i>Gastrolobium latifolium</i>		
<i>Gastrolobium musaceum</i>		
<i>Gastrolobium parviflorum</i>		
<i>Gastrolobium racemosum</i>		
<i>Gastrolobium tetragonophyllum</i>		
<i>Gastrolobium venulosum</i>		
<i>Gompholobium baxteri</i>		
<i>Gompholobium confertum</i>		
<i>Gompholobium cyaninum</i>		
<i>Gompholobium knightianum</i>		
<i>Gompholobium marginatum</i>		
<i>Gompholobium viscidulum</i>		
<i>Goodia medicaginea</i>		
<i>Hovea acanthoclada</i>		
<i>Hovea trisperma</i>		
<i>Indigofera australis</i>		
<i>Jacksonia alata</i>		
<i>Jacksonia elongata</i>		
<i>Jacksonia viscosa</i>		
<i>Kennedia nigricans</i>		
<i>Labichea lanceolata</i> subsp. <i>brevifolia</i>		
<i>Mirbelia multicaulis</i>		
<i>Mirbelia ovata</i>		
<i>Otion microphyllum</i>		
<i>Pultenaea calycina</i> subsp. <i>proxena</i>	(P4)	
<i>Pultenaea craigiana</i>	(P3)	
<i>Pultenaea indira</i> subsp. <i>indira</i>		
<i>Pultenaea indira</i> subsp. <i>monstrosita</i>	(P3)	
<i>Pultenaea purpurea</i>		
<i>Pultenaea rotundifolia</i>		
<i>Pultenaea wudjariensis</i>	(P1)	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>		
<i>Senna artemisioides</i> subsp. x <i>artemisioides</i>		
<i>Sphaerolobium daviesioides</i>		
<i>Sphaerolobium racemulosum</i>		
<i>Templetonia battii</i>		
<i>Templetonia neglecta</i>		
Frankeniaceae		
		<i>Frankenia tetrapetala</i>
Goodeniaceae		
		<i>Anthotium humile</i>
		<i>Anthotium rubriflorum</i>
		<i>Coopernochla polygalacea</i>
		<i>Coopernochla strophiolata</i>
		<i>Dampiera angulata</i> subsp. <i>angulata</i>
		(P4)
		<i>Dampiera deltoidea</i>
		<i>Dampiera fasciculata</i>
		<i>Dampiera juncea</i>
		<i>Dampiera lavandulacea</i>
		<i>Dampiera sacculata</i>
		<i>Goodenia affinis</i>
		<i>Goodenia concinna</i>
		<i>Goodenia incana</i>
		<i>Goodenia laevis</i> subsp. <i>humifusa</i>
		<i>Goodenia phillipsiae</i>
		<i>Goodenia pinifolia</i>
		<i>Goodenia scapigera</i> subsp. <i>scapigera</i>
		(P4)
		<i>Goodenia stenophylla</i>
		<i>Goodenia trichophylla</i>
		<i>Goodenia tripartita</i>
		<i>Lechenaultia formosa</i>
		<i>Scaevola bursariifolia</i>
		<i>Scaevola myrtifolia</i>
Gyrostemonaceae		
		<i>Gyrostemon racemiger</i>
		<i>Gyrostemon sessilis</i>
		<i>Gyrostemon subnudus</i>
Haemodoraceae		
		<i>Conostylis argentea</i>
		<i>Conostylis bealliana</i>
		<i>Haemodorum discolor</i>
Haloragaceae		
		<i>Glischrocaryon angustifolium</i>
		<i>Glischrocaryon flavescens</i>
Hemerocallidaceae		
		<i>Agrostocrinum scabrum</i> subsp. <i>scabrum</i>
		<i>Dianella brevicaulis</i>
		<i>Dianella revoluta</i> var. <i>revoluta</i>
		<i>Stypandra glauca</i>
Iridaceae		
		<i>Patersonia juncea</i>
		<i>Patersonia lanata</i> forma <i>lanata</i>
		<i>Patersonia limbata</i>
Lamiaceae		
		<i>Hemigenia teretiuscula</i>
		<i>Microcorys barbata</i>
		<i>Microcorys glabra</i>
		<i>Microcorys loganiacea</i>
		<i>Microcorys obovata</i>
		<i>Microcorys pimeleoides</i>
		<i>Microcorys subcanescens</i>
		<i>Prostanthera baxteri</i>
		<i>Teucrium sessiliflorum</i>
		<i>Westringia cephalantha</i>
		<i>Westringia dampieri</i>
Lauraceae		
		<i>Cassytha glabella</i> forma <i>dispar</i>
		<i>Cassytha melantha</i>
		<i>Cassytha pomiformis</i>

<i>Cassytha racemosa</i>		
Loganiaceae		
<i>Logania buxifolia</i>		
<i>Logania micrantha</i>		
<i>Logania tortuosa</i>		
Malvaceae		
<i>Alyogyne huegelii</i> var. <i>grossulariifolia</i>		
<i>Guichenotia anota</i>	(P1)	
<i>Guichenotia apetala</i>	(P1)	
<i>Guichenotia micrantha</i>		
<i>Lasiopetalum compactum</i>		
<i>Lasiopetalum rosmarinifolium</i>		
<i>Lysiosepalmus involucratum</i>		
<i>Sida calyxhymenia</i>		
<i>Thomasia angustifolia</i>		
<i>Thomasia foliosa</i>		
<i>Thomasia microphylla</i>		
Myrtaceae		
<i>Baeckea corynophylla</i>		
<i>Baeckea crisipflora</i>		
<i>Baeckea latens</i>		
<i>Baeckea latens</i> x <i>corynophylla</i>		
<i>Baeckea pachyphylla</i>		
<i>Baeckea preissiana</i>		
<i>Beaufortia micrantha</i> var. <i>micrantha</i>		
<i>Beaufortia orbifolia</i>		
<i>Beaufortia schaueri</i>		
<i>Calothamnus gibbosus</i>		
<i>Calothamnus gracilis</i>		
<i>Calothamnus pinifolius</i>		
<i>Calothamnus quadrifidus</i>		
<i>Calothamnus roseus</i>	(P1)	
<i>Calytrix depressa</i>		
<i>Calytrix leschenaultii</i>		
<i>Calytrix tetragona</i>		
<i>Chamelaucium ciliatum</i>		
<i>Cyathostemon tenuifolius</i>		
<i>Darwinia diosmoides</i>		
<i>Darwinia</i> sp. Ravensthorpe (GJ Keighery 8030)		
<i>Darwinia vestita</i>		
<i>Eucalyptus aff. rigidula</i> (A Markey 7000)		
<i>Eucalyptus astringens</i> subsp. <i>astringens</i>		
<i>Eucalyptus astringens</i> subsp. <i>redacta</i>		
<i>Eucalyptus</i> x <i>bennettiae</i>	(P4)	
<i>Eucalyptus brachycalyx</i>		
<i>Eucalyptus</i> x. <i>brachycalyx</i>		
<i>Eucalyptus calycogona</i> subsp. <i>calycogona</i>		
<i>Eucalyptus celastroides</i> subsp. <i>virella</i>		
<i>Eucalyptus cernua</i>		
<i>Eucalyptus</i> x. <i>cernua</i> (S Kern & R Jasper LCH 17803)		
<i>Eucalyptus cernua</i> x <i>dielsii</i>		
<i>Eucalyptus clivicola</i>		
<i>Eucalyptus desmondensis</i>	(P4)	
<i>Eucalyptus dielsii</i>		
<i>Eucalyptus eremophila</i> subsp. <i>eremophila</i>		
<i>Eucalyptus</i> x <i>erythrandra</i>		
<i>Eucalyptus extensa</i>		
<i>Eucalyptus falcata</i> subsp. <i>falcata</i>		
<i>Eucalyptus flocktoniae</i> subsp. <i>flocktoniae</i>		
<i>Eucalyptus gardneri</i> subsp. <i>ravensthorpensis</i>		
<i>Eucalyptus incrassata</i> (A Markey 7680)		
<i>Eucalyptus</i> x <i>incrassata</i>		
<i>Eucalyptus indurata</i>		
<i>Eucalyptus kesselii</i> subsp. <i>eugnosta</i>		
<i>Eucalyptus lehmannii</i> subsp. <i>parallela</i>		
<i>Eucalyptus leptocalyx</i>		
<i>Eucalyptus megacornuta</i>		
<i>Eucalyptus myriadena</i> subsp. <i>myriadena</i>		
<i>Eucalyptus occidentalis</i>		
<i>Eucalyptus oleosa</i> subsp. <i>corvina</i>		
<i>Eucalyptus olivina</i>		
<i>Eucalyptus ovularis</i>		
<i>Eucalyptus phaenophylla</i>		
<i>Eucalyptus phenax</i> subsp. <i>phenax</i>		
<i>Eucalyptus pileata</i>		
<i>Eucalyptus platypus</i> subsp. <i>congregata</i>		
<i>Eucalyptus platypus</i> subsp. <i>platypus</i>		
<i>Eucalyptus pleurocarpa</i>		
<i>Eucalyptus pluricaulis</i> subsp. <i>pluricaulis</i>		
<i>Eucalyptus preissiana</i> subsp. <i>preissiana</i>		
<i>Eucalyptus proxima</i>		
<i>Eucalyptus</i> x. <i>proxima</i> (S Kern & R Jasper LCH 16464)		
<i>Eucalyptus proxima</i> x <i>suggrandis</i>		
<i>Eucalyptus purpurata</i>	(R)	
<i>Eucalyptus salmonophloia</i>		
<i>Eucalyptus salubris</i>		
<i>Eucalyptus</i> sp. Ravensthorpe (AS George 616)		
<i>Eucalyptus sporadica</i>		
<i>Eucalyptus stoatei</i>	(P4)	
<i>Eucalyptus suggrandis</i> subsp. <i>suggrandis</i>		
<i>Eucalyptus tenera</i>		
<i>Eucalyptus thamnooides</i> subsp. <i>megista</i>		
<i>Eucalyptus uncinata</i>		
<i>Hypocalymma strictum</i> subsp. <i>strictum</i>		
<i>Kunzea affinis</i>		
<i>Kunzea cincinnata</i>		
<i>Kunzea micromera</i>		
<i>Kunzea preissiana</i>		
<i>Kunzea similis</i> subsp. <i>mediterranea</i>	(R)	
<i>Kunzea strigosa</i>		
<i>Leptospermum erubescens</i>		
<i>Leptospermum maxwellii</i>		
<i>Leptospermum nitens</i>		
<i>Leptospermum</i> sp. Bandalup Hill (G Cockerton 11001)		
<i>Leptospermum spinescens</i>		
<i>Melaleuca acuminata</i> subsp. <i>acuminata</i>		
<i>Melaleuca</i> aff. <i>coccinea</i> (S Kern & R Jasper LCH 18374)		
<i>Melaleuca bracteosa</i>		
<i>Melaleuca brevifolia</i>		
<i>Melaleuca calycina</i>		
<i>Melaleuca carrii</i>		
<i>Melaleuca cliffortioides</i>		
<i>Melaleuca coronicarpa</i>		
<i>Melaleuca cucullata</i>		
<i>Melaleuca eleuterostachya</i>		
<i>Melaleuca elliptica</i>		
<i>Melaleuca glaberrima</i>		
<i>Melaleuca hamata</i>		
<i>Melaleuca johnsonii</i>		
<i>Melaleuca lateriflora</i> subsp. <i>lateriflora</i>		
<i>Melaleuca pauperiflora</i> subsp. <i>pauperiflora</i>		
<i>Melaleuca penicula</i>	(P4)	
<i>Melaleuca pentagona</i> var. <i>latifolia</i>		
<i>Melaleuca pentagona</i> var. <i>pentagona</i>		
<i>Melaleuca pomphostoma</i>		
<i>Melaleuca pungens</i>		
<i>Melaleuca rigidifolia</i>		
<i>Melaleuca societatis</i>		
<i>Melaleuca ulicoides</i>		
<i>Melaleuca stramentosa</i>		
<i>Melaleuca striata</i>		
<i>Melaleuca strobophylla</i>		
<i>Melaleuca subfalcata</i>		
<i>Melaleuca subtrigona</i>		
<i>Melaleuca teuthidooides</i>		
<i>Melaleuca thapsina</i>		

<i>Melaleuca torquata</i>	<i>Austrostipa</i> sp. Carlingup Rd (S Kern & R Jasper LCH 18459)
<i>Melaleuca tuberculata</i> var. <i>macrophylla</i>	<i>Austrostipa</i> sp. Ravensthorpe Range (A Markey & J Allen 6261)
<i>Melaleuca undulata</i>	<i>Austrostipa variabilis</i>
<i>Melaleuca villosipala</i>	<i>Avena fatua</i> *
<i>Micromyrtus imbricata</i>	<i>Lolium perenne</i> *
<i>Micromyrtus navicularis</i>	<i>Neurachne alopecuroidea</i>
<i>Micromyrtus obovata</i>	<i>Spartochloa scirpoidea</i>
<i>Rinzia communis</i>	
<i>Taxandria spathulata</i>	
<i>Verticordia acerosa</i> var. <i>preissii</i>	
<i>Verticordia chrysantha</i>	
<i>Verticordia densiflora</i> var. <i>cespitosa</i>	
<i>Verticordia endlicheriana</i> var. <i>major</i>	
<i>Verticordia grandiflora</i>	
<i>Verticordia inclusa</i>	
<i>Verticordia oxylepis</i>	
<i>Verticordia picta</i>	
<i>Verticordia roei</i> subsp. <i>roei</i>	
Olacaceae	
<i>Olax benthamiana</i>	
Orchidaceae	
<i>Caladenia falcata</i>	<i>Polygalaceae</i>
<i>Caladenia graminifolia</i>	<i>Comesperma ciliatum</i>
<i>Corunastylis tepperi</i>	<i>Comesperma drummondii</i>
<i>Cyanicula aperta</i>	<i>Comesperma polygaloides</i>
<i>Elythranthera brunonis</i>	<i>Comesperma spinosum</i>
<i>Eriochilus dilatatus</i> subsp. <i>undulatus</i>	<i>Comesperma volubile</i>
<i>Pterostylis mutica</i>	
<i>Pterostylis recurva</i>	
<i>Pterostylis sanguinea</i>	
<i>Pterostylis aff barbata</i> (A Markey 7223)	Portulacaceae
<i>Pterostylis</i> sp. Ongerup (KR Newbey 4874) (P4)	<i>Calandrinia eremaea</i>
<i>Pterostylis vittata</i>	
<i>Thelymitra campanulata</i>	Proteaceae
<i>Thelymitra graminea</i>	<i>Adenanthes flavidiflorus</i>
<i>Thelymitra occidentalis</i>	<i>Adenanthes glabrescens</i> subsp. <i>exasperatus</i>
Oxalidaceae	<i>Adenanthes oreophilus</i>
<i>Oxalis perennans</i>	<i>Banksia cirsoides</i>
Phyllanthaceae	(P3)
<i>Phyllanthus calycinus</i>	<i>Banksia corvijuga</i>
<i>Phyllanthus scaber</i>	<i>Banksia erythrocephala</i> var. <i>erythrocephala</i>
<i>Poranthera drummondii</i>	(P4)
<i>Poranthera microphylla</i>	<i>Banksia foliosissima</i>
Pittosporaceae	(P4)
<i>Billardiera coriacea</i>	<i>Banksia heliantha</i>
<i>Billardiera fusiformis</i>	<i>Banksia laevigata</i> subsp. <i>laevigata</i>
<i>Billardiera venusta</i>	(P4)
<i>Cheiranthera brevifolia</i>	<i>Banksia lemanniana</i>
<i>Cheiranthera filifolia</i> var. <i>filifolia</i>	<i>Banksia media</i>
<i>Marianthus bicolor</i>	<i>Banksia obovata</i>
<i>Marianthus microphyllus</i>	<i>Banksia pallida</i>
<i>Marianthus mollis</i>	<i>Banksia violacea</i>
	<i>Grevillea acuaria</i>
	<i>Grevillea anethifolia</i>
	<i>Grevillea coccinea</i> subsp. <i>coccinea</i>
	<i>Grevillea concinna</i> subsp. <i>leemanniana</i>
	<i>Grevillea dolichopoda</i>
	(P4)
	<i>Grevillea fastigiata</i>
	(P3)
	<i>Grevillea fulgens</i>
	<i>Grevillea huegelii</i>
	<i>Grevillea nudiflora</i>
	<i>Grevillea oligantha</i>
	<i>Grevillea patentiloba</i> subsp. <i>patentiloba</i>
	<i>Grevillea patentiloba</i> subsp. <i>platypoda</i>
	<i>Grevillea pectinata</i>
	(P3)
	<i>Grevillea punctata</i>
	<i>Grevillea rigidia</i> subsp. <i>distans</i>
	<i>Grevillea shuttleworthiana</i> subsp. <i>obovata</i>
	(P1)
	<i>Grevillea sulcata</i>
	<i>Hakea commutata</i>
	<i>Hakea corymbosa</i>
	<i>Hakea cygna</i> subsp. <i>cygna</i>
	<i>Hakea gilbertii</i>
	<i>Hakea ilicifolia</i>
	<i>Hakea incrassata</i>
	<i>Hakea laurina</i>
	<i>Hakea lissocarpa</i>
	<i>Hakea marginata</i>
	<i>Hakea multilineata</i>
	<i>Hakea nitida</i>
	<i>Hakea obtusa</i>
	<i>Hakea pandanicarpa</i> subsp. <i>crassifolia</i>
	<i>Hakea pandanicarpa</i> subsp. <i>pandanica</i>
	<i>Hakea scoparia</i> subsp. <i>scoparia</i>
	<i>Hakea subsulcata</i>
	<i>Hakea trifurcata</i>
	<i>Hakea verrucosa</i>

<i>Hakea victoria</i>	<i>Microcybe albiflora</i>
<i>Isopogon polyccephalus</i>	<i>Microcybe multiflora</i> subsp. <i>baccharoides</i>
<i>Isopogon</i> sp. Fitzgerald River (DB Foreman 813)	<i>Microcybe pauciflora</i> subsp. <i>grandis</i>
<i>Isopogon</i> sp. Newdegate (DB Foreman 771)	<i>Microcybe pauciflora</i> subsp. <i>pauciflora</i>
<i>Isopogon</i> sp. Ravensthorpe (DB Foreman 1207)	<i>Nematolepis phebaliodies</i>
<i>Isopogon trilobus</i>	<i>Phebalium ambiguum</i>
<i>Persoonia dillwynioides</i>	<i>Phebalium lepidotum</i>
<i>Persoonia helix</i>	<i>Phebalium obovatum</i>
<i>Persoonia striata</i>	<i>Phebalium tuberculosum</i>
<i>Persoonia teretifolia</i>	<i>Philoteca gardneri</i> subsp. <i>gardneri</i>
<i>Petrophile fastigiata</i>	<i>Rhadinothamnus rufus</i> subsp. <i>amblycarpus</i>
<i>Petrophile glauca</i>	
<i>Petrophile seminuda</i>	
<i>Petrophile squamata</i> subsp. northern (J Monks 40)	
<i>Petrophile teretifolia</i>	
<i>Synaphea</i> sp Southern Ranges (S Kern et al. LCH 17378)	Santalaceae
<i>Synaphea drummondii</i>	<i>Choretrum glomeratum</i> var. <i>glomeratum</i>
<i>Synaphea flabelliformis</i>	<i>Exocarpos aphyllus</i>
<i>Synaphea interioris</i>	<i>Exocarpos sparteus</i>
<i>Synaphea petiolaris</i>	<i>Leptomeria lehmannii</i>
	<i>Leptomeria pachyclada</i>
	<i>Leptomeria pauciflora</i>
	<i>Santalum acuminatum</i>
Pteridaceae	Sapindaceae
<i>Cheilanthes lasiophylla</i>	<i>Dodonaea amblyphylla</i>
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	<i>Dodonaea bursariifolia</i>
Restionaceae	<i>Dodonaea caespitosa</i>
<i>Desmocladus flexuosus</i>	<i>Dodonaea concinna</i>
<i>Harperia lateriflora</i>	<i>Dodonaea pinifolia</i>
<i>Hypolaena fastigiata</i>	<i>Dodonaea ptarmicaefolia</i>
<i>Lepidobolus chaetocephalus</i>	<i>Dodonaea trifida</i>
<i>Lepidobolus preissianus</i> subsp. <i>arcuatus</i>	<i>Dodonaea viscosa</i> subsp. <i>angustissima</i> x subsp. <i>spathulata</i>
Rhamnaceae	Scrophulariaceae
<i>Cryptandra graniticola</i>	<i>Eremophila densifolia</i> subsp. <i>densifolia</i>
<i>Cryptandra minutifolia</i> subsp. <i>brevistyla</i>	<i>Eremophila densifolia</i> subsp. <i>erecta</i>
<i>Cryptandra myriantha</i>	<i>Eremophila densifolia</i> subsp. <i>pubiflora</i>
<i>Cryptandra nutans</i>	<i>Eremophila glabra</i>
<i>Cryptandra pungens</i>	<i>Glycocystis beckeri</i>
<i>Cryptandra wilsonii</i>	
<i>Pomaderris brevifolia</i>	Styliidiaceae
<i>Pomaderris paniculosa</i> subsp. <i>paniculosa</i>	<i>Levenhookia pusilla</i>
<i>Siegfriedia darwinioides</i>	<i>Levenhookia stipitata</i>
<i>Spyridium cordatum</i>	<i>Styliodium albomontis</i>
<i>Spyridium glaucum</i>	<i>Styliodium breviscapum</i>
<i>Spyridium majoranifolium</i>	<i>Styliodium dichotomum</i>
<i>Spyridium</i> sp. Jerdacuttup (A Williams 332)	<i>Styliodium piliferum</i>
<i>Trymalium elachophyllum</i>	<i>Styliodium repens</i>
<i>Trymalium myrtillus</i> subsp. <i>myrtillus</i>	<i>Styliodium spinulosum</i> subsp. <i>spinulosum</i>
Rubiaceae	<i>Styliodium stowardii</i>
<i>Opercularia apiciflora</i>	<i>Styliodium zeicolor</i>
<i>Opercularia vaginata</i>	
Rutaceae	Thymelaeaceae
<i>Boronia albiflora</i>	<i>Pimelea argentea</i>
<i>Boronia crassifolia</i>	<i>Pimelea brachyphylla</i>
<i>Boronia crenulata</i> subsp. <i>obtusa</i>	<i>Pimelea cracens</i>
<i>Boronia inconspicua</i>	<i>Pimelea erecta</i>
<i>Boronia inornata</i> subsp. <i>inornata</i>	<i>Pimelea physodes</i>
<i>Boronia inornata</i> subsp. <i>leptophylla</i>	<i>Pimelea sulphurea</i>
<i>Boronia oxyantha</i> var. <i>brevicalyx</i>	
<i>Boronia penicillata</i>	
<i>Boronia ramosa</i> subsp. <i>anethifolia</i>	Violaceae
<i>Boronia scabra</i> subsp. <i>scabra</i>	<i>Hybanthus floribundus</i> subsp. <i>adpressus</i>
<i>Boronia spathulata</i>	<i>Hybanthus floribundus</i> subsp. <i>floribundus</i>
<i>Boronia subsessilis</i>	
<i>Boronia ternata</i> var. <i>elongata</i>	Xanthorrhoeaceae
	<i>Xanthorrhoea platyphylla</i>
	Zygophyllaceae
	<i>Zygophyllum glaucum</i>

APPENDIX 2

Sorted two-way table of Ravensthorpe Ranges floristic data showing species frequency in the 21 communities described. Only species that occur at frequencies of ≥20% in at least one community are included in the table. Species groups are shown by horizontal lines.

Taxon	Woodlands							Mallee shrublands					Species-rich mallee shrublands							
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
<i>Thomasia microphylla</i>	-	-	-	40	-	-	25	-	-	-	-	-	-	-	-	-	-	14	13	-
<i>Lepidosperma</i> sp. (GF Craig 8249)	-	-	3	-	-	-	25	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Acrotriche plurilocularis</i>	-	-	-	-	-	-	25	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Dodonaea trifida</i>	-	-	3	-	-	-	50	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Daviesia lancifolia</i>	-	-	-	-	-	-	25	-	-	-	-	-	-	17	-	-	-	14	6	-
<i>Melaleuca stramentosa</i>	-	-	-	-	-	-	75	-	-	-	-	-	-	33	-	-	-	-	9	-
<i>Dodonaea amblyophylla</i>	-	-	-	-	-	-	25	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Melaleuca bracteosa</i>	-	-	-	-	-	-	25	-	-	-	-	-	-	17	-	3	-	-	3	-
<i>Acacia patagiata</i>	-	-	-	-	-	-	25	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus astringens</i>	-	-	-	-	-	-	75	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus thamnoides</i> subsp. <i>megista</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	-	-	-	3	-
<i>Calothamnus roseus</i>	-	-	-	-	-	-	2	-	-	-	-	-	-	-	25	-	6	10	-	-
<i>Daviesia benthamii</i>	-	20	6	40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Goodenia laevis</i> subsp. <i>humifusa</i>	-	20	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Dillwynia divaricata</i>	-	-	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Otion microphyllum</i>	-	20	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus eremophila</i> subsp. <i>eremophila</i>	-	-	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Melaleuca johnsonii</i>	-	-	-	-	40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Melaleuca societatis</i>	-	-	-	-	60	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Cryptandra minutifolia</i> subsp. <i>brevistyla</i>	-	-	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Phебалium obovatum</i>	-	-	-	20	8	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-
<i>Eucalyptus tenera</i>	-	-	3	80	8	-	-	-	-	-	-	-	-	4	-	-	-	-	-	-
<i>Eucalyptus pileata</i>	-	20	21	80	50	20	-	-	-	-	-	-	25	-	-	-	-	-	3	-
<i>Melaleuca teuthidoides</i>	-	-	3	60	17	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Melaleuca coronicarpa</i>	-	-	6	40	8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Melaleuca eleuterostachya</i>	-	-	3	20	8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Melaleuca undulata</i>	-	-	3	60	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Microcybe albiflora</i>	-	-	3	40	25	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus platypus</i>	-	-	6	20	50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus clivicola</i>	-	-	6	-	67	-	75	-	-	-	-	25	39	4	-	-	-	-	-	-
<i>Lepidosperma</i> sp. Ravensthorpe (GF Craig 5188)	-	20	3	20	25	-	25	-	-	-	-	-	9	4	-	-	3	-	19	-
Acrotriche sp. Ravensthorpe (S Kern et al. LCH 16953)	13	-	-	-	-	1	-	-	-	-	-	-	-	13	-	-	22	-	-	-
<i>Eucalyptus purpurata</i>	-	-	3	-	-	-	-	-	-	-	-	20	-	-	-	-	-	-	-	-

Appendix 2 (cont.)

Taxon	Woodlands							Mallee shrublands					Species-rich mallee shrublands									
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
<i>Pimelea erecta</i>	-	-	3	-	-	-	-	-	20	-	-	-	4	-	-	-	-	-	-	-	-	-
<i>Beyeria cockertonii</i>	-	-	-	-	-	3	-	60	-	-	-	-	-	-	60	-	-	20	-	-	-	-
<i>Olearia imbricata</i>	-	-	-	-	-	-	-	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pultenaea wudjariensis</i>	-	-	-	-	-	1	-	20	-	-	-	-	-	-	20	-	-	20	-	-	-	-
<i>Melaleuca calycina</i>	-	-	-	-	-	-	-	-	20	-	-	-	-	-	-	-	-	-	-	3	-	-
<i>Cryptandra nutans</i>	-	-	-	-	-	-	-	-	20	-	-	-	-	4	-	-	-	-	-	-	-	-
<i>Leucopogon infuscatus</i>	-	-	-	-	-	-	-	-	20	-	-	-	-	-	50	-	6	-	43	16	-	-
<i>Melaleuca pomphostoma</i>	-	-	3	-	-	-	-	-	40	100	-	-	-	4	-	-	-	-	14	6	-	-
<i>Pultenaea rotundifolia</i>	-	-	9	-	-	-	-	-	40	-	-	-	-	-	-	-	-	-	-	14	6	-
<i>Choretrum glomeratum</i> var. <i>glomeratum</i>	-	-	3	-	-	-	-	-	20	-	7	-	-	-	-	-	-	-	-	6	-	-
<i>Melaleuca pentagona</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	33	50	-	6	-	6	-	-
<i>Eriochilus dilatatus</i> subsp. <i>undulatus</i>	-	-	-	-	-	-	-	17	20	-	7	-	4	-	-	-	-	-	14	-	25	-
<i>Banksia media</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	12	43	9	-	-
<i>Hibbertia exasperata</i>	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	-	-	-	43	-	-	-
<i>Comesperma spinosum</i>	-	-	-	-	-	-	-	-	-	-	7	-	-	-	-	-	-	-	43	-	-	-
<i>Eucalyptus kesselii</i> subsp. <i>eugnosta</i>	-	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	43	3	-	-
<i>Styphelia intertexta</i>	-	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	43	-	-	-
<i>Dodonaea concinna</i>	13	-	38	20	8	-	-	-	20	-	40	25	-	-	-	-	-	-	-	-	-	-
<i>Westringia dampieri</i>	13	-	18	-	-	-	-	17	-	-	40	25	4	-	-	-	-	-	-	-	-	-
<i>Exocarpos sparteus</i>	-	-	9	-	-	-	-	-	-	-	27	25	-	4	-	-	6	-	14	9	-	-
<i>Leptomeria pachyclada</i>	-	-	6	-	-	-	-	17	-	-	40	-	-	-	-	-	-	-	14	3	-	-
<i>Nematolepis pheballoides</i>	-	-	-	-	-	-	25	-	40	-	33	25	-	-	-	-	-	-	-	3	-	-
<i>Beyeria villosa</i>	53	-	-	-	-	3	-	-	-	-	-	-	-	-	53	-	-	22	-	-	-	-
<i>Pultenaea calycina</i> subsp. <i>proxena</i>	-	-	-	-	-	-	-	17	20	-	67	-	-	-	-	-	-	-	3	-	-	-
<i>Boronia inornata</i> subsp. <i>inornata</i>	25	20	12	-	-	20	-	-	-	-	67	100	-	-	-	-	-	-	14	3	-	-
<i>Melaleuca pauperiflora</i> subsp. <i>pauperiflora</i>	-	-	18	40	-	-	-	-	20	-	80	25	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus brachycalyx</i>	-	-	12	-	-	-	-	17	20	-	87	50	-	-	-	-	-	-	14	-	-	-
<i>Eucalyptus indurata</i>	-	-	9	-	-	-	-	-	-	-	80	25	-	-	-	-	-	-	-	-	-	-
<i>Logania buxifolia</i>	-	-	-	-	-	-	-	17	40	-	7	-	4	8	-	-	12	-	29	6	-	-
<i>Eucalyptus leptocalyx</i>	-	-	3	-	-	25	-	-	60	-	7	-	-	-	-	-	-	12	71	16	-	-
<i>Templetonia retusa</i>	-	20	6	-	-	25	-	-	20	-	20	-	4	4	-	-	-	-	29	3	-	-
<i>Hybanthus floribundus</i> subsp. <i>adpressus</i>	-	-	15	-	-	20	-	58	40	50	20	-	-	8	17	-	-	-	14	13	-	-
<i>Melaleuca cliffortioides</i>	-	-	3	20	-	-	-	42	60	50	27	-	-	4	-	-	-	-	14	-	-	-
<i>Dodonaea pinifolia</i>	25	-	18	20	-	-	-	92	80	100	13	-	43	8	-	-	-	-	29	13	50	-
<i>Hakea verrucosa</i>	13	-	24	-	-	20	-	75	100	100	47	-	17	31	-	25	-	-	71	6	-	-
<i>Acacia ophiolithica</i>	-	-	12	-	-	-	-	50	60	-	20	-	-	4	-	-	-	-	57	3	-	-
<i>Eucalyptus</i> sp. Ravensthorpe (AS George 616)	20	-	4	-	-	9	19	20	3	-	3	-	-	17	58	-	-	19	-	-	-	-
<i>Grevillea oligantha</i>	-	-	21	20	-	-	-	25	80	-	7	-	-	-	-	-	-	-	57	16	-	-
<i>Halgania andromedifolia</i>	-	20	24	20	-	-	-	25	40	-	60	25	-	-	-	-	-	-	43	-	-	-
<i>Pomaderis brevifolia</i>	-	-	18	-	-	-	-	8	40	-	47	-	13	12	-	-	3	-	71	3	-	-

Taxon	Woodlands							Mallee shrublands					Species-rich mallee shrublands									
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
<i>Grevillea fastigiata</i>	-	-	3	-	-	-	-	42	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Allocasuarina hystricosa</i>	-	-	-	-	-	-	-	25	-	-	-	-	-	-	-	-	-	-	-	-	25	
<i>Wurmbea cernua</i>	-	-	-	-	-	-	-	33	40	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Acacia mutabilis</i> subsp. <i>mutabilis</i>	-	-	3	-	8	-	-	-	-	-	27	-	-	-	-	-	-	-	-	-	-	
<i>Glischrocaryon flavescens</i>	-	-	-	-	-	-	-	17	20	-	13	-	-	-	-	-	-	-	14	3	-	
<i>Gahnia lanigera</i>	-	-	6	-	-	-	-	8	-	-	20	-	-	-	-	-	-	-	-	-	-	
<i>Melaleuca brevifolia</i>	-	-	-	-	-	-	-	-	-	-	20	-	-	-	-	-	-	-	-	-	-	
<i>Verticordia oxylepis</i>	-	-	-	-	-	-	-	8	20	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Goodenia scapigera</i> subsp. <i>scapigera</i>	-	-	-	-	-	-	-	8	-	50	-	-	9	8	-	-	3	24	-	3	-	
<i>Thysanotus patersonii</i>	-	20	9	-	8	-	-	17	20	50	-	-	26	8	-	-	6	6	-	13	25	
<i>Lepidosperma</i> sp. Carracarrup Creek (S Kern, R Jasper, D Brassington LCH 16738)	-	-	-	-	-	4	4	-	3	-	-	19	-	-	20	20	-	15	-	3	13	
<i>Lomandra effusa</i>	-	20	-	-	8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9	-	
<i>Acacia sulcata</i>	13	20	3	-	-	-	-	8	-	100	7	-	17	4	-	-	-	-	-	13	25	
<i>Lepidosperma</i> sp. Mt Short (S Kern et al. LCH 17510)	-	-	-	-	-	-	-	-	-	-	-	-	-	4	33	-	12	6	-	13	-	
<i>Coopernochla polygalacea</i>	-	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	57	19	-	-	
<i>Melaleuca subfalcata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	29	13	-	
<i>Eucalyptus pluricaulis</i> subsp. <i>pluricaulis</i>	13	-	-	-	-	-	-	-	-	-	-	-	-	15	33	-	-	6	-	19	25	
<i>Eucalyptus proxima</i>	-	-	3	-	-	-	-	8	20	-	-	-	-	4	-	-	-	-	-	16	-	
<i>Boronia scabra</i> subsp. <i>scabra</i>	-	-	-	-	-	-	-	17	-	100	-	-	-	-	-	-	-	-	-	9	-	
<i>Melaleuca elliptica</i>	-	-	-	-	-	-	-	17	-	100	-	-	-	-	-	-	-	-	-	-	25	
<i>Pimelea brachyphylla</i>	-	-	-	-	-	-	-	8	-	50	-	-	4	-	-	-	-	-	14	-	-	
<i>Grevillea anethifolia</i>	-	-	-	-	-	-	-	8	-	50	-	-	-	-	-	-	-	-	-	-	-	
<i>Agrostocrinum scabrum</i> subsp. <i>scabrum</i>	-	-	4	-	-	2	-	-	-	-	3	-	-	25	-	6	11	25	-	-	-	
<i>Lepidosperma</i> sp. Maydon (S Kern et al. LCH 17844)	-	-	-	40	-	-	-	8	-	-	-	-	-	-	-	-	-	-	-	-	25	
<i>Stypandra glauca</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	
<i>Mirbelia multicaulis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	
<i>Pterochaeta paniculata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	
<i>Daviesia pachyphylla</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	15	-	-	3	50	
<i>Stylium dichotomum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	
<i>Allocasuarina huegeliana</i>	-	-	-	-	-	-	-	-	-	50	-	-	4	-	-	-	-	-	-	-	25	
<i>Baeckea crispiflora</i>	-	-	-	-	-	-	-	-	-	50	-	-	-	-	-	-	-	-	-	-	50	
<i>Gompholobium marginatum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	6	-	9	75	
<i>Grevillea dolichopoda</i>	-	-	-	-	-	-	-	8	-	-	-	-	-	4	-	-	6	12	-	13	100	
<i>Schoenus sesquispiculus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	12	-	-	3	75	
<i>Melaleuca carrii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	-	3	50	-	
<i>Acacia mimica</i> var. <i>angusta</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	75	
<i>Eucalyptus desmondensis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	75	
<i>Allocasuarina campestris</i>	-	-	-	-	-	-	-	25	-	-	-	-	9	12	-	-	3	-	-	3	50	
<i>Lepidobolus preissianus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	12	-	-	6	50	

<i>Eucalyptus megacornuta</i>	-	-	-	-	20	-	-	-	-	-	-	4	-	-	25	-	-	-	-	-	-
<i>Persoonia teretifolia</i>	-	20	3	20	8	40	-	8	-	50	20	-	9	27	-	-	3	-	14	6	-
<i>Eucalyptus lehmannii</i> subsp. northern (M French 425)	-	-	-	-	-	-	-	-	-	-	-	-	8	50	-	6	12	-	3	-	-
<i>Boronia oxyantha</i> var. <i>brevicalyx</i>	-	-	-	-	-	80	75	-	-	-	-	9	-	67	-	3	6	-	6	-	-
<i>Spyridium glaucum</i>	-	-	9	-	-	60	25	-	-	-	-	13	12	17	-	3	6	-	3	-	-
<i>Andersonia</i> aff. <i>lehmanniana</i> (S Kern & R Jasper LCH 18374)	-	-	-	-	-	-	-	-	-	-	-	-	-	33	-	3	-	-	-	-	-
<i>Daviesia mollis</i>	-	-	-	-	-	-	-	-	-	-	-	-	4	33	-	-	6	-	6	-	-
<i>Microcorys glabra</i>	-	-	15	-	-	-	-	-	20	50	-	-	4	-	-	-	-	29	9	-	-
<i>Siegfriedia darwiniooides</i>	-	-	6	-	-	-	25	-	-	-	-	22	12	33	-	-	-	29	16	-	-
<i>Billardiera coriacea</i>	-	20	-	-	-	-	-	-	-	-	-	-	4	17	25	-	-	-	-	-	-
<i>Spyridium majoranifolium</i>	-	-	-	-	-	-	-	-	-	-	-	-	4	33	25	-	-	-	-	-	-
<i>Pterostylis sanguinea</i>	-	-	-	20	-	-	-	-	-	-	-	-	-	-	-	6	6	-	3	-	-
<i>Boronia ternata</i> var. <i>elongata</i>	-	-	-	-	-	-	-	-	-	-	-	4	27	-	-	9	-	-	-	-	-
<i>Hakea multilineata</i>	-	-	-	-	-	-	-	-	-	-	-	4	23	-	-	-	-	-	3	-	-
<i>Hibbertia atrichosepala</i>	-	-	-	-	-	2	15	-	-	-	-	-	-	25	-	-	10	-	-	-	-
<i>Kunzea strigosa</i>	-	-	-	-	-	2	-	-	-	-	-	-	-	-	25	-	6	11	25	-	-
<i>Grevillea concinna</i> subsp. <i>leemanniana</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	33	-	-	-	-	-	-	-
<i>Lepidosperma</i> sp. (AS George 9935)	-	-	-	-	-	-	-	-	-	-	-	-	-	33	25	-	6	-	-	-	-
<i>Drosera macrantha</i> subsp. <i>macrantha</i>	-	-	-	20	-	-	-	8	-	-	-	-	-	-	-	6	24	-	3	-	-
<i>Beyeria brevifolia</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	17	25	18	29	-	9	-	-
<i>Dampiera fasciculata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	17	-	-	29	-	-	-	-
<i>Calothamnus gracilis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	29	-	-	-	-
<i>Jacksonia elongata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	17	-	-	41	-	3	-	-
<i>Xanthorrhoea platyphylla</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	53	-	-	-	-	-
<i>Isopogon trilobus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	29	-	3	-	-	-
<i>Schoenus sublaxus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	17	-	6	53	-	-	-	-
<i>Adenanthes oreophilus</i>	-	-	-	-	-	-	-	-	-	-	-	-	8	17	-	-	35	-	-	-	-
<i>Melaleuca striata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	17	-	-	24	-	-	-
<i>Chorizema trigonum</i>	-	-	-	-	-	-	25	-	-	-	-	-	-	-	50	25	-	29	-	19	-
<i>Grevillea nudiflora</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	33	-	-	18	-	6	-
<i>Leucopogon carinatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	50	25	-	41	-	13	-
<i>Patersonia limbata</i>	-	-	-	-	-	-	-	-	-	50	-	-	-	-	-	-	-	12	-	-	-
<i>Calothamnus pinifolius</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	-	35	-	-	-	-
<i>Chorizema uncinatum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	-	-	-	-
<i>Hibbertia mucronata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	50	-	6	-	-	-	-
<i>Pterostylis vittata</i>	-	-	6	-	8	-	-	8	-	-	7	-	-	-	-	-	-	29	-	6	-
<i>Chamelaucium ciliatum</i>	-	-	-	-	-	-	-	17	-	50	-	-	4	8	17	-	12	41	-	19	-
<i>Banksia heliantha</i>	-	-	-	-	-	10	-	-	-	-	-	-	-	83	83	-	82	6	-	3	-
<i>Eucalyptus preissiana</i> subsp. <i>preissiana</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	50	-	-	24	-	-	-	-
<i>Lepidosperma</i> sp. Archer Drive (S Kern & R Jasper LCH 18300)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	-	-	25	-	-
<i>Isopogon</i> sp. Fitzgerald River (DB Foreman 813)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	-	6	-	-	-
<i>Gompholobium viscidulum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	-	-	-	-
<i>Kunzea cincinnata</i>	-	-	-	-	-	-	-	8	-	-	-	4	23	-	-	18	-	-	16	-	-
<i>Melaleuca thapsina</i>	-	-	-	20	-	20	-	-	-	-	-	22	8	-	-	-	-	-	3	-	-

Appendix 2 (cont.)

230

Taxon	Woodlands							Mallee shrublands						Species-rich mallee shrublands							
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
<i>Haemodorum discolor</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9	-	-	-	25
<i>Boronia subsessilis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	33	-	6	-	-	6	-
<i>Logania tortuosa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	25
<i>Hakea marginata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	-	-	19	25
<i>Lepidosperma</i> sp. (RL Barrett 2770)	-	-	-	-	8	-	-	-	-	-	-	-	-	-	-	-	-	12	-	16	25
<i>Lepidosperma carpoides</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	21	24	-	6	-
<i>Conostylis bealiana</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	29	-	13	-	-
<i>Dampiera lavandulacea</i>	-	-	-	-	-	-	-	33	-	-	-	-	-	4	-	25	6	41	-	25	25
<i>Stachystemon virgatus</i>	-	-	-	-	-	-	-	8	-	-	-	-	-	-	-	-	-	41	-	-	25
<i>Petrophile squamata</i> subsp. northern (J Monks 40)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9	53	-	13	-
<i>Argentipallium niveum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	-	6	-	-
<i>Pultenaea indira</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	35	-	6	-	-
<i>Allocasuarina acutivalvis</i> subsp. <i>acutivalvis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	12	17	25	9	-	-	-	25
<i>Cassytha racemosa</i>	-	-	-	-	-	-	-	-	20	-	-	-	-	-	-	-	3	-	-	-	-
<i>Banksia corvijuga</i>	-	-	-	-	-	1	-	-	-	-	-	-	-	21	-	-	7	-	21	-	-
<i>Astrolobia serratifolium</i>	-	-	-	-	-	-	-	33	-	-	-	-	-	4	-	25	18	6	-	9	50
<i>Calytrix leschenaultii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	12	-	-	15	18	-	28	75
<i>Schoenus racemosus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	12	24	14	38	50
<i>Lepidosperma</i> sp. Bandalup Scabrid (N Evelegh 10798)	-	-	9	-	40	11	4	-	-	-	-	41	25	-	41	20	6	5	25	-	-
<i>Melaleuca glaberrima</i>	-	-	-	40	-	-	-	8	-	50	-	-	-	4	-	-	-	-	14	44	25
<i>Rinzia communis</i>	-	-	-	20	-	-	25	-	-	-	-	-	-	15	-	-	-	-	28	-	-
<i>Grevillea patentiloba</i> subsp. <i>patentiloba</i>	-	-	6	-	25	20	25	8	-	-	13	-	4	8	17	-	3	24	29	19	-
<i>Acacia gonophylla</i>	-	-	-	-	-	-	25	-	-	-	-	-	-	4	-	25	-	41	-	41	-
<i>Eutaxia cuneata</i>	-	-	-	20	-	-	-	-	-	-	-	-	-	4	12	-	-	6	-	47	-
<i>Dianella revoluta</i> var. <i>revoluta</i>	-	40	18	20	-	-	-	42	20	-	7	-	4	4	17	25	3	6	43	44	25
<i>Boronia inconspicua</i>	-	-	15	40	17	20	25	-	-	-	-	-	17	15	50	-	6	12	29	31	-
<i>Eucalyptus phaenophylla</i>	-	-	9	60	-	-	25	-	-	-	-	-	9	23	17	-	6	12	57	59	-
<i>Eucalyptus suggrandis</i> subsp. <i>suggrandis</i>	-	20	21	20	8	-	25	-	-	-	-	-	-	-	17	-	6	-	71	41	-
<i>Gompholobium confertum</i>	-	20	-	20	-	-	25	-	-	-	-	-	-	15	-	-	3	6	-	41	-
<i>Lasiopteratum rosmarinifolium</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	12	17	75	9	29	-	44	-
<i>Baeckea corynophylla</i>	-	-	-	20	8	-	50	8	20	-	-	-	-	12	-	-	6	12	14	50	-
<i>Melaleuca rigidifolia</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	8	33	-	21	35	14	56	25
<i>Spyridium cordatum</i>	-	-	3	60	-	-	-	8	-	-	-	-	9	19	17	-	30	-	86	47	-
<i>Cassytha glabella</i>	13	-	3	-	-	-	-	42	20	-	33	-	-	8	33	25	39	59	29	44	25
<i>Hibbertia pungens</i>	13	-	3	40	-	-	-	75	-	100	13	-	43	19	67	-	24	-	43	69	100
<i>Melaleuca hamata</i>	25	40	21	40	25	-	50	67	-	-	13	-	65	65	50	-	3	6	86	94	100
<i>Hakea lissocarpa</i>	-	-	3	-	-	-	-	8	-	-	-	-	-	12	-	25	18	24	-	66	75
<i>Lomandra micrantha</i> subsp. <i>teretifolia</i>	-	20	-	-	-	-	-	8	-	-	-	-	-	-	33	25	27	41	57	53	-
<i>Hakea laurina</i>	-	-	12	20	-	-	25	-	20	-	-	-	26	15	-	25	9	29	57	69	-
<i>Tetraeria capillaris</i>	-	-	3	-	-	-	-	17	-	-	27	-	4	8	50	50	39	18	57	78	-

<i>Neurachne alopecuroides</i>	-	40	-	20	-	-	-	83	-	-	-	-	4	19	-	25	64	24	14	75	75
<i>Eucalyptus incrassata</i>	-	-	-	-	-	-	25	-	-	-	-	-	13	15	33	25	39	71	14	75	-
<i>Eucalyptus uncinata</i>	-	-	-	-	-	-	-	-	40	-	7	-	4	31	67	-	48	29	71	84	50
<i>Gahnia ancistrophylla</i>	-	40	9	40	17	-	-	25	20	-	7	-	-	19	17	-	42	29	57	81	75
<i>Lepidosperma</i> sp. Mt Benson (RL Barrett 3553)	-	-	-	-	-	-	-	17	-	-	-	-	4	46	-	25	18	12	-	9	50
<i>Brachyloma geissoloma</i> subsp. <i>geissoloma</i>	-	-	-	-	-	3	38	-	-	-	-	3	-	-	38	-	6	3	-	3	-
<i>Philotheeca gardneri</i> subsp. <i>Ravensthorpe</i> (GF Craig 6902)	7	-	26	-	-	6	62	-	-	50	-	6	8	-	62	-	-	3	-	3	-
<i>Lepidosperma</i> sp. Cordingup (GF Craig 6138)	-	-	-	-	-	20	-	8	-	-	-	-	26	42	67	25	18	-	14	-	25
<i>Acrotriche ramiiflora</i>	25	-	-	20	-	-	-	-	-	-	-	-	22	35	50	25	21	18	14	9	-
<i>Rhadinothamnus rufus</i> subsp. <i>amblycarpus</i>	-	-	-	-	-	-	-	-	-	-	-	-	22	31	33	-	21	-	-	6	-
<i>Banksia laevigata</i> subsp. <i>laevigata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	31	-	-	21	-	-	-	-
<i>Acacia</i> sp. <i>Ravensthorpe</i> (RS Cowan & BR Maslin RSC A-760)	-	-	-	20	-	-	-	-	-	-	-	-	26	54	-	-	33	12	-	19	-
<i>Hakea obtusa</i>	-	-	-	-	-	-	-	-	-	-	-	-	13	65	100	25	45	18	14	19	-
<i>Beaufortia orbifolia</i>	-	-	-	-	-	20	-	-	-	-	-	-	-	69	67	-	18	-	-	6	-
<i>Isopogon</i> sp. <i>Ravensthorpe</i> (DB Foreman 1207)	-	-	-	-	-	-	-	-	-	-	-	-	4	46	33	-	48	6	-	6	-
<i>Leucopogon cuneifolius</i>	-	-	-	20	-	-	-	-	-	100	-	-	-	35	17	-	52	-	-	6	-
<i>Goodenia pinifolia</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	35	-	-	55	-	-	3	25
<i>Persoonia helix</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	23	-	-	33	-	-	3	25
<i>Santalum acuminatum</i>	25	40	12	-	-	-	-	17	40	50	13	50	43	50	17	-	3	6	-	6	-
<i>Gastrolobium parviflorum</i>	50	20	3	40	17	80	25	-	-	-	-	-	74	62	67	-	6	-	-	16	-
<i>Lasiopetalum compactum</i>	38	20	18	-	33	20	25	8	-	-	-	50	43	38	33	-	18	6	71	25	-
<i>Stylium albomontis</i>	-	-	-	-	8	20	-	17	-	-	-	-	30	65	83	25	39	12	14	44	-
<i>Lepidosperma</i> sp. Saltbush Hill (KR Newbey 4118)	-	25	17	-	-	16	15	-	3	-	-	38	-	50	50	20	24	6	-	21	25
<i>Calothamnus quadrifidus</i>	-	-	-	-	-	-	-	67	-	-	7	-	22	31	-	50	9	6	-	38	25
<i>Platysace maxwellii</i>	25	-	6	20	-	-	-	17	-	50	13	25	61	31	17	50	21	18	-	41	-
<i>Grevillea coccinea</i> subsp. <i>coccinea</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	15	-	-	24	6	-	-	-
<i>Thysanotus sparteus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	12	-	-	21	-	-	9	-
<i>Dampiera angulata</i> subsp. <i>angulata</i>	13	-	-	-	-	-	-	-	-	-	-	-	-	27	17	-	15	-	14	16	-
<i>Marianthus bicolor</i>	-	-	3	-	17	20	-	-	-	-	-	-	17	12	-	25	18	18	29	38	-
<i>Cassytha pomiformis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	8	-	-	36	-	-	6	-
<i>Guichenotia anota</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	30	-	-	-	-
<i>Hakea pandanicarpa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	35	-	6	-
<i>Daviesia teretifolia</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	21	35	-	-	25
<i>Hakea trifurcata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	21	12	-	-	-
<i>Schoenus pleiostemoneus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	18	-	3	-
<i>Adenanthes flavidiflora</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	33	-	-	-	-
<i>Drosera grievei</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	33	-	-	-	-
<i>Lepidobolus chaetocephalus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	21	-	-	3	-
<i>Hakea cygna</i> subsp. <i>cygna</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	30	-	-	-	-
<i>Hemigenia teretiuscula</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	30	-	-	3	-
<i>Hakea incrassata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	18	-	-	-	25
<i>Stachystemon brachyphyllus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	-	-	-	-
<i>Astroloomba prostratum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	27	-	-	-	-
<i>Lepidosperma</i> sp. (RL Barrett 3522)	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	21	18	-	-	-
<i>Schoenus obtusifolius</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	18	-	9	-
<i>Chorizema aciculare</i> subsp. <i>aciculare</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	18	6	-	13	50

Taxon	Woodlands							Mallee shrublands					Species-rich mallee shrublands									
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
<i>Leucopogon concinnus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	33	12	-	3	75	
<i>Acacia heterochroa</i> subsp. <i>heterochroa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	4	33	-	30	29	-	3	-	
<i>Gompholobium knightianum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	27	18	-	3	-	
<i>Billardiera venusta</i>	-	-	-	-	-	5	12	-	-	-	-	25	-	17	33	-	6	7	-	33	-	
<i>Dampiera sacculata</i>	-	-	-	-	-	-	-	8	-	-	-	-	-	4	-	-	24	12	-	22	-	
<i>Leptospermum spinescens</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	19	17	-	30	29	-	9	-	
<i>Lepidosperma</i> sp. Mt Burdett (MA Burgman & C Layman MAB 3287)	13	-	-	-	-	7	4	-	-	-	-	16	-	17	53	-	53	9	-	27	-	
<i>Taxandria spathulata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	8	67	25	45	88	-	22	-	
<i>Banksia lemanniana</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	9	19	50	25	58	71	-	25	-
<i>Beaufortia schaueri</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	15	50	-	76	41	14	38	25	
<i>Leucopogon conostephoides</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	15	-	25	36	76	-	44	25	
<i>Eucalyptus falcata</i> subsp. <i>falcata</i>	13	-	-	-	-	-	25	-	-	-	-	-	-	74	92	33	75	70	82	-	34	-
<i>Eucalyptus pleurocarpa</i>	-	-	-	-	-	-	-	8	-	-	-	-	-	13	85	67	75	94	100	29	50	-
<i>Hibbertia gracilipes</i>	13	-	-	-	-	-	-	8	-	-	-	-	-	4	46	17	-	85	76	14	44	50
<i>Lomandra mucronata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	8	50	100	70	88	-	44	25	
<i>Jacksonia viscosa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	15	-	-	48	18	-	-	-	
<i>Conostylis argentea</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	4	17	25	58	6	-	31	100	
<i>Boronia crassifolia</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	8	33	-	70	53	-	9	-	
<i>Leptospermum</i> sp. Bandalup Hill (G Cockerton 11001)	-	-	-	-	-	9	4	-	-	-	-	6	-	33	79	-	53	7	-	79	-	
<i>Beaufortia micrantha</i> var. <i>micrantha</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	58	29	-	13	-	
<i>Lysinema ciliatum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	61	65	-	9	-	
<i>Amphipogon turbinatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	17	25	61	76	-	31	50	-	
<i>Allocasuarina humilis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	17	25	73	82	-	6	-	-	
<i>Mesomelaena stygia</i> subsp. <i>stygia</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	64	82	-	9	50	
<i>Persoonia striata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	42	24	-	3	-	
<i>Petrophile seminuda</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	39	35	-	3	25	
<i>Schoenus brevisetis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	36	-	-	-	-	
<i>Banksia pallida</i>	-	-	-	-	-	3	-	-	-	-	-	-	-	-	61	-	-	7	-	61	-	
<i>Lepidosperma</i> sp. (RL Barrett 3570)	-	-	-	-	-	-	-	8	-	-	-	-	-	8	-	-	61	-	-	13	-	
<i>Banksia cirsoides</i>	-	-	-	-	-	6	8	-	-	-	-	13	-	33	73	-	-	7	-	73	-	
<i>Petrophile glauca</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	4	33	-	55	-	-	6	-	
<i>Dampiera juncea</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	67	-	-	3	-	
<i>Melaleuca villosisepala</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	8	-	-	85	-	-	16	25	
<i>Banksia foliosissima</i>	-	-	-	-	-	1	-	-	-	-	-	3	-	-	21	-	-	7	-	21	-	
<i>Daviesia euryloba</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	30	-	-	-	-	
<i>Stylium piliferum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	18	24	-	9	25	
<i>Corunastylis tepperi</i>	7	-	-	-	-	-	2	-	-	9	-	-	-	-	-	20	20	6	15	-	-	
<i>Eucalyptus celastroides</i> subsp. <i>virella</i>	-	-	3	-	-	-	-	-	-	-	-	25	-	-	-	-	-	-	-	-	-	
<i>Eucalyptus myriadena</i> subsp. <i>myriadena</i>	-	20	-	-	-	-	-	-	-	-	-	25	-	-	-	-	-	-	-	-	-	
<i>Maireana marginata</i>	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

<i>Austrostipa exilis</i>	13	-	3	-	17	-	-	-	-	-	-	25	4	-	-	-	-	-	-	-	
<i>Acacia</i> sp. Ravensthorpe Range (BR Maslin 5463)	25	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Atriplex semibaccata</i>	-	-	-	-	-	-	-	-	-	-	-	25	4	-	-	-	-	-	-	-	
<i>Melaleuca acuminata</i> subsp. <i>acuminata</i>	-	-	-	40	-	-	-	-	-	-	7	25	13	-	-	-	-	-	3	-	
<i>Acacia cyclops</i>	13	-	-	-	-	-	-	17	-	-	-	25	4	-	-	-	-	-	-	-	
<i>Acacia subcaerulea</i>	13	-	-	-	8	-	-	-	-	-	-	-	13	4	-	25	-	18	-	-	
<i>Hovea acanthoclada</i>	-	-	-	-	17	20	-	-	-	-	-	50	17	8	67	100	-	6	-	-	
<i>Labichea lanceolata</i> subsp. <i>brevifolia</i>	13	-	-	-	-	-	-	-	-	-	-	-	30	-	-	75	-	-	-	-	
<i>Beyeria sulcata</i> var. <i>brevipes</i>	-	-	39	40	-	8	35	-	6	-	-	3	-	17	40	-	12	-	-	13	
<i>Eucalyptus gardneri</i> subsp. <i>ravensthorpensis</i>	25	20	3	-	-	100	-	-	-	-	-	-	39	-	-	-	-	-	-	-	
<i>Grevillea patentiloba</i> subsp. <i>platypoda</i>	50	-	-	-	-	-	-	-	-	-	-	-	13	4	17	25	-	-	3	-	
<i>Phebalium tuberculosum</i>	13	-	-	-	-	-	-	-	-	50	-	-	65	12	-	-	-	-	-	-	
<i>Trymalium elachophyllum</i>	13	-	-	-	-	-	-	-	-	-	-	-	26	4	-	-	-	-	-	-	
<i>Billardiera fusiformis</i>	-	-	13	-	-	3	-	-	9	-	-	-	-	25	20	-	14	-	-	25	
<i>Austrodanthonia setacea</i>	63	40	6	-	8	-	-	8	-	-	-	-	39	4	-	25	-	-	9	25	
<i>Austrostipa hemipogon</i>	63	60	3	-	8	-	-	8	-	-	-	-	30	-	-	-	-	-	3	-	
<i>Sonchus oleraceus</i>	25	-	-	-	-	-	-	-	-	-	7	-	-	-	-	-	-	-	-	-	
<i>Daucus glochidiatus</i>	25	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Sclerolaena diacantha</i>	25	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Enchytraea tomentosa</i> var. <i>tomentosa</i>	38	20	3	-	-	-	-	-	-	-	-	25	4	-	-	-	-	-	-	-	
<i>Austrostipa acrociliata</i>	38	20	6	-	-	-	-	-	-	-	-	-	4	-	-	-	-	-	-	-	
<i>Austrostipa</i> sp. Carlingup Rd (S Kern & R Jasper LCH 18459)	63	20	9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Vittadinia gracilis</i>	25	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Oxalis perennans</i>	50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Ptilotus holosericeus</i>	63	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Eucalyptus oleosa</i> subsp. <i>corvina</i>	25	-	6	-	17	-	-	-	-	-	7	100	-	-	-	-	-	-	-	-	
<i>Senna artemisioides</i> subsp. x <i>artemisioides</i>	75	40	12	-	8	20	-	8	-	-	7	75	22	-	-	-	-	-	3	-	
<i>Acacia erinacea</i>	50	100	12	-	17	40	-	-	-	-	-	25	-	-	-	-	-	-	-	-	
<i>Austrostipa elegantissima</i>	38	80	3	-	-	-	-	-	-	-	7	50	17	-	-	-	-	-	-	-	
<i>Olearia muelleri</i>	38	100	3	-	25	-	-	-	-	-	-	25	-	-	-	-	-	-	-	-	
<i>Asparagus asparagoides</i>	-	20	-	-	-	-	-	-	-	-	-	-	4	-	-	-	-	-	3	-	
<i>Disphyma crassifolium</i>	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Lepidium rotundum</i>	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Cassinia arcuata</i>	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Rhagodia crassifolia</i>	-	60	-	-	-	-	-	-	-	-	7	50	-	-	-	-	-	-	-	-	
<i>Lepidosperma</i> sp. Mt Chester (S Kern et al. LCH 16596)	-	-	-	-	6	-	-	3	-	25	3	-	-	80	80	-	15	-	-	-	
<i>Austrostipa pycnostachya</i>	-	60	3	-	-	-	-	-	-	-	-	-	4	-	-	-	-	-	-	-	
<i>Dianella brevicaulis</i>	13	60	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Eucalyptus salmonophloia</i>	-	80	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Acacia chrysella</i>	-	20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Lepidosperma</i> sp. (GF Craig 8243)	-	20	3	40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	-	
Number of plots	8	5	34	5	12	5	4	12	5	2	15	4	23	26	6	4	33	17	7	32	4

APPENDIX 3

Significant indicator taxa of the 21 floristic community types recognised on the Ravensthorpe Range. Indicator values (%) are shown only for taxa which were significant at $p \leq 0.05$ (from Monte Carlo permutation test). Species with high indicator values (≥ 14) are indicated by shading. Community 10 was not included as the low number of plots (2) precluded this type of analysis.

Taxon	Woodlands							Mallee shrublands						Species-rich mallee shrublands						
	1	2	3	4	5	6	7	8	9	11	12	13	14	15	16	17	18	19	20	21
Acacia sp. Ravensthorpe Range	22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Austrodanthonia setacea</i>	17	7	-	-	-	-	-	-	-	-	-	7	-	-	3	-	-	-	-	3
<i>Austrostipa acrociliata</i>	21	6	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Austrostipa hemipogon</i>	22	20	-	-	-	-	-	-	-	-	-	5	-	-	-	-	-	-	-	-
<i>Austrostipa</i> sp. Ravensthorpe Range	43	4	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Daucus glochidiatus</i>	22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Enchytraea tomentosa</i> var. <i>tomentosa</i>	16	4	-	-	-	-	-	-	-	-	7	-	-	-	-	-	-	-	-	-
<i>Eucalyptus phenax</i> subsp. <i>phenax</i>	19	3	14	1	-	7	-	1	12	-	1	-	-	-	-	-	3	1	1	-
<i>Grevillea patentiloba</i> subsp. <i>platypoda</i>	22	-	-	-	-	-	-	-	-	-	2	-	2	6	-	-	-	-	-	-
<i>Oxalis perennans</i>	50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Ptilotus holosericeus</i>	63	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Sonchus oleraceus</i>	20	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-
<i>Vittadinia gracilis</i>	25	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Acacia erinacea</i>	10	41	1	-	1	7	-	-	-	-	3	-	-	-	-	-	-	-	-	-
<i>Acacia glaucoptera</i>	9	24	6	-	4	-	-	1	15	-	6	-	-	-	-	-	-	-	-	-
<i>Austrostipa elegantissima</i>	7	33	-	-	-	-	-	-	-	-	13	2	-	-	-	-	-	-	-	-
<i>Austrostipa pycnostachya</i>	-	54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Dianella brevicaulis</i>	2	50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus salmonophloia</i>	-	80	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Lepidosperma</i> sp. Mt Chester	-	58	-	-	-	-	6	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Olearia muelleri</i>	7	53	-	-	3	-	-	-	-	-	3	-	-	-	-	-	-	-	-	-
<i>Rhagodia crassifolia</i>	-	31	-	-	-	-	-	-	-	-	21	-	-	-	-	-	-	-	-	-
<i>Wilsonia humilis</i>	-	19	3	8	-	-	-	-	2	-	-	-	-	-	-	-	4	-	-	-
<i>Acacia binata</i>	2	-	18	-	1	-	-	-	4	-	-	-	-	-	-	-	-	2	-	-
<i>Eucalyptus calycogona</i> subsp. <i>calycogona</i>	14	-	16	9	-	-	-	-	2	-	-	-	-	-	-	-	-	-	-	-
<i>Lepidosperma fimbriatum</i>	1	2	15	-	-	-	-	-	2	1	4	-	-	-	-	-	1	-	-	-
<i>Melaleuca sophisma</i>	1	-	20	-	1	-	-	-	16	1	-	-	-	-	-	-	14	-	-	-
<i>Cassytha melantha</i>	-	4	11	17	-	-	-	-	1	-	-	-	-	3	-	-	14	1	-	-
<i>Daviesia benthamii</i>	-	6	1	24	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus tenera</i>	-	-	-	67	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus flocktoniae</i> subsp. <i>flocktoniae</i>	16	1	13	16	5	1	4	-	-	-	-	-	-	-	-	-	12	2	-	-
<i>Eucalyptus pileata</i>	-	2	2	29	11	2	-	-	-	-	3	-	-	-	-	-	-	-	-	-
<i>Grevillea pectinata</i>	-	8	1	18	3	-	-	-	-	-	-	-	-	-	-	-	17	-	-	-
<i>Lepidosperma</i> sp. (GF Craig 8243)	-	6	-	23	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-
<i>Lepidosperma</i> sp. Maydon	-	-	-	22	-	-	-	1	-	-	-	-	-	-	-	-	-	-	9	-
<i>Melaleuca acuminata</i> subsp. <i>acuminata</i>	-	-	-	18	-	-	-	-	-	1	7	2	-	-	-	-	-	-	-	-
<i>Melaleuca coronicarpa</i>	-	-	-	1	30	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Appendix 3 (cont.)

Taxon	Woodlands							Mallee shrublands						Species-rich mallee shrublands						
	1	2	3	4	5	6	7	8	9	11	12	13	14	15	16	17	18	19	20	21
<i>Boronia inornata</i> subsp. <i>inornata</i>	2	2	1	-	-	2	-	-	-	17	38	-	-	-	-	-	-	1	-	-
<i>Eucalyptus celastroides</i> subsp. <i>virella</i>	-	-	-	-	-	-	-	-	-	22	-	-	-	-	-	-	-	-	-	-
<i>Eucalyptus oleosa</i> subsp. <i>corvina</i>	4	-	-	-	2	-	-	-	-	-	65	-	-	-	-	-	-	-	-	-
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	21	6	1	-	-	1	-	-	-	-	21	2	-	-	-	-	-	-	-	-
<i>Phebalium tuberculosum</i>	2	-	-	-	-	-	-	-	-	-	-	48	1	-	-	-	-	-	-	-
<i>Trymalium elachophyllum</i>	4	-	-	-	-	-	-	-	-	-	16	-	-	-	-	-	-	-	-	-
<i>Acacia</i> sp. Ravensthorpe	-	-	-	2	-	-	-	-	-	-	4	18	-	-	7	1	-	2	-	-
<i>Banksia laevigata</i> subsp. <i>laevigata</i>	-	-	-	-	-	-	-	-	-	-	-	18	-	-	9	-	-	-	-	-
<i>Beaufortia orbifolia</i>	-	-	-	-	-	2	-	-	-	-	-	27	25	-	2	-	-	-	-	-
<i>Boronia ternata</i> var. <i>elongata</i>	-	-	-	-	-	-	-	-	-	-	-	18	-	-	2	-	-	-	-	-
<i>Brachyloma geissoloma</i> subsp. <i>geissoloma</i>	-	-	-	-	-	-	-	-	-	-	-	29	-	-	1	-	-	-	-	-
<i>Eucalyptus falcata</i> subsp. <i>falcata</i>	-	-	-	-	-	-	1	-	-	-	11	17	2	11	10	14	-	2	-	-
<i>Hakea multilineata</i>	-	-	-	-	-	-	-	-	-	-	1	17	-	-	-	-	-	-	-	-
<i>Philotheeca gardneri</i> subsp. <i>gardneri</i>	-	-	-	-	1	-	-	-	-	-	6	34	-	-	-	-	-	-	-	-
<i>Andersonia</i> aff. <i>lehmanniana</i>	-	-	-	-	-	-	-	-	-	-	-	-	31	-	-	-	-	-	-	-
<i>Banksia heliantha</i>	-	-	-	-	-	-	-	-	-	-	-	-	36	3	-	35	-	-	-	-
<i>Boronia subsessilis</i>	-	-	-	-	-	-	-	-	-	-	-	-	24	-	1	-	-	1	-	-
<i>Chorizema trigonum</i>	-	-	-	-	-	-	4	-	-	-	-	-	17	4	-	6	-	2	-	-
<i>Daviesia mollis</i>	-	-	-	-	-	-	-	-	-	-	-	-	23	-	-	1	-	1	-	-
<i>Eucalyptus lehmannii</i> subsp. <i>parallela</i>	-	-	-	-	-	-	-	-	-	-	-	-	1	32	-	-	2	-	-	-
<i>Eucalyptus preissiana</i> subsp. <i>preissiana</i>	-	-	-	-	-	-	-	-	-	-	-	-	34	-	-	8	-	-	-	-
<i>Grevillea concinna</i> subsp. <i>leemanniana</i>	-	-	-	-	-	-	-	-	-	-	-	-	33	-	-	-	-	-	-	-
<i>Grevillea nudiflora</i>	-	-	-	-	-	-	-	-	-	-	-	-	19	-	-	5	-	1	-	-
<i>Hakea obtusa</i>	-	-	-	-	-	-	-	-	-	-	1	14	33	2	7	1	1	1	-	-
<i>Lepidosperma</i> sp. Cordingup	-	-	-	-	-	2	-	-	-	-	3	7	18	3	1	-	1	-	3	-
<i>Leucopogon carinatus</i>	-	-	-	-	-	-	-	-	-	-	-	-	19	5	-	13	-	1	-	-
<i>Leucopogon infuscatus</i>	-	-	-	-	-	-	-	3	-	-	-	-	19	-	-	-	14	2	-	-
<i>Spyridium majoranifolium</i>	-	-	-	-	-	-	-	-	-	-	-	-	18	10	-	-	-	-	-	-
<i>Stylium albomontis</i>	-	-	-	-	-	1	-	1	-	-	3	12	19	2	4	-	1	5	-	-
<i>Hibbertia mucronata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	45	-	1	-	-	-	-
<i>Hovea acanthoclada</i>	-	-	-	-	1	1	-	-	-	9	1	-	16	35	-	-	-	-	-	-
<i>Labichea lanceolata</i> subsp. <i>brevifolia</i>	1	-	-	-	-	-	-	-	-	-	8	-	-	48	-	-	-	-	-	-
<i>Lasiopteratum rosmarinifolium</i>	-	-	-	-	-	-	-	-	-	-	-	1	1	30	-	5	-	10	-	-
<i>Lomandra mucronata</i>	-	-	-	-	-	-	-	-	-	-	-	-	7	26	13	20	-	5	2	-
<i>Melaleuca pentagona</i>	-	-	-	-	-	-	-	-	-	-	-	-	12	26	-	-	-	-	-	-
<i>Adenanthes flavidiflorus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	33	-	-	-	-	-
<i>Astrolobium prostratum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	27	-	-	-	-	-
<i>Banksia cirsoides</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	9	-	42	-	-	1	-
<i>Banksia corvijuga</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	21	-	-	-	-	-
<i>Banksia foliosissima</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	18	-	-	-	-	-
<i>Banksia pallida</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	61	-	-	-	-	-

<i>Beaufortia micrantha</i> var. <i>micrantha</i>	-	-	-	-	-	-	-	-	-	-	-	-	33	9	-	2	2	
<i>Beaufortia schaueri</i>	-	-	-	-	-	-	-	-	-	-	1	10	22	7	1	5	2	
<i>Boronia crassifolia</i>	-	-	-	-	-	-	-	-	-	-	6	-	28	16	-	1	-	
<i>Cassytha pomiformis</i>	-	-	-	-	-	-	-	-	-	-	1	-	26	-	-	1	-	
<i>Dampiera juncea</i>	-	-	-	-	-	-	-	-	-	-	-	-	60	-	-	-	-	
<i>Daviesia euryloba</i>	-	-	-	-	-	-	-	-	-	-	-	-	30	-	-	-	-	
<i>Drosera grievei</i>	-	-	-	-	-	-	-	-	-	-	-	-	33	-	-	-	-	
<i>Goodenia pinifolia</i>	-	-	-	-	-	-	-	-	-	-	10	-	25	-	-	-	5	
<i>Guichenotia anota</i>	-	-	-	-	-	-	-	-	-	-	-	-	27	-	-	-	-	
<i>Hakea cygna</i> subsp. <i>cygna</i>	-	-	-	-	-	-	-	-	-	-	-	-	30	-	-	-	-	
<i>Hakea subsulcata</i>	-	-	-	-	-	-	-	-	-	-	-	-	18	-	-	-	-	
<i>Hemigenia teretiuscula</i>	-	-	-	-	-	-	-	-	-	-	-	-	27	-	-	-	-	
<i>Hibbertia gracilipes</i>	-	-	-	-	-	-	-	-	-	-	6	1	-	20	16	1	5	7
<i>Isopogon</i> sp. Ravensthorpe (DB Foreman 1207)	-	-	-	-	-	-	-	-	-	-	15	8	-	16	-	-	-	-
<i>Jacksonia viscosa</i>	-	-	-	-	-	-	-	-	-	-	3	-	-	29	4	-	-	-
<i>Lepidobolus chaetocephalus</i>	-	-	-	-	-	-	-	-	-	-	-	-	18	-	-	-	-	
<i>Lepidosperma</i> sp. (RL Barrett 3570)	-	-	-	-	-	-	-	1	-	-	-	1	-	41	-	-	2	-
<i>Leptospermum</i> sp. Bandalup Hill (G Cockerton 110-1)	-	-	-	-	-	-	-	-	-	-	-	6	-	35	16	-	-	-
<i>Leucopogon cuneifolius</i>	-	-	-	3	-	-	-	-	-	-	9	2	-	21	-	-	-	-
<i>Melaleuca villosisepala</i>	-	-	-	-	-	-	-	-	-	-	-	-	54	-	-	2	5	
<i>Persoonia striata</i>	-	-	-	-	-	-	-	-	-	-	-	-	26	8	-	-	-	
<i>Petrophile glauca</i>	-	-	-	-	-	-	-	-	-	-	11	-	30	-	-	-	-	
<i>Petrophile seminuda</i>	-	-	-	-	-	-	-	-	-	-	-	-	15	12	-	-	6	
<i>Schoenus brevisetis</i>	-	-	-	-	-	-	-	-	-	-	-	-	36	-	-	-	-	
<i>Stachystemon brachyphyllus</i>	-	-	-	-	-	-	-	-	-	-	-	-	24	-	-	-	-	
<i>Adenanthera oreophilus</i>	-	-	-	-	-	-	-	-	-	-	1	5	-	-	21	-	-	-
<i>Allocasuarina humilis</i>	-	-	-	-	-	-	-	-	-	-	-	1	3	26	33	-	-	-
<i>Amphipogon turbinatus</i>	-	-	-	-	-	-	-	-	-	-	-	1	2	14	22	-	4	10
<i>Argentipallium niveum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	19	-	1	-
<i>Banksia lemanniana</i>	-	-	-	-	-	-	-	-	-	-	1	10	2	13	19	-	2	-
<i>Calothamnus gracilis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	27	-	-	-
<i>Calothamnus pinifolius</i>	-	-	-	-	-	-	-	-	-	-	-	-	10	-	21	-	-	-
<i>Chorizema uncinatum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	-	-	-
<i>Conostylis bealiana</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	21	-	4	-
<i>Dampiera fasciculata</i>	-	-	-	-	-	-	-	-	-	-	-	6	-	-	19	-	-	-
<i>Eucalyptus pleurocarpa</i>	-	-	-	-	-	-	-	-	-	-	14	9	11	17	19	2	5	-
<i>Gompholobium viscidulum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	24	-	-	-
<i>Hakea pandanicarpa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	9	19	-	1	-
<i>Isopogon</i> sp. Fitzgerald River	-	-	-	-	-	-	-	-	-	-	-	-	-	-	19	-	1	-
<i>Isopogon trilobus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	27	-	-	-
<i>Jacksonia elongata</i>	-	-	-	-	-	-	-	-	-	-	-	5	-	-	28	-	-	-
<i>Lepidosperma</i> sp. Mt Burdett	-	-	-	-	-	-	-	1	-	-	-	2	-	5	19	1	2	-
<i>Leucopogon conostephoides</i>	-	-	-	-	-	-	-	-	-	-	1	-	3	6	26	-	9	3
<i>Lysinema ciliatum</i>	-	-	-	-	-	-	-	-	-	-	-	-	27	31	-	1	-	-
<i>Mesomelaena stygia</i> subsp. <i>stygia</i>	-	-	-	-	-	-	-	-	-	-	-	-	20	33	-	-	12	-
<i>Petrophile squamata</i> subsp. <i>northern</i>	-	-	-	-	-	-	-	-	-	-	-	-	1	38	-	2	-	-
<i>Pultenaea indira</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	30	-	1	-

Taxon	Woodlands							Mallee shrublands						Species-rich mallee shrublands							
	1	2	3	4	5	6	7	8	9	11	12	13	14	15	16	17	18	19	20	21	
<i>Schoenus sublaxus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	37	-	-	-	
<i>Stachystemon virgatus</i>	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	23	-	-	8	
<i>Taxandria spathulata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	17	2	8	31	-	2	-	
<i>Xanthorrhoea platyphylla</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	53	-	-	-	-	
<i>Acacia ingrata</i>	-	-	4	-	-	-	-	1	-	19	7	-	-	-	-	-	-	30	6	-	
<i>Acrotriche cordata</i>	-	-	1	-	-	-	-	1	6	1	-	1	-	9	-	1	-	18	2	-	
<i>Banksia media</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	29	1	-	
<i>Boronia inornata</i> subsp. <i>leptophylla</i>	-	-	4	7	3	-	-	1	2	-	-	-	-	-	-	-	-	32	-	-	
<i>Comesperma spinosum</i>	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	37	-	-	
<i>Coopernookia polygalacea</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	41	4	-	
<i>Daviesia anceps</i>	-	-	1	-	-	-	-	7	10	3	-	-	-	1	-	-	-	21	9	2	
<i>Daviesia articulata</i>	-	2	3	-	-	2	4	-	-	-	-	-	-	-	-	-	-	20	2	-	
<i>Eucalyptus kesselii</i> subsp. <i>eugnosta</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	38	-	-	
<i>Eucalyptus leptocalyx</i>	-	-	-	-	-	-	3	-	19	-	-	-	-	-	-	-	-	1	26	1	-
<i>Eucalyptus suggrandis</i> subsp. <i>suggrandis</i>	-	2	2	2	-	-	3	-	-	-	-	-	-	-	1	-	-	-	22	7	-
<i>Hibbertia exasperata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	39	-	-	
<i>Melaleuca subfalcata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	20	4	-	
<i>Pomaderris brevifolia</i>	-	-	1	-	-	-	-	-	7	10	-	1	1	-	-	-	-	24	-	-	
<i>Spyridium cordatum</i>	-	-	-	13	-	-	-	-	-	-	-	-	-	1	1	-	3	-	26	8	-
<i>Styphelia intertexta</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	40	-	-	
<i>Eucalyptus incrassata</i>	-	-	-	-	-	-	2	-	-	-	1	1	1	4	2	5	16	1	18	-	
<i>Eucalyptus uncinata</i>	-	-	-	-	-	-	-	-	4	-	-	-	2	10	-	5	2	12	16	6	
<i>Eutaxia cuneata</i>	-	-	-	5	-	-	-	-	-	-	-	-	2	-	-	-	-	-	25	-	
<i>Gahnia ancistrophylla</i>	-	3	-	3	1	-	-	1	1	-	-	-	1	1	-	4	2	7	14	12	
<i>Hakea laurina</i>	-	-	-	1	-	-	2	-	1	-	-	2	1	-	2	-	3	11	15	-	
<i>Harperia lateriflora</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	19	-	
<i>Melaleuca rigidifolia</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	6	-	2	6	1	16	3	
<i>Tetralonia capillaris</i>	-	-	-	-	-	-	-	1	-	2	-	-	-	7	7	4	1	9	17	-	
<i>Acacia mimica</i> var. <i>angusta</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	72	-	
<i>Allocasuarina campestris</i>	-	-	-	-	-	-	-	6	-	-	-	1	1	-	-	-	-	-	-	25	
<i>Astroloma serratifolium</i>	-	-	-	-	-	-	-	8	-	-	-	-	-	-	4	2	-	-	1	17	
<i>Baeckea crispiflora</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	50	-	
<i>Calytrix leschenaultii</i>	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	2	2	-	5	38	
<i>Chorizema aciculare</i> subsp. <i>aciculare</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	2	29	-	
<i>Conostylis argentea</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	1	3	14	-	-	4	42	
<i>Daviesia pachyphylla</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	37	-	
<i>Eucalyptus desmondensis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	75	-	
<i>Gompholobium marginatum</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	60	
<i>Grevillea dolichopoda</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	1	70	
<i>Haemodorum discolor</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	18	
<i>Hakea lissocarpa</i>	-	-	-	-	-	-	-	-	-	-	-	-	1	-	3	1	2	-	19	24	

<i>Hibbertia pungens</i>	-	-	-	3	-	-	-	11	-	-	-	4	1	9	-	1	-	4	9	20
<i>Lepidobolus preissianus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	1	35
<i>Leucopogon concinnus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9	1	-	-	-	46
<i>Logania tortuosa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	22
<i>Melaleuca carrii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	42
<i>Melaleuca hamata</i>	1	2	1	2	1	-	3	6	-	-	-	6	6	3	-	-	-	10	12	13
<i>Schoenus racemosus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	4	1	10	18	18
<i>Schoenus sesquispiculus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	62
Number of plots	8	5	34	5	12	5	4	12	5	4	15	23	26	6	4	33	17	7	32	4