




TESTING A PROTOCOL FOR  
FORESHORE ASSESSMENT  
IN METROPOLITAN WATERWAYS



WATER RESOURCE MANAGEMENT SERIES

WATER AND RIVERS COMMISSION REPORT NO. WRM 13  
1999



WATER AND RIVERS  
COMMISSION

# Testing a protocol for foreshore assessment in Metropolitan Waterways

Prepared by  
Kelly Shepherd and Nicole Siemon  
Ecosystem Management Services

Jointly funded by



Natural Heritage Trust



WATER AND RIVERS  
COMMISSION

WATER AND RIVERS COMMISSION  
REPORT NO. WRM 13  
OCTOBER 1999



# Acknowledgments

This report has been jointly funded by the Natural Heritage Trust and the Water and Rivers Commission.

This report was prepared by Nicole Siemon, Senior Environmental Consultant and Kelly Shepherd, Consultant, Ecosystem Management Services (EMS).

Methodology development by Nicole Siemon, EMS in consultation with Dr Luke Pen and Jodie Oates, Water and Rivers Commission.

Surveys and mapping were undertaken by Kelly Shepherd, EMS.

Many members of the community and catchment group representatives provided valuable input and feedback in the development of the methodology and surveys.

Digital map production by Brett Harrison, Banksia Environmental Mapping.

# Reference Details

The recommended reference for this publication is:  
Water and Rivers Commission 1999, *Testing a Protocol for Foreshore Assessment in Metropolitan Waterways*.  
Water and Rivers Commission, Water Resource Management Series, No WRM 13.

ISBN 0-7309-7338-7

ISSN 1326-6934

*Text printed on 100gsm recycled stock,  
Cover; 250gsm Sapphire Dull  
October 1999*



# Foreword

Landcare groups in Western Australia have been concerned with the protection and rehabilitation of river systems for some time. However, with such large areas to cover, and many streams being in private ownership, there is a lack of information available to many groups to assist them in making management decisions.

In 1995 Pen and Scott developed a technique for '*Stream Foreshore Assessment in Farming Areas*'.

This provided a standardised assessment technique that can be performed by groups and individual landholders themselves. It has been widely accepted and used to successfully assess many streams throughout south-west WA. As use of the technique has expanded from farm to catchment scale surveys, some users began to express a need for a modification of the methodology that would enable them to assess streams in urban and semi-rural environments, where there are a different suite of issues to be considered. In 1997 the Water and Rivers Commission obtained Natural Heritage Trust funding to assist in the development of a foreshore condition assessment methodology suitable for use in urban areas and to undertake surveys on several major tributaries of the Swan-Canning Catchment.

Nicole Siemon and Kelly Shepherd of Ecosystem Management Services (EMS), in consultation with the Water and Rivers Commission, have developed a technique for '*Foreshore Condition Assessment in Urban and Semi-rural Areas*'. The assessment technique is comprehensive yet, like that of Pen and Scott, does not require specialised knowledge or expensive technical assistance and hence assessment can be performed by groups and individuals themselves.

The methodology considers overall stream condition to be comprised of four major parameters that are independently assessed and the results are then combined to determine the overall stream condition.

**Bank stability** includes assessment of bank slope, erosion, slumping, sedimentation and stabilising structures.

**Foreshore vegetation** structure and composition, includes the use of tables with native and weed species commonly found in the region. This allows for straightforward yet comprehensive vegetation surveys

looking at abundance, health and regeneration of individual species.

**Stream cover** recognises the importance of overhanging native vegetation and in-stream cover, and notes the abundance of native and exotic vegetation and the presence of deciduous trees.

**Habitat diversity** includes stream form, water quality and identifies habitat requirements for a variety of terrestrial and aquatic fauna.

Along with recording information on stream condition at the time of the survey the methodology also ensures that information is collected that will aid groups in making management decisions. This information includes disturbance factors, surrounding land use, evidence of existing management and special cultural or spiritual significance.

The condition assessment technique that has been developed has several features that are particularly important in helping groups to make their own river management decisions. The techniques:

- do not require specialised knowledge or expensive technical assistance and surveys can therefore be undertaken by individual landholders or by community groups;
- immediately provide managers with data to aid them in their decision making, especially in prioritisation of works;
- provide standardised data suitable for compilation and comparative assessment, even when using data collected by a variety of groups and individuals; and
- provide standardised data suitable for ongoing monitoring and evaluation.

The methodology has been tested on several tributaries in the Swan-Canning catchment. These tributaries have active catchment groups working on, or planning, rehabilitation works. Reaches surveyed were those identified by the catchment groups as priority areas in which they plan to be undertaking works. It is hoped that this report will assist in the long-term management of these tributaries.





# Contents

1. Introduction .....	1
1.1 Need for this study .....	1
1.2 Community involvement process.....	1
1.3 This report .....	2
2. Methodology.....	3
2.1 Site selection within tributaries.....	3
2.2 Implementing the assessment method.....	3
2.2.1 Undertaking foreshore surveys.....	3
2.2.2 Environmental parameters of foreshore condition .....	4
2.2.3 Collating the results .....	17
3. Key findings for all survey sites .....	8
3.1 Bank stability .....	8
3.2 Vegetation.....	8
3.2.1 Native species .....	8
3.2.2 Weeds .....	9
3.3 Stream cover.....	9
3.4 Habitat diversity .....	10
3.5 Overall summary conditions for all surveyed sites.....	11
3.5.1 Summary results for Bennett Brook (Bennett Brook Catchment) .....	11
3.5.2 Summary results for Bannister Creek (Canning Catchment).....	13
3.5.3 Summary results for Canning River (Canning Catchment) .....	14
3.5.4 Summary results for Roley Pool (Canning Catchment).....	16
3.5.5 Summary results for Southernwood Creek (Canning Catchment).....	16
3.5.6 Summary results for Wright Brook (Canning Catchment).....	17
3.5.7 Summary results for Breera Brook (Ellen Brook Catchment).....	18
3.5.8 Summary results for Ellen Brook (Ellen Brook Catchment) .....	19
4. Specific site reports .....	21
5. General recommendations.....	22
6. Common issues .....	24
6.1 Ownership and access .....	24
6.2 Developing management and rehabilitation plans .....	24
6.3 Long term management.....	25
6.4 Surrounding landuse.....	25
6.5 Gaining support from state and local government.....	26
6.7 Fire management .....	26
6.8 Access to information.....	26
7. Summary .....	27
8. References and recommended reading.....	28



## Appendices

<b>Appendix 1:</b> Native and weed species identified during the foreshore assessment process (1998).....	29
<b>Appendix 2:</b> Suggested weed control methods.....	37
<b>Appendix 3:</b> Suggested species for revegetation works .....	63
<b>Appendix 4:</b> Suggested soft engineering works .....	69
<b>Appendix 5:</b> Condition mapping symbols .....	71

## List of tables

<b>Table 1:</b> Colour codes and points value for ranking stream conditions.....	4
<b>Table 2:</b> The summary conditions of the environmental parameters assessed to determine foreshore health.....	5
<b>Table 3:</b> Summary of Stream Condition Index .....	7



# 1. Introduction

The riparian zone adjacent to natural watercourses acts as a buffer to the surrounds. Healthy foreshore vegetation stabilises the foreshore banks, slows and filters water thus reducing erosion of the banks and sedimentation of major channels. Foreshore vegetation also provides stream cover and suitable habitats for aquatic and terrestrial animals. Often these areas are a haven for native fauna, particularly during the dry summer months.

Riparian areas have always been a focus for development and as a consequence are often highly degraded. The major threats to foreshore health are the loss of native vegetation or a decline in health due to weed invasion. The loss of deep-rooted native plants often causes the destabilisation of foreshore banks, leaving these areas prone to erosion particularly during peak flow events.

Gaining an understanding of the health of river foreshores is the first step towards developing appropriate management strategies to protect and enhance these areas:

## 1.1 Need for this study

Community groups are becoming increasingly interested in foreshore management and are taking an active role in this process. This interest in foreshores provides opportunities to collect substantial data about waterways.

The need for a standard methodology to assess foreshore condition was recognised to ensure consistency of information gathering, in the early 1990s. Procedures for recording information on foreshore condition have been available in rural areas for a number of years (Pen and Scott 1995), however, this system had limited applications in urban and semi-rural environments. Recognition of the need to modify this methodology

occurred in 1997, and resulted in a funding application being developed for the Natural Heritage Trust. This successful application required the development of a standard foreshore assessment method based on the rural system (Water and Rivers Commission 1999), testing of the new methodology and developing a reporting technique for this work. Ecosystem Management Services (EMS) undertook this project on behalf of Water and Rivers Commission (WRC) and the Natural Heritage Trust (NHT).

## 1.2 Community involvement process

The intended audience for the foreshore assessment method is state and local government officers and the community. In order to ensure that the information included on the assessment form was relevant to these groups, and captured most of the data required, EMS and the WRC implemented a community involvement process for development of the form.

A preliminary draft of the foreshore assessment method was developed and presented to representatives from many of the catchment groups in the metropolitan area. The comments from this meeting were assimilated into the method. This second draft was then presented at a subsequent series of meetings with each catchment group, to canvass further comments. Again, suggestions recorded were collated and incorporated into the document.

Discussion was also held at the second series of meetings to determine specific areas of interest for each catchment group. Each group identified priority foreshore areas to undergo assessment, to enable further refinement of the standard methodology. The locations selected included areas that were already a focus or are potential sites for future rehabilitation works.





The sites nominated by groups to be surveyed were as follows:

**Bennett Brook Catchment**

- Bennett Brook

**Upper Canning Catchment**

- Bannister Creek
- Canning River
- Roley Pool
- Southernwood Creek
- Wright Brook

**Ellen Brook Catchment**

- Breera Brook
- Ellen Brook

As a result of time constraints and access difficulties not all of the foreshore areas that were nominated by the community groups were surveyed.

## **1.3 This report**

This report summarises the results of the preliminary surveys using the revised (draft) foreshore assessment method (Water and Rivers Commission 1999). These surveys were conducted to verify and refine the methodology. Recommended strategies for appropriate management of future works on the focus foreshore areas are also detailed in the document. Information is provided on weed control techniques, recommended native species for rehabilitation work and methods to undertake soft engineering works.



## 2. Methodology

### 2.1 Site selection within tributaries

Following the community involvement process the nominated sections of the selected waterways were assessed to determine the most appropriate areas to undertake the foreshore survey. This was based on the need to assess a complete range of foreshore health to ensure that the assessment method was sufficiently balanced to cover all situations ranging from rural to urban zones.

Following is a summary of the extent of the nominated waterways that were surveyed to assess and refine the foreshore assessment methodology.

#### Bennett Brook Catchment Group

Waterway	Extent of Survey
Bennett Brook	North of Benara Road upstream to Mussel Pool, Whiteman Park.

#### Bannister Creek Catchment Group

Waterway	Extent of Survey
Bannister Creek	South of Adenia Road to Iveston Road

#### Upper Canning/Southern-Wungong Integrated Catchment Group

Waterway	Extent of Survey
Roley Pool	Soldiers Road to a few hundred metres upstream of the stairway at Collins Road.
Southernwood Creek	Confluence of Southern River to the entry drain near Michel Crescent.
Canning River	South of Parkside Drive to a few hundred metres upstream of Albany Highway.
Wright Brook	Confluence of Canning River with Wright Brook to a few hundred metres upstream of Turner Road.

#### Ellen Brook Integrated Catchment Group

Waterway	Extent of Survey
Breera Brook	~ 80 m upstream of railway line to eastern boundary of Lot M13.
Ellen Brook	Railway line at Almeria Parade upstream to the end of Lot 501.

### 2.2 Implementing the assessment method

The foreshore assessment survey method has been developed to enable community groups to assess the condition of foreshores in urban and semi-rural areas. For detailed information on the methodology used to assess foreshore condition refer to Water and Rivers Commission (1999).

As outlined above, this process ensures consistency of information gathering allowing the information collected from multiple surveys, conducted by various people over time, to be collated. The accumulated information can then be used to prepare management plans and focus on priority areas for rehabilitation. The results can also be used to monitor changes over time and to compare different foreshore areas, and be shared amongst state and local government authorities and the community.

#### 2.2.1 Undertaking foreshore surveys

Each of the foreshore areas selected were traversed prior to the survey being conducted. The foreshore was then divided into relatively homogeneous sections of similar vegetation structure and land use. A survey was conducted for each of these sections, and the condition of the foreshore parameters was calculated and the overall Stream Condition Index was determined.

In areas where foreshore vegetation was very dense on both banks, both sides were surveyed separately and a form was completed for each side. On highly degraded rivers where the foreshore along both banks was easily observed from one side, and the vegetation and disturbance factors were similar, a single survey form was completed.



Scaled baseline maps were prepared by WRC showing cadastral boundaries and the waterway. The cadastral information assists in gaining bearings out in the field. As each homogeneous section was identified, information was sketched onto the baseline maps. Other information such as the extent of vegetative overstorey along the foreshore, the location and extent of predominant middlestorey native species and weeds and the presence of disturbance factors such as discharge pipes and infrastructure such as fences present, were detailed on each map. This ensured that each form completed for a specific section also had all relevant information marked on the correct map.

Note that the left and right side of the main channel are defined by looking upstream.

### 2.2.2 Environmental Parameters of Foreshore Condition

Principal environmental parameters are used as indicators of foreshore condition and are assessed during the foreshore survey to determine the overall Stream Condition Index.

These parameters are;

- Bank stability
- Foreshore vegetation
- Stream cover
- Habitat diversity

A colour coded system has been developed to summarise the condition of each of the above environmental parameters. This system allows the information to be provided in an immediately recognizable form. The status of each of the parameters are assessed and graded from Blue (Excellent) to Black (Very Poor) (Table 1) using the criteria outlined in Table 2. For example, the Bank Stability of an area is determined by assessing the level of erosion, slumping and sedimentation along the foreshore. In a pristine area where there is no discernable decline in condition, and no obvious erosion the Bank Stability may be graded as Blue. In a highly modified system where the foreshore is highly degraded and subject to severe erosion and bank collapse, Bank Stability may be graded as Red or Black. A scoring system is linked to this process to provide a quantitative method of calculating stream health.

**Table 1: Colour codes and points value for ranking stream conditions**

<b>Condition</b>	Excellent	Good	Moderate	Poor	Very Poor
<b>Colour rating</b>	<b>Blue</b>	<b>Green</b>	<b>Yellow</b>	<b>Red</b>	<b>Black</b>
<b>Score</b>	8	6	4	2	0

From: Water and Rivers Commission (1999).



Table 2: Determining summary foreshore health

	<b>Blue - Excellent 8 points</b>	<b>Green - Good 6 points</b>	<b>Yellow - Moderate 4 points</b>	<b>Red - Poor 2 points</b>	<b>Black - Very poor 0 points</b>
<b>Bank Stability</b>	No erosion, slumping or sediment deposits; dense native vegetation cover on banks and verge; no evidence of disturbance or areas of exposed soil.	No significant erosion, slumping or sediment deposits in floodway or on lower banks; good native vegetation cover; only isolated areas of exposed soil or thinning vegetation.	Some localised erosion, slumping and sediment deposits; native vegetation cover on verges may be patchy and interspersed with patches of exposed soil.	Extensive active erosion slumping and sediment desposition particularly during peak flows; bare banks and verges common.	Almost continuous erosion; over 50% of banks slumping; sediment heaps line or fill much of the floodway; little or no vegetation cover.
<b>Foreshore vegetation</b>	Healthy, undisturbed native vegetation with structure intact and verges more than 20 m wide; no weed or signs of disturbance evident.	Vegetation structure dominated by native plants that comprise 80 - 100% of the total number of species; only scattered weeds or rarely evident in small clusters; nil or minor signs of disturbance (i.e. tracks, rubbish dumping).	Some changes in vegetation structure, native plants comprising of 50 - 80% of the total species composition; little regeneration of trees and shrubs; weeds occurring occasionally; moderate levels of disturbance.	Modified vegetation structure with native plants comprising only 20 - 50% of the total species composition. Trees remain with only scattered shrubs and an understorey dominated by weeds; high prevalence of disturbance.	Insufficient vegetation to control erosion; natural vegetation structure absent with occasional native trees and shrubs comprising less than 20% of the total species composition; weeds abundant; very high prevalence of disturbance and extensive areas of exposed soil.

	<b>Blue - Excellent 8 points</b>	<b>Green - Good 6 points</b>	<b>Yellow - Moderate 4 points</b>	<b>Red - Poor 2 points</b>	<b>Black - Very poor 0 points</b>
<b>Stream Cover</b>	Abundant stream cover from dense overhanging vegetation providing almost continuous shade; frequent instream cover from aquatic vegetation and/or leaf litter, rocks or logs.	Abundant shade from overhanging vegetation; occasional instream cover from patches of aquatic vegetation and isolated heaps of leaf litter or rocks and logs.	Scattered fringing vegetation with occasional patches of shade; infrequent instream cover with little aquatic vegetation, very infrequent rocks and logs.	Stream channel mainly clear; fringing vegetation almost absent providing very little permanent shade; instream cover almost absent with generally no instream vegetation and very infrequent rocks and logs.	Zero or minimal stream cover with no permanently shaded areas and no instream cover.
<b>Habitat Diversity</b>	Excellent water quality with permanent water (i.e: pools and creeks); three or more aquatic and terrestrial habitats including diverse vegetation types, edge waters, instream cascades, riffles, pools and woody debris.	Good water quality and some permanent water; at least three aquatic habitat types; at least one habitat type for terrestrial invertebrates; at least one habitat type for each terrestrial vertebrate category (frogs, reptiles and birds).	No apparent problems with water quality (i.e: muddy or cloudy in winter); at least two aquatic habitat types; at least one habitat type for terrestrial invertebrates; at least one habitat type for any two of the terrestrial vertebrate categories.	Possible seasonal problems with water quality and no permanent water; at least one aquatic habitat type; at least one habitat type for terrestrial invertebrates; at least one habitat type for one of the terrestrial vertebrates.	Poor water quality; almost no healthy habitats available for aquatic and terrestrial organisms.

The Stream Condition Index is a summary of the foreshore environmental parameters (Table 3) and is an indication of the overall stream condition.

**Table 3: Summary of Stream Condition Index**

Colour Code	Parameter Rating	Description
<b>Blue (32 points)</b>	<b>Excellent</b>	All parameters blue.
<b>Green (22-30 points)</b>	<b>Good</b>	Three to four parameters rated green or better with only one parameter rated yellow; no red or black ratings.
<b>Yellow (14-20 points)</b>	<b>Moderate</b>	Three parameters rated yellow or better with no more than one red; no black
<b>Red (6-12 points)</b>	<b>Poor</b>	Two or three parameters rated red with no more than one black.
<b>Black (0-4 points)</b>	<b>Very Poor</b>	Two or more parameters rated black.

### 2.2.3 Collating the results

The results compiled from the foreshore surveys of the selected sites were collated and a series of maps produced. These maps were digitised to enable presentation of the foreshore information in a visual format with corresponding text.

The summary codes of the condition of each environmental parameter and the Stream Condition Index are included on the summary map for each site.

This report contains a detailed description of the key findings of the four environmental parameters assessed for each survey section within the nominated survey sites. The recommended strategies for appropriate remedial works are discussed for each section.



## 3. Key findings for all sites

### 3.1 Bank stability

Bank stability is determined by the extent of erosion and slumping occurring along foreshore banks and the level of sedimentation within stream channels. Erosion is evident at almost all sites, generally at low to moderate levels.

There are however, a number of urban foreshores that are prone to severe erosion due to a lack of foreshore vegetation. For example severe erosion is occurring along much of Wright Brook and is particularly evident at Section C/Map 2 where the surrounding vegetation is reduced to a few trees and an understorey of maintained grass (lawn).

Localised disturbance frequently occurs along steep banks near the entry points of drainage channels or near outflow points of discharge pipes. Erosion also increases where infrastructure works have been undertaken for example near crossovers and bridges.

The impact of a decrease in the extent of dense emergent species along most of the foreshores surveyed is evidenced by increased erosion, particularly near the base of trees that grow immediately along the banks. As the soil is scoured away, roots are exposed and trees are less supported. Subsequently, there is an increased likelihood of trees collapsing and exacerbating the erosion problem.

Currently there is little evidence of severe bank collapse along the surveyed foreshore areas. Artificial stabilization structures have been utilised in some areas for example along Bannister Creek (Section A/Map 1). Recommended strategies include the use of geotextile matting to support areas cleared of weeds prior to planting native species to minimise the effect of further destabilizing foreshore banks (Appendix 4).

Sedimentation levels vary along the main channels of the surveyed watercourses. Sites such as the lower reaches of Ellen Brook (Map 1 and 2) show high levels of sedimentation. Large sandbars have become stabilised and vegetated along sections of Ellen Brook and Bennett Brook. Significant levels of sedimentation and increasing particle load in the water column are

indicative of erosion occurring further upstream. This highlights the need to understand processes occurring upstream of any waterway and shows that no site can be considered in isolation.

### 3.2 Vegetation

The foreshore vegetation along the majority of the surveyed tributaries is highly modified with remnant overstorey and typically weed dominated middlestorey and understorey. In some areas, often where residential housing extends almost to the edge of the watercourse, or alternatively in semi-rural areas where stock have access to the foreshore, the overstorey is completely absent or present for a few metres only on either side of the main channel. Breera Brook is the only example of a relatively healthy system within the survey sections with an overall Stream Condition of Yellow (Moderate) or Green (Good).

#### 3.2.1 Native species

Swamp paperbark (*Melaleuca raphiophylla*) and Flooded gum (*Eucalyptus rudis*) are the predominant overstorey trees along all of the foreshore sites. Other less common tree species include Modong (*Melaleuca preissiana*) and Marri (*Corymbia calophylla*) which often occur on drier soils.

Native shrubs are generally infrequent in the middlestorey and usually present in areas where there is an extensive and relatively healthy overstorey. The native shrubs often persist in low numbers away from the immediate foreshore as they are typically excluded by dense stands of weeds such as Watsonia (*Watsonia bulbifera*) which dominate these areas.

Species present at a number of the survey sites include Swamp peppermint (*Agonis linearifolia*), Coojong (*Acacia saligna*), Narrow leaved oxlybium (*Oxlybium lineare*), Prickly moses (*Acacia pulchella*), Swishbush (*Viminaria juncea*) and Blackboy (*Xanthorrhoea preissii*) (Appendix 1A).

There are very few native understorey species persisting along foreshore areas. Ground creepers such as Running postman (*Kennedia prostrata*) and Native wisteria



(*Hardenbergia comptoniana*) are present infrequently. Native sedges and rushes such as the Pale rush (*Juncus pallidus*) occur in clumps along the foreshore channels and in low-lying damp areas. The small herb Centella (*Centella cordifolia*) persists often in highly weed-infested areas.

There are a few native species that are abundant and form dense stands in the middlestorey and understorey. The native Bracken fern (*Pteridium esculentum*) for example, is often associated with an increased frequency of Marri trees in the overstorey and is located away from the immediate foreshore as seen at Bennett Brook (Section E/Map 2) and Breera Brook (Section D/Map 5). The Pithy sword sedge (*Lepidosperma longitudinale*) also forms dense stands running parallel to the waterway. This species predominates along most of Breera Brook.

### 3.2.2 Weeds

Exotic deciduous trees such as Fig (*Ficus* spp.), Willow (*Salix* spp.) and the Coral Tree (*Erythrina x sykesii*) are common along degraded foreshores in urban areas. These trees were originally planted as ornamentals or have escaped from nearby gardens. Deciduous trees threaten foreshore health as sudden leaf fall during winter decreases available stream cover and often introduces large amounts of vegetative material into the water column. The breakdown of large amounts of soft leaves may cause a sudden decline in the amount of available oxygen in the water column affecting instream organisms.

The control and removal of exotic trees is often difficult as species such as Willow produce numerous suckers. These trees often grow in areas with limited foreshore cover and the removal of these large trees may threaten bank stability.

Weeds in the middlestorey often form dense stands in clumps or in narrow strips along the edge of the watercourse. The most widespread weed species that form dense stands within the sections surveyed include Blackberry (*Rubus fruticosus*), Watsonia (*Watsonia bulbifera*) and the Giant reed (*Arundo donax*). Arum lily (*Zantedeschia aethiopica*) is frequently present in high numbers along foreshore areas and in low lying winter wet depressions in the floodplain. Other species that are present at a number of the survey sites but often

in low numbers include Castor oil (*Ricinus communis*), Deadly nightshade (*Solanum nigrum*) and Cotton bush (*Gomphocarpus fruticosus*) (Appendix 1B).

The greatest threat to revegetation is the presence of dominant understorey weeds including grasses such as Kikuyu (*Pennisetum clandestinum*), Couch (*Cynodon dactylon*), Perennial veldt grass (*Ehrharta calycina*), African lovegrass (*Eragrostis curvula*) and Wild oats (*Avena fatua*). Frequent annual weeds include Soursob (*Oxalis pes-caprae*), Whiteflower fumitory (*Fumaria capreolata*) and Fleabane (*Conyza* spp.). The introduced rush *Juncus microcephalus* occurs along foreshore banks and within low lying flooded areas. Creepers such as Bridal creeper (*Myrsiphyllum asparagoides*) and less frequently Morning glory (*Ipomoea* spp.) are present in a number of weed dominated foreshore areas.

The aquatic weed Watercress (*Rorippa nasturtium-aquaticum*) is present in the main channel at a number of the survey sites.

### 3.3 Stream cover

The level of overhanging vegetation and the abundance of native and exotic species along the foreshore determines the level of cover and permanent shade along a waterway. Instream emergent and submerged vegetation and the presence of rocks and logs also provide cover for aquatic organisms.

The condition of Stream Cover varied from Green (Good) to Black (Very Poor) along the selected survey sites. In relatively healthy, undisturbed areas along sections of Breera Brook the Stream Cover is graded as Yellow (Moderate) to Green (Good) due to the presence of a healthy overstorey and dense stream side vegetation including stands of emergent sedges and rushes. Stream Cover along sections of Bannister Creek (Section B/Map 1) and Ellen Brook (Section C/Map 4 and 5) were also graded as Yellow (Moderate) due to the presence of dense stands of weeds such as Watsonia (*Watsonia bulbifera*) or Blackberry (*Rubus fruticosus*) that overhang the waterway and provide patches of permanent shade.

Sections along Wright Brook (Section C/Map 2 and Section I/Map 8) and the Canning River (Section F/Maps 8 and 9 or Section G/Map 10) for example, were graded as Red (Poor) or Black (Very Poor). These areas are highly modified and the absence of dense overstorey





trees and the predominance of small weeds in the understorey minimise the amount of available stream cover at these sites.

The presence of large numbers of exotic trees in the overstorey also minimises available cover as a number of the most common species such as Fig, Willows and Coral trees are deciduous, dropping their leaves in autumn.

### **3.4 Habitat diversity**

Instream habitat diversity is affected by the quality and permanency of water and by the presence of instream rocks, submerged and emergent vegetation and logs. These features provide substrates for attachment for aquatic invertebrates, cover for fish and potential basking sites for turtles. Healthy, diverse streamside vegetation provides suitable habitats for terrestrial organisms and overstorey trees provide roosting and nesting sites for birds.

Many of the survey sites assessed were scored as having either Red (Poor) or Black (Very Poor) Habitat Diversity. Water flow in sites such as Wright Brook (Section C/Map 2) is seasonal and therefore unable to support aquatic organisms throughout the year. Frequently streams are narrow and shallow and are generally not suitable for fish and turtles.

The frequent lack of healthy, diverse native vegetation often limits the number of suitable habitats available for terrestrial animals.

The Habitat Diversity was graded as Yellow (Moderate) for Bennett Brook (Sections E - G/Maps 2 - 5) and Breera Brook indicating that there are no apparent problems with water quality and there are suitable sites for aquatic organisms such as logs and rocks instream. Further, diverse habitats for terrestrial organisms such as a variety of vegetation types, deep leaf litter and dense streamside vegetation are also present at these sites.



### 3.5 Overall summary conditions for all surveyed sites

The overall summary conditions of the foreshore sections surveyed for each of the tributaries is provided below. Most sites surveyed were diagnosed as poor Red (Poor) or Black (Very Poor) with only Breera Brook having Green (Good) foreshore condition.

#### 3.5.1 Summary results for Bennett Brook (Bennett Brook Catchment)

##### Bennett Brook (Section A)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Black	Black	Black
Moderate	Very Poor	Very Poor	Very Poor
4	0	0	0

Stream Condition
Black
Very Poor
4

##### Bennett Brook (Section B)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Black	Red	Red
Moderate	Very Poor	Poor	Poor
4	0	2	2

Stream Condition
Red
Poor
8

##### Bennett Brook (Section C)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Red	Yellow	Red
Poor	Poor	Moderate	Poor
2	2	4	2

Stream Condition
Red
Poor
10

##### Bennett Brook (Section D)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Red	Red
Moderate	Poor	Poor	Poor
4	2	2	2

Stream Condition
Red
Poor
10



**Bennett Brook (Section E)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Yellow
Moderate	Poor	Moderate	Moderate
4	2	4	4

Stream Condition
Yellow
Moderate
14

**Bennett Brook (Section F)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Yellow
Moderate	Poor	Moderate	Moderate
4	2	4	4

Stream Condition
Yellow
Moderate
14

**Bennett Brook (Section G)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Yellow	Yellow	Yellow
Moderate	Moderate	Moderate	Moderate
4	4	4	4

Stream Condition
Yellow
Moderate
16

**Bennett Brook (Section H)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Black	Black	Black
Poor	Very Poor	Very Poor	Very Poor
2	0	0	0

Stream Condition
Black
Very Poor
2

**Bennett Brook (Section I)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Black	Black	Red	Red
Very Poor	Very Poor	Poor	Poor
0	0	2	2

Stream Condition
Black
Very Poor
4



**Bennett Brook (Section J)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Black	Black	Black
Poor	Very Poor	Very Poor	Very Poor
2	0	0	0

Stream Condition
Black
Very Poor
2

**Bennett Brook (Section K)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Black	Red	Red
Poor	Very Poor	Poor	Poor
2	0	2	2

Stream Condition
Red
Poor
6

**Bennett Brook (Section L)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Red	Red
Moderate	Poor	Poor	Poor
4	2	2	2

Stream Condition
Red
Poor
10

**3.5.2 Summary results for Bannister Creek (Canning Catchment)**

**Bannister Creek (Section A)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Red	Red
Moderate	Poor	Poor	Poor
4	2	2	2

Stream Condition
Red
Poor
10

**Bannister Creek (Section B)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Red
Moderate	Poor	Moderate	Poor
4	2	4	2

Stream Condition
Red
Poor
12



**Bannister Creek (Section C)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Black	Black	Black
Poor	Very Poor	Very Poor	Very Poor
2	0	0	0

Stream Condition
Black
Very Poor
2

**Bannister Creek (Section D Left Bank)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Red	Red
Moderate	Poor	Poor	Poor
4	2	2	2

Stream Condition
Red
Poor
10

**Bannister Creek (Section E)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Red	Red
Moderate	Poor	Poor	Poor
4	2	2	2

Stream Condition
Red
Poor
10

**3.5.3 Summary results for Canning River (Canning Catchment)****Canning River (Section A)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Red	Red
Moderate	Poor	Poor	Poor
4	2	2	2

Stream Condition
Red
Poor
10

**Canning River (Section B)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Red
Moderate	Poor	Moderate	Poor
4	2	4	2

Stream Condition
Red
Poor
12



**Canning River (Section C)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Red
Moderate	Poor	Moderate	Poor
4	2	4	2

Stream Condition
Red
Poor
12

**Canning River (Section D)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Red	Red
Moderate	Poor	Poor	Poor
4	2	2	2

Stream Condition
Red
Poor
10

**Canning River (Section E)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Red	Red	Red
Poor	Poor	Poor	Poor
2	2	2	2

Stream Condition
Red
Poor
8

**Canning River (Section F)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Black	Red	Red
Poor	Very Poor	Poor	Poor
2	0	2	2

Stream Condition
Red
Poor
6

**Canning River (Section G)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Black	Black	Black
Moderate	Very Poor	Very Poor	Very Poor
4	0	0	0

Stream Condition
Black
Very Poor
4



### 3.5.4 Summary results for Roley Pool (Canning Catchment)

#### Roley Pool (Section A)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Green
Moderate	Poor	Moderate	Good
4	2	4	6

Stream Condition
Yellow
Moderate
16

#### Roley Pool (Section B)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Green
Moderate	Poor	Moderate	Good
4	2	4	6

Stream Condition
Yellow
Moderate
16

### 3.5.5 Summary results for Southernwood Creek (Canning Catchment)

#### Southernwood Creek (Section A)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Black	Black	Black
Moderate	Very Poor	Very Poor	Very Poor
4	0	0	0

Stream Condition
Black
Very Poor
4

#### Southern River (Section B)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Red
Moderate	Poor	Moderate	Poor
4	2	4	2

Stream Condition
Red
Poor
12



### 3.5.6 Summary results for Wright Brook (Canning Catchment)

#### Canning River upstream of Wright Brook confluence (Section A)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Red	Yellow	Red
Poor	Poor	Moderate	Poor
2	2	4	2

Stream Condition
Red
Poor
10

#### Canning River downstream of Wright Brook confluence (Section B)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Red	Red	Red
Poor	Poor	Poor	Poor
2	2	2	2

Stream Condition
Red
Poor
8

#### Wright Brook (Section C)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Black	Black	Black	Black
Very Poor	Very Poor	Very Poor	Very Poor
0	0	0	0

Stream Condition
Black
Very Poor
0

#### Wright Brook (Section D)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Red	Yellow	Red
Poor	Poor	Moderate	Poor
2	2	4	2

Stream Condition
Red
Poor
10

#### Wright Brook (Section E)

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Red	Yellow	Red
Poor	Poor	Moderate	Poor
2	2	4	2

Stream Condition
Red
Poor
10





**Wright Brook (Section F)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Red
Moderate	Poor	Moderate	Poor
4	2	4	2

Stream Condition
Red
Poor
12

**Wright Brook (Section G)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Red	Yellow	Red
Poor	Poor	Moderate	Poor
2	2	4	2

Stream Condition
Red
Poor
10

**Wright Brook (Section H)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Red	Red	Yellow	Red
Poor	Poor	Moderate	Poor
2	2	4	2

Stream Condition
Red
Poor
10

**Wright Brook (Section I)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Black	Black	Black
Moderate	Very Poor	Very Poor	Very Poor
4	0	0	0

Stream Condition
Black
Very Poor
4

**3.5.7 Summary results for Breera Brook (Ellen Brook Catchment)**

**Breera Brook (Section A)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Yellow	Yellow	Yellow
Moderate	Moderate	Moderate	Moderate
4	4	4	4

Stream Condition
Yellow
Moderate
16



**Breera Brook (Section B)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Green	Yellow	Green
Moderate	Good	Moderate	Good
4	6	4	6

Stream Condition
Yellow
Moderate
20

**Breera Brook (Section C)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Green	Green	Green	Yellow
Good	Good	Good	Moderate
6	6	6	4

Stream Condition
Green
Good
22

**Breera Brook (Section D)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Green	Green	Green	Yellow
Good	Good	Good	Moderate
6	6	6	4

Stream Condition
Green
Good
22

**Breera Brook (Section E)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Green	Yellow	Green	Yellow
Good	Moderate	Good	Moderate
6	4	6	4

Stream Condition
Yellow
Moderate
20

**3.5.8 Summary results for Ellen Brook (Ellen Brook Catchment)**

**Ellen Brook (Section A)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Black	Red	Red	Black
Very Poor	Poor	Poor	Very Poor
0	2	2	0

Stream Condition
Black
Very Poor
4



**Ellen Brook (Section B)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Black	Red	Red	Black
Very Poor	Poor	Poor	Very Poor
0	2	2	0

Stream Condition
Black
Very Poor
4

**Ellen Brook (Section C)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Yellow
Moderate	Poor	Moderate	Moderate
4	2	4	4

Stream Condition
Yellow
Moderate
14

**Ellen Brook (Section D)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Yellow
Moderate	Poor	Moderate	Moderate
4	2	4	4

Stream Condition
Yellow
Moderate
14

**Ellen Brook (Section E)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Yellow	Red	Yellow	Red
Moderate	Poor	Moderate	Poor
4	2	4	2

Stream Condition
Red
Poor
12

**Ellen Brook (Section F)**

Bank Stability	Foreshore Vegetation	Stream Cover	Habitat Diversity
Black	Red	Red	Black
Very Poor	Poor	Poor	Very Poor
0	2	2	0

Stream Condition
Black
Very Poor
4



## 4. Specific site reports

The following Water and Rivers Commission reports contain the detailed foreshore condition assessment and recommended strategies for tributaries surveyed and referred to in this report.

**Water Resource Management Series, No WRM 14.**  
*Foreshore Assessment in the Bennett Brook Catchment.*

This report contains survey results for:

- Bennett Brook, from Mussel Pool to Benara Road, Caversham.

**Water Resource Management Series, No WRM 15.**  
*Foreshore Assessment in the Canning Catchment.*

This report contains survey results for :

- Bannister Creek, Adenia Street to Iveston Road.
- Canning River, Pioneer Park to Herbert Street.
- Roley Pool, Soldiers Road to Thompson Road.
- Southernwood Creek, Southernwood Park.
- Wright Brook, Kelmscott.

**Water Resource Management Series, No WRM 16.**  
*Foreshore Assessment in the Ellen Brook Catchment.*

This report contains survey results for :

- Ellen Brook, north Ellen Brook Nature Reserve to Almeria Pde, Upper Swan.
- Breera Brook



## 5. General recommendations

A number of general recommendations can be identified which apply to all of the sites. These are divided into the core activities, which will be required for groups to successfully develop and implement rehabilitation strategies.

### 5.1 Planning

- Determine cadastral boundaries and landowner/manager and ensure that they support the foreshore assessment process, and are involved in the development and implementation of any remedial strategies.
- Collate as much existing information about the focus waterway and catchment as possible.
- Focus initial foreshore assessment survey work in areas where future rehabilitation projects may be undertaken.
- Extend future foreshore assessment work from previously surveyed areas along the foreshore, eventually mapping all sites. Future surveys may also include re-assessment of earlier surveys to assess changes to the environment.
- Create herbariums of native and weed species to teach group members and other interested parties to distinguish between native and introduced plants present in the rehabilitation zone.
- Ensure that all works are planned well in advance and that a long-term strategy has been developed and is amended as new information becomes available.
- Ensure that all agencies with statutory responsibilities such as the relevant local government authority, Water Corporation and Swan River Trust is advised of any works within their management areas, to ensure that the works meet all of the legislative requirements.
- Develop information brochures to increase community awareness of the importance of foreshore areas and to encourage community involvement in managing their own foreshores and surrounding reserves.
- Develop an information brochure for the landholder to suggest methods of improved land management practices and encourage rehabilitation of the foreshore area.

- Endeavour to source funds from outside sources to assist both the group and any private landholders that are willing to implement rehabilitation activities.

### 5.2 Site preparation

#### 5.2.1 Weed control

- Ensure weed control activities are undertaken in manageable sized nodes, reinforcing overstorey species and restoring the middlestorey and understorey species (using species recommended in Appendix 3 of this report) once weeds have been eradicated.
- Tag any native plants present to protect them from weed control activities.
- Hand weed where possible, especially annual weeds and instream weeds.
- Use a qualified herbicide operator if chemical control is undertaken near waterways.
- Always consider the impacts that weed control will have on habitat, particularly for reptiles and small mammals such as bandicoots. Maintain vegetated corridors for animals to move within until sufficient native plants have re-established.
- Ensure that all weeds are removed from the site to limit re-infestation.
- Create buffers around existing clumps of native vegetation to encourage natural regeneration of existing plants e.g. spray Fusillade around native rushes to control introduced grasses and enable the clumps of rushes to spread naturally.
- Ensure the impact on bank stability is considered before weed control works are undertaken. Consider potential for use of erosion control matting as an option to reduce weed re-emergence, support plants installed and improve bank stability on steeper gradient banks.



### 5.2.2 General site preparation

- Encourage landholders throughout the rural and semi-rural catchments to fence off waterways and tributaries and implement broadscale revegetation program.
- Provide financial support or material assistance to landholders willing to implement rehabilitation activities.
- Define access tracks to weed management areas or where there are planting programs, to minimise disturbance and limit damage to existing vegetation and the substrate.
- Implement intensive weed control activities in manageable sized nodes where planting will be undertaken.
- Remove flower heads prior to seeding to limit reinforcement of the weed seed bank.
- In broadscale areas proposed for future works or in high-risk areas of dense weeds with few native plants where complete removal is inappropriate, ensure either flower removal or repeated brushcutting occurs prior to seeding.

### 5.3 Planting out

- Ensure planted areas within streamlines are artificially stabilised and planted in low-flow conditions to enable sufficient time for establishment, to reduce the chance for plants to be washed out during peak flows.
- Plant native species only in areas where weeds have been effectively controlled and managed for a preferred minimum of two seasons.
- Encourage landholders to ensure all strata of vegetation including understorey, middlestorey and upperstorey species are included in revegetation works to reinforce bank stability.
- Plant overstorey species initially in highly exposed regions lacking vegetation, to create a level of cover and protection for future plantings.
- Plant emergent and wetland plants in permanent water between September and March, securing those planted in flowing water with 600mm steel “U” shaped pegs.

- Plant dryland plants and seasonally inundated areas in May to July.
- Plant in higher densities than ultimately required to create instant habitat and improve weed exclusion; particularly in the inner urban environments.
- Obtain professional advice about planting densities for each recommended species, to optimise chances of success and re-creating a more natural ecosystem.

### 5.4 Maintenance

- Ensure the works program includes ongoing intensive maintenance of areas where weed control and planting works have previously been undertaken.
- Implement ongoing weed management, prior to commencing site preparation and planting works in new areas.
- Monitor for any natural regeneration on a regular basis, and undertake weed control around any emerging native plant seedlings.
- Assess the effectiveness of any river restoration works or installation of any products such as hemp matting and modify as required.
- Determine the impact of vandalism if any, and develop and implement strategies to manage this problem.

### 5.5 Monitoring

- Continue to use the method to assess changes and improvement to foreshore health over time.
- Assess the effectiveness and relative benefits of different management techniques utilised and update the works program accordingly.
- Document the results and learn from experience.
- Monitor the effectiveness of sustaining interest within the project at both the management and implementation level. Develop techniques to support community groups and individuals in undertaking this work.
- Minimise the potential for burnout by not over-extending limited resources, particularly labour.



## 6. Common issues

### 6.1 Ownership and access

It is essential that cadastral boundaries are determined at each site and that the people implementing the foreshore assessment are aware of who owns the land. Permission is required from the landowners, who may be State or local government agencies or private landholders, prior to undertaking any survey work. Gaining access to private property may prove to be difficult, whilst permission to enter most government managed lands is generally readily available.

Often property boundaries are fenced and landowners may be suspicious that any information collected during surveys along their foreshore will eventually be used against them. It is important that people implementing the survey are clear about the process and the reasons for the survey and approach all landholders. Where landholder agreement cannot be readily obtained, it is important not to waste time and resources in excessive negotiations. Locate landholders that are interested in improving the health of their foreshore and assist these properties to enhance their land. Healthy foreshores can increase property values and through discussion within communities can ultimately result in peer pressure on others to work on protecting their waterways.

There are often conflicting perceptions about the requirements for managing riparian zones and determining what is a healthy foreshore. Many landholders consider lawn to the high water mark with occasional trees to be healthy and providing sufficient habitat for example, as large numbers of birds e.g. black ducks, may frequently use the foreshore. It is very difficult to articulate foreshore management issues until a common perception of a stable, intact waterway is developed between the group doing the work and the wider community.

A further conflict can arise when landholders consider that their current foreshore management program is adequate. For example, as well maintained lawns reduce the fire hazard, limits uncontrolled weed growth and keep the streambed free of debris, it is argued by these private landholders to be an appropriate management technique to protect the waterway. Frequently this

management regime is in contrast to management practices in neighbouring foreshore reserves that are managed by State and Local government authorities. Extensive weeds, limited access and considerable fire risk are often features of these reserves. As a result it is perceived that there is little management effort. In situations where State and Local government authorities are not demonstrating best management practice, it is difficult to discourage landholders from maintaining their own inappropriate management program.

Both State and Local government and the wider community need to implement improved foreshore management.

### 6.2 Developing management and rehabilitation plans

Management plans are an important tool used to strike a balance between the multiple use demands of foreshore areas and the protection of flora, fauna and water quality. These documents should have clearly defined aims, objectives and visions as ultimately, the final use of the land will affect how, where and if, rehabilitation plans need development and implementation.

If, for example, a grassed area occurs adjacent to a waterway which is a high use recreation zone, then extensive revegetation works are likely to impinge on the purpose of the land and therefore may be inappropriate. A compromise position may need to be negotiated such as establishing a narrow buffer zone immediately along the stream banks, with well defined access points for viewing the waterway. The buffer zone needs to have a clearly defined boundary between any lawn areas and native vegetation to avoid trampling of native seedlings.

All issues associated with development, conservation and management of the waterway and associated land need to be addressed prior to the development of any plans. Community needs and visions for particular areas need to be canvassed to ensure the document reflects community attitude, which affects whether or not plans get implemented.



At the next level, following management planning there is a need to develop a complete rehabilitation plan for the waterway. It is essential to extend the assessment of foreshore condition to the length of the waterway, prior to any works to gain a complete understanding of current health. This may be limited by access issues, however the broader the understanding of the waterway and their tributaries, the better.

An ecosystem approach to management will ensure that appropriate rehabilitation plans are developed minimising the impact of any activities. For example complete eradication of dense weeds along the immediate foreshore results in acute loss of habitat and may destabilise foreshore banks increasing the danger of severe erosion and bank collapse. It is necessary to undertake weed control in small, manageable sized nodes to ensure that eradicated weeds are immediately replaced with deep-rooted native species, to minimise the impact on bank stability, and protect native fauna.

Developing detailed management and rehabilitation plans and having a clear understanding of the works required over the long term, enables the development of detailed budgets, allocation of funding or opportunities to raise funds to ensure the completion of any project.

### 6.3 Long term management

The rehabilitation planning process should include a maintenance schedule for any existing works as well as directing future projects. The importance of continued maintenance within current project sites prior to beginning any new works, can not be emphasised enough. Ongoing management in the long term must be scheduled to ensure the success of any rehabilitation works. Weed control needs to be continued indefinitely as there will always be the threat of reinfestation.

Undertaking works on crown land and reserves requires ongoing community commitment and an interest from state and local government agencies to provide assistance such as fire break maintenance and provision of qualified herbicide operators to undertake weed control.

Private landowners must be strongly committed to any project undertaken on their property to ensure ongoing maintenance. Any change in ownership may require a negotiation with the new owners to determine if management will continue.

Once a rehabilitation project has commenced on a property it will require a significant amount of time to implement weed control, planting and maintenance. Setting manageable areas for work and achievable targets is the most effective way to ensure success. Over-extension of limited resources frequently causes the areas to degrade further, resulting in a situation that is worse than prior to any rehabilitation effort.

There is nothing more disillusioning than having put considerable effort into developing and implementing works for little or no benefit in the medium to long term.

### 6.4 Surrounding landuse

Adjacent landuse can have a considerable impact on the riparian zone and waterway health. Different landuses have different implications for stream health and therefore the appropriate management regimes required will vary.

Riparian zones are often highly degraded. Foreshore vegetation is frequently reduced to a few metres either side of the watercourse. It is important to provide information to landholders and land managers about the benefits of undertaking remedial works along foreshores, emphasizing the importance of fencing off riparian areas and excluding stock. Sourcing funds and providing support may encourage interested landholders to undertake intensive weed control and revegetation works.

Foreshores in urban areas are frequently high use recreation sites. Traditionally large open areas of maintained lawn were favoured over dense stands of native vegetation. Advertising campaigns and creating signage around project sites are useful tools to increase community awareness. Providing detailed information on the benefits of replanting native species such as stabilizing foreshore banks and increasing stream cover and habitat diversity will increase awareness and may encourage local residents to become involved in the projects.

Sedimentation of watercourses is generally an indication of erosion occurring further upstream. No system can be considered in total isolation, as there will always be impacts from activities further upstream. When undertaking any projects it is essential that groups have a clear understanding of the surrounding landuse and the condition of any tributaries feeding into the main waterway.





The impact of new subdivisions or earthworks upstream should be carefully monitored. Weeds may invade from nearby residential housing. Subdivisions can also have a huge impact on water regimes and sediment loads entering streams and tributaries. Early detection of potential threats minimises the impact on foreshore health in the long term if remedial action is undertaken immediately.

## 6.5 Gaining support from state and local government

State and local government have a significant role to play in supporting foreshore rehabilitation. Many agencies are also directly involved in managing waterways and foreshore areas. Water Corporation, Water and Rivers Commission, Swan River Trust, Agriculture WA and local government authorities all actively manage some waterways within the State.

Many of these agencies also have statutory requirements to meet, that relate to management of these areas. The Swan River Trust Management Area, for example, relates to the bed and banks of the Swan and Canning Rivers extending across the riparian zone to the limit of the Parks and Recreation Reservation. It is illegal to undertake any works within the SRT Management Area without notifying the Swan River Trust.

Some agencies also have community support functions to assist groups to undertake hands on work, prepare management and rehabilitation plans and can also provide some support for administrative and information requirements.

Key contacts include:

Contact	Agency Contact	Number
Ecoplan	Department of Environmental Protection	9222 7000
	Swan Catchment Centre	9221 3840
	Water and Rivers Commission	9278 0300
	Swan River Trust	9278 0400
	Agriculture WA	9368 3333
	Relevant local government authority	White pages

There may be contacts within each agency for on-ground support. The Swan Catchment Centre has a Landcare trailer that is fully rigged for landcare activities and provides the relevant equipment for site preparation, weed control and planting.

Where reserves are managed by a state or local government authority it is essential that the community liaises with the land manager to develop and implement any assessment and rehabilitation projects.

Support from agencies also improves the opportunities for gaining funding from external sources such as Greening WA, Lotteries WA and the Natural Heritage Trust.

## 6.7 Fire management

Fire is not recommended as a management technique for riparian zones, particularly in the Scarp region and areas with peaty soils. Should fire occur as a result of vandalism or an accidental burn, then advantage should be taken of the increased access to the area for weed control activities.

Prescribed burns are likely to do significant damage to fringing vegetation, the seed bank and may result in reduced bank stability and higher levels of erosion. Fire also often encourages further weed invasion and spread of existing weed species. Autumn burns are particularly risky.

## 6.8 Access to information

State and local government authorities have considerable information resources about waterways and should be contacted. Many agencies also have libraries that the community can access, however borrowing books is generally not permitted.

Existing information about any particular waterway should be collated prior to development of management plans.

General information about weed control techniques, site preparation and stream and foreshore restoration needs to be obtained prior to the development of rehabilitation plans.



## 7. Summary

The foreshore assessment process has been developed to aid interested community groups, officers of State and local government authorities and private landholders in urban and semi-rural areas to gain an understanding of the condition of foreshore areas within their own community. By using a standard method to gather information it is possible to compare and contrast foreshore condition of the same area over time, or between different sites in the same survey season to prioritise works.

This document provides the results of the first series of foreshore assessments undertaken in accordance with the Water and Rivers Commission (1999) foreshore condition assessment method. Testing and refining the assessment protocol in this work was intended to identify any shortcomings or limitations of the method.

The assessments were undertaken along sections of the Canning and Southern Rivers, Bannister Creek, Bennett Brook and Ellen Brook.

Implementing the technique has resulted in a limited number of modifications to the methodology and provided considerable documentation for the surveyed sections of the waterways listed above.

The foreshore sites selected for this baseline study ranged in condition and current management practices. The detailed recommended strategies outlined for each of these sites aim to give suggestions for hands-on works for rehabilitation of degraded foreshore systems. General recommendations have been provided for broadscale long term planning which emphasise the need to consider the implications of any works, and the commitment required to sustain these activities in the long term.

This report of foreshore condition will be the first of many, as the process continues to evolve and be implemented across urban and semi-rural areas statewide.



## 8. References and recommended reading

- APACE Green Skills and Pen, L.J. (1997) *Survey of river foreshores in the Oyster Harbour Catchment*. Water and Rivers Commission Report, Water Resource Technical Series No. 17.
- Bush, B. Maryan, B. Browne-Cooper, R. and Robinson, D. (1995) *A guide to the reptiles and frogs of the Perth region*. University of Western Australia Press, Nedlands, Western Australia.
- Hussey, B.M.J. Hobbs, R.J. and Saunders, D.A. (1989) *Guidelines for bush corridors*. Surrey Beatty & Sons.
- Hussey, B.M.J. and Wallace, K.J. (1993) *Managing your bushland*. Department of Conservation and Land Management.
- Land and Water Resources Research and Development Corporation (1996) *Managing riparian land*. Riparian Management Series (1), September 1996.
- Land and Water Resources Research and Development Corporation (1996) *Water quality*. Riparian Management Series (3), November 1996.
- Land and Water Resources Research and Development Corporation (1996) *River ecosystems*. Riparian Management Series (4), November 1996.
- Land and Water Resources Research and Development Corporation (1996) *Land-based ecosystems*. Riparian Management Series (5), November 1996.
- Land and Water Resources Research and Development Corporation (1998) *Managing snags in rivers*. Riparian Management Series (7), February 1998.
- Pen, L.J. (1992) *Fringing vegetation of the Lower Collie and Brunswick River 1992*. Leschenault Management Authority, Waterways Commission Report No. 37.
- Pen, L.J. (1993) *Fringing vegetation of the Canning, Southern and Wungong Rivers*. Swan River Trust Report No. 7.
- Pen, L.J. (1994) *Condition of the foreshore vegetation of the Kalgan River 1992*. Albany Waterways Management Authority, Waterways Commission Report.
- Pen, L.J. and Majer, K. (1993) *Living streams - a guide to bringing watercourses back to life in south-west Western Australia*. Waterways Commission, Perth, Western Australia.
- Pen, L.J. and Scott, M. (1995) *Stream foreshore assessment in farming areas*. Blackwood Catchment Coordinating Group, Western Australia.
- Riding, T. and Carter, R. (1992) *The importance of the riparian zone in water resource management*. Water Resources, Canberra.
- Scheltema, M. and Harris, J. (Ed) (1996) *Managing Perth's Bushland*. Greening WA, Perth.
- Siemon, N. (1994) *Lower Canning River Management Plan*. Report No 23, Swan River Trust, Perth WA.
- Siemon, N., Davis, G., Hubbard, P. and Duckworth, A. (1993) *Fringing vegetation of the Serpentine River in the Shire of Serpentine-Jarrahdale and City of Rockingham*. Waterways Commission Report. No. 38.
- Water and Rivers Commission (1996) *Macroinvertebrates and water quality*. Water facts 2, September 1996.
- Water and Rivers Commission (1997) *River and estuary pollution*. Water facts 3, July 1997.
- Water and Rivers Commission (1998) *Living streams*. Water facts 4, January 1998.
- Water and Rivers Commission (1998) *Taking water from streams and lakes*. Water facts 5, May 1998.
- Water and Rivers Commission (1998) *Water words*. Water facts 1, January 1998.
- Water and Rivers Commission (1999) *Planning and Management: Foreshore Condition Assessment in urban and semi-rural areas of south-west Western Australia*. Water and Rivers Commission Report No. RR2.
- Woodcock, S. (1994) *Collie and Brunswick Rivers foreshore study*. Leschenault Inlet Management Authority, Waterways Commission Report No. 48.



# Appendix 1

Native and weed species identified  
during the foreshore assessment  
process (1998)



## Appendix 1a: Native Species identified during the foreshore assessment process (1998)

Scientific name	Common Name	Bennett Brook	Bannister Creek	Canning River	Roley Pool	Southernwood Creek	Wright Brook	Ellen Brook	Breera Brook
<i>Acacia alata</i>	Winged wattle		Y		Y		Y		
<i>Acacia pulchella</i>	Prickly moses	Y			Y		Y		
<i>Acacia saligna</i>	Coojong	Y		Y	Y	Y	Y	Y	Y
<i>Acanthocarpus preissii</i>					Y				
<i>Agonis linearifolia</i>	Swamp peppermint	Y	Y	Y	Y		Y	Y	Y
<i>Astartea fascicularis</i>	Common astartea				Y			Y	
<i>Banksia menziesii</i>	Firewood banksia	Y							
<i>Baumea juncea</i>	Bare twigrush		Y					Y	Y
<i>Bolboschoenus caldwellii</i>	Marsh club rush		Y						
<i>Baumea rubiginosa</i>	River twigrush		Y						
<i>Burchardia umbellata</i>	Milkmaid								Y
<i>Caladenia</i> spp.	Orchids	Y							
<i>Carex appressa</i>	Tall sedge		Y					Y	Y
<i>Carex divisa</i>	Divided sedge		Y						
<i>Carex fascicularis</i>	Tassel sedge		Y					Y	Y
<i>Centella cordifolia</i>	Centella	Y	Y	Y				Y	Y
<i>Centrolepis</i> spp.	Centrolepis		Y						
<i>Chenopodium glaucum</i>	Glaucous goosefoot		Y						
<i>Corynotheca micrantha</i>	Sand lily		Y						Y
<i>Conostylis</i> spp.		Y							
<i>Corynotheca micrantha</i>					Y				
<i>Corymbia calophylla</i>	Marri	Y	Y	Y	Y	Y	Y	Y	Y
<i>Cotula coronopifolia</i>	Button weed	Y	Y						
<i>Darwinia citriodora</i>	Lemon-scented darwinia								
<i>Drosera erythrorhiza</i>	Red ink sundew								Y
<i>Drosera glanduligera</i>	Pimpernel sundew								Y
<i>Dryandra nivea</i>	Couch honeypots				Y				Y

<i>Dryandra sessilis</i>	Parrot bush								Y
<i>Eucalyptus rudis</i>	Flooded gum	Y	Y	Y	Y	Y	Y	Y	Y
<i>Eucalyptus wandoo</i>	Wandoo				Y				
<i>Gahnia decomposita</i>								Y	Y
<i>Gastrolobium spinosum</i>	Prickly poison				Y		Y		
<i>Gompholobium tomentosum</i>	Hairy yellow pea	Y							
<i>Grevillea</i> spp.					Y		Y		Y
<i>Hakea prostrata</i>	Harsh hakea								Y
<i>Hakea varia</i>	Variable leaf hakea		Y						
<i>Hardenbergia comptoniana</i>	Native wisteria	Y	Y						
<i>Hemiandra pungens</i>	Snake bush								
<i>Hibbertia</i> spp.	Native buttercups	Y			Y		Y		Y
<i>Hypocalymma angustifolium</i>	White myrtle	Y	Y					Y	Y
<i>Isolepis setiformis</i>	Tufted sedge		Y						
<i>Jacksonia furcellata</i>	Grey stinkwood	Y	Y					Y	Y
<i>Jacksonia sternbergiana</i>	Green stinkwood	Y	Y				Y		Y
<i>Juncus amabilis</i>	Blue rush		Y						
<i>Juncus kraussii</i>	Sea Rush		Y						
<i>Juncus pallidus</i>	Pale rush	Y	Y	Y	Y		Y	Y	Y
<i>Kennedia prostrata</i>	Running postman	Y	Y				Y		
<i>Kunzea</i> spp.								Y	Y
<i>Lasiopetalum bracteatum</i>	Helena velvet bush				Y				
<i>Lepidosperma effusum</i>	Spreading sword sedge	Y	Y					Y	Y
<i>Lepidosperma longitudinale</i>	Pithy sword-sedge	Y						Y	Y
<i>Lepidosperma tetraquetrum</i>	Angle sword-sedge	Y			Y				Y
<i>Lomandra</i> spp.								Y	Y
<i>Lyginia barbata</i>								Y	Y
<i>Macrozamia riedlei</i>	Zamia	Y			Y		Y		Y
<i>Melaleuca lateritia</i>	Robin redbreast bush		Y						
<i>Melaleuca preissiana</i>	Modong	Y	Y					Y	Y
<i>Melaleuca raphiophylla</i>	Swamp paperbark	Y	Y	Y	Y	Y	Y	Y	Y

<i>Mesomeleana tetragona</i>	Semaphore sedge		Y			Y		
<i>Oxylobium lineare</i>	Narrow-leaved oxylobium	Y	Y		Y		Y	Y
<i>Paraserianthes lophantha</i>	Albizia	Y					Y	Y
<i>Patersonia occidentalis</i>	Purple flag						Y	Y
<i>Persicaria decipiens</i>	Slender knotweed		Y					
<i>Phyllanthus calycinus</i>	False boronia							Y
<i>Potamogeton crispus</i>	Curly pondweed		Y					
<i>Pteridium esculentum</i>	Bracken fern	Y		Y		Y	Y	Y
<i>Regelia ciliata</i>			Y					
<i>Schoenoplectus validus</i>	Lake club rush		Y					
<i>Schoenus</i> spp.							Y	Y
<i>Schoenus subfascicularis</i>	Bog rush						Y	Y
<i>Thomasia foliosa</i>					Y	Y		
<i>Thomasia macrocarpa</i>					Y	Y		
<i>Verticordia</i> spp.	Feather flowers							Y
<i>Viminaria juncea</i>	Swishbush		Y		Y	Y	Y	Y
<i>Xanthorrhoea preissii</i>	Blackboy	Y			Y	Y	Y	Y

## Appendix 1b: Weed Species identified during the foreshore assessment process (1998)

Scientific name	Common Name	Bennett Brook	Bannister Creek	Canning River	Roley Pool	Southernwood Creek	Wright Brook	Ellen Brook	Breera Brook
<i>Acacia</i> spp.	Weed wattles		Y	Y	Y		Y		
<i>Allium triquetrum</i>	Three-cornered garlic								
<i>Alopecurus myosuroides</i>	Slender foxtail		Y						
<i>Alternanthera nodiflora</i>	Joyweed	Y	Y						
<i>Anagallis arvensis</i>	Pimpernel					Y		Y	
<i>Aponogeton elongatus</i>		Y						Y	
<i>Arctotheca calendula</i>	Capeweed	Y		Y		Y	Y	Y	
<i>Arundo donax</i>	Giant reed	Y	Y	Y	Y	Y	Y	Y	
<i>Aster subulatus</i>	Bushy starwort		Y						
<i>Avena fatua</i>	Wild oats		Y	Y		Y	Y	Y	
<i>Briza</i> spp.	Blowfly grass, shivery grass		Y		Y			Y	
<i>Bromus diandrus</i>	Great brome					Y		Y	
<i>Canna</i> spp.	Canna lily			Y			Y		
<i>Centaurea</i> spp.	Thistles								
<i>Chenopodium album</i>	Fathen		Y						
<i>Conyza</i> spp.	Fleabane	Y	Y	Y		Y	Y	Y	
<i>Cortaderia selloana</i>	Pampas grass		Y	Y			Y		
<i>Cynodon dactylon</i>	Couch grass			Y		Y	Y	Y	
<i>Cyperus difformis</i>	Dirty dora							Y	
<i>Cyperus involucreatum</i>	Cyperus	Y	Y				Y		
<i>Cytisus proliferus</i>	Tree lucerne						Y		
<i>Cyperus</i> spp.			Y	Y					
<i>Dipogon lignosus</i>	Dolichos pea						Y		
<i>Echinochloa telmatophila</i>	Swamp barnyard grass		Y						
<i>Echium plantagineum</i>	Paterson's curse						Y	Y	
<i>Ehrharta calycina</i>	Perennial veldt grass	Y	Y		Y			Y	
<i>Ehrharta longiflora</i>	Annual veldt grass					Y	Y	Y	



<i>Eragrostis curvula</i>	African lovegrass			Y	Y	Y	Y	Y	
<i>Erodium moschatum</i>	Musky crowfoot					Y			
<i>Erythrina x sykesii</i>	Coral tree		Y	Y			Y		
<i>Ferraria crispa</i>	Black flag iris	Y							
<i>Foeniculum vulgare</i>	Fennel		Y						
<i>Ficus</i> spp.	Edible Fig Tree	Y	Y	Y			Y		
<i>Freesia aff. leichtlinii</i>	Freesia	Y							
<i>Fumaria capreolata</i>	Whiteflower fumitory	Y		Y		Y	Y	Y	
<i>Gladiolus</i> spp.	Gladiolus								
<i>Gomphocarpus fruticosus</i>	Cotton bush			Y			Y		
<i>Hesperantha falcata</i>		Y						Y	
<i>Homeria flaccida</i>	One leaf cape tulip	Y						Y	
<i>Hordeum leporinum</i>	Barley grass					Y			
<i>Hyparrhenia hirta</i>	Tambookie grass						Y	Y	
<i>Hypochaeris radicata</i>	Flatweed								
<i>Ipomoea</i> spp	Morning glory		Y	Y			Y		
<i>Isolepis prolifera</i>			Y						
<i>Juncus articulatus</i>									
<i>Juncus capitatus</i>				Y					
<i>Juncus microcephalus</i>		Y	Y		Y		Y	Y	Y
<i>Lantana camara</i>	Lantana		Y				Y		
<i>Leptospermum laevigatum</i>	Victorian tea-tree								
<i>Lolium</i> spp.	Ryegrass							Y	
<i>Lupinus</i> spp.	Lupins	Y					Y	Y	
<i>Medicago</i> spp.	Medics							Y	
<i>Monopsis debilis</i>			Y						
<i>Myrsiphyllum asparagoides</i>	Bridal creeper		Y	Y	Y	Y	Y		
<i>Narcissus tazetta</i>	Jonquils					Y			
<i>Nerium oleander</i>	Oleander						Y		
<i>Olea europaea</i>	Olive Tree								
<i>Oxalis pes-caprae</i>	Soursob	Y	Y	Y	Y	Y	Y	Y	

<i>Panicum capillare</i>	Witchgrass		Y						
<i>Paspalum</i> spp.	Paspalum	Y		Y	Y	Y	Y		
<i>Pelargonium capitatum</i>	Geranium	Y							
<i>Pennisetum clandestinum</i>	Kikuyu	Y	Y	Y	Y	Y	Y	Y	
<i>Plantago lanceolata</i>	Ribwort plantain	Y		Y	Y		Y	Y	
<i>Populus</i> spp.	Poplars		Y	Y					
<i>Raphanus raphanistrum</i>	Wild radish			Y	Y	Y			
<i>Rhynchelytrum repens</i>	Red natal grass				Y		Y	Y	
<i>Ricinus communis</i>	Castor oil	Y	Y	Y		Y	Y		
<i>Romulea rosea</i>	Guildford grass					Y		Y	
<i>Rorippa nasturtium-aquaticum</i>	Watercress	Y	Y		Y	Y			
<i>Rubus fruticosus</i>	Blackberry	Y	Y	Y	Y				
<i>Rumex</i> spp.	Dock	Y	Y	Y				Y	
<i>Salix</i> spp.	Willows	Y	Y	Y		Y			
<i>Schinus terebinthifolia</i>	Japanese pepper	Y	Y	Y					
<i>Solanum nigrum</i>	Deadly nightshade	Y	Y	Y			Y	Y	Y
<i>Stachys arvensis</i>	Staggerweed							Y	
<i>Stenotaphrum secundatum</i>	Buffalo grass								Y
<i>Taraxacum officinale</i>	Dandelion	Y	Y	Y		Y	Y	Y	
<i>Thunbergia alata</i>	Black-eyed Susan			Y					
<i>Trifolium</i> spp.	Clover	Y						Y	
<i>Tropeolum</i> spp.	Nasturtium		Y	Y					
<i>Typha orientalis</i>	Bulrush	Y	Y	Y			Y		Y
<i>Ursinia anthemoides</i>	Ursinia							Y	
<i>Vicia sativa</i>	Vetch	Y	Y	Y		Y		Y	
<i>Vinca major</i>	Periwinkle								
<i>Watsonia bulbifera</i>	Watsonia	Y		Y	Y	Y	Y	Y	
<i>Zantedeschia aethiopica</i>	Arum lily	Y	Y	Y					



# Appendix 2

## Suggested weed control methods



## Appendix 2: Suggested weed control methods

Some of the information contained in this report has been taken from Dixon and Keighery (1995) in *Managing Perth's Bushlands* or referenced to Kings Park Board.

<i>Species Name:</i>	<i>Acacia spp</i>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Weed wattles	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input checked="" type="checkbox"/>
<i>Seed Form:</i>	Light seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input checked="" type="checkbox"/>
<i>Seeding Time:</i>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<i>Method of Spread:</i>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>	Species dependent - prior to flowering				Grass <input type="checkbox"/>
<i>Method of Control:</i>	Hand weed juvenile plants. Small plants means they are relatively easy to remove. Once plants are mature or woody stemmed, cut the main trunk/stem below the widest part of the stem beneath the ground. This will effectively kill all wattles.				
<i>Species Name:</i>	<i>Allium triquetrum</i>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Three cornered garlic	3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<i>Seed Form:</i>			Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<i>Method of Spread:</i>	Spreads by bulb or corm growth				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>					Grass <input type="checkbox"/>
<i>Method of Control:</i>	Apply Glyphosate 1 in 50 or Glean whilst plants are in flower. Repeat applications will be necessary.				
<i>Species Name:</i>	<i>Alopecurus myosuroides</i>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Slender foxtail	3	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<i>Seed Form:</i>			Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input type="checkbox"/>
<i>Method of Spread:</i>					Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>					Grass <input checked="" type="checkbox"/>
<i>Method of Control:</i>	Hand weeding prior to seeding is effective. Herbicides are not recommended as this plant occurs in wetlands and there is a threat of contamination.				
	Repeated brushcutting prior to seeding is effective and reduces the rate of spread of this plant.				

*Control priority 1 - Major environmental weed, urgent control required*

*Control priority 2 - Nuisance weed, control as soon as possible*

*Control priority 3 - Minor weed, control as resources become available*

*Ecosystem Management Services 1999*



<b>Species Name:</b>	<b><i>Alternanthera nodiflora</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Joyweed	1	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	March-April		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Oct-Nov				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weed plants in strips up to 2 m perpendicular to water flow and replace immediately with native emergent species. Carefully bag and remove weed material from the site.				Climber <input type="checkbox"/>

Any segment which is broken from this plant is likely to regenerate into a new plant, so using a floating bund with netting or similar device downstream to trap any segments missed.

<b>Species Name:</b>	<b><i>Anagallis arvensis</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Pimpernel	3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>					Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weeding small populations is effective. Alternatively treat with Glyphosate or Glean at 15g per ha.				Climber <input type="checkbox"/>

<b>Species Name:</b>	<b><i>Aponogeton elongatus</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>		2	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input checked="" type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Nov - Mar (access dependent)				Grass <input type="checkbox"/>
<b>Method of Control:</b>	This aquatic weed is difficult to control because it slows water movement, increases sedimentation and reduces erosion which affects bed and bank stability following removal. The recommended removal technique involves manual clearing of a channel and also clearing 5 to 10 m wide bands, 20 metres apart which are perpendicular to the stream flow. This will minimise the potential for de-stabilising the stream bed.				Climber <input type="checkbox"/>
	Seek expert advice and approvals from the relevant government agencies prior to implementing broad scale works. Herbicides should not be used for this weed. Shading out and planting dense clumps of indigenous plants are the most effective management techniques.				

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**

<b>Species Name:</b>	<b>Arctotheca calendula</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Capeweed	3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Oct - Feb				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weeding small populations of this plant is effective. Rotary hoeing broadscale infestations repeatedly can also work. Kings Park Board recommends glyphosate at 100ml in 15l water. Lontrel 1 in 100 has been successful on larger plants in areas without any native vegetation.				

<b>Species Name:</b>	<b>Arundo donax</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Giant reed	2	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Sept - Dec		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads readily from rhizome growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	All year				Grass <input checked="" type="checkbox"/>
<b>Method of Control:</b>	Cut down and spray regrowth when 0.5 - 1.0m high with Glyphosate 360 100ml in 10l of water. An alternative technique is to remove bulk of plant material and pour herbicide down each tube.  Ensure removal of seed heads prior to ripening if plant control is not possible. Generally this plant occurs on the banks of streams and rivers. It is important not to dig this plant out if there is a risk of increasing erosion. Onsite poisoning is the preferred option leaving the dense rhizome mat intact.				

<b>Species Name:</b>	<b>Aster subulatus</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Bushy starwort	3	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light and easily spread by wind		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Aug - Mar				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weeding these plants is easy and effective. It is essential to weed them prior to flowering and fruiting to reduce their spread.				

<b>Species Name:</b>	<b>Avena spp.</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Wild Oats	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	March - June		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Aug - Oct				Grass <input checked="" type="checkbox"/>
<b>Method of Control:</b>	Hand weeding small plants in winter is effective for small populations. Blanket/Spot spraying at 2l Fusillade per ha is effective. Brushcutting plants with immature seed heads will aid control in the longer term by minimizing seed spread.  Dense populations represent a significant fire hazard and threat to remnant vegetation, so repeated brushcutting also assists in reduction of fire hazard.				

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



**Species Name:** *Briza maxima* **Control Priority** **Location** **Habit** **Form**

**Common Name:** Blowfly grass **2** Dryland  Bulb/Corm  Tree   
 Riparian  Perennial  Shrub   
 Aquatic  Annual  Herb   
 Rush/Sedge   
 Grass   
 Climber

**Seed Form:** Light, easily spread by wind

**Seeding Time:** Sept - Nov

**Method of Spread:** Spreads mostly from seed

**Best Time of Control:** June - Aug

**Method of Control:** Hand weeding is effective.

Control may be achieved by spot/blanket spraying Sertin or similar at 2l per ha.

**Species Name:** *Briza minor* **Control Priority** **Location** **Habit** **Form**

**Common Name:** Shivery grass **2** Dryland  Bulb/Corm  Tree   
 Riparian  Perennial  Shrub   
 Aquatic  Annual  Herb   
 Rush/Sedge   
 Grass   
 Climber

**Seed Form:** Light, easily spread by wind

**Seeding Time:** Sept - Oct

**Method of Spread:** Spreads mostly from seed

**Best Time of Control:** June - Aug

**Method of Control:** Hand weeding is effective.

Control may be achieved by spot/blanket spraying Sertin or similar at 2l per ha.

**Species Name:** *Bromus diandrus* **Control Priority** **Location** **Habit** **Form**

**Common Name:** Great brome **2** Dryland  Bulb/Corm  Tree   
 Riparian  Perennial  Shrub   
 Aquatic  Annual  Herb   
 Rush/Sedge   
 Grass   
 Climber

**Seed Form:** Coarse seed

**Seeding Time:** Sept - Nov

**Method of Spread:** Spreads mostly from seed

**Best Time of Control:** June - Aug

**Method of Control:** Hand weeding is easy and effective for small populations. The most frequently recommended treatment is Fusillade at between 2-4l per ha, when the plants are actively growing in winter. Repeated brushcutting can also be effective.

Note: Correct identification of grasses is important to protect native grasses from removal. The presence of native grasses should be investigated prior to spraying herbicides.

**Species Name:** *Canna spp.* **Control Priority** **Location** **Habit** **Form**

**Common Name:** Canna **3** Dryland  Bulb/Corm  Tree   
 Riparian  Perennial  Shrub   
 Aquatic  Annual  Herb   
 Rush/Sedge   
 Grass   
 Climber

**Seed Form:** Heavy seed

**Seeding Time:**

**Method of Spread:** Spreads readily from rhizome growth

**Best Time of Control:** Sept - Apr

**Method of Control:** Dig out small infestations. Selectively spraying the leaves with a systemic herbicide can be effective.

Encourage residents to harvest the flowers to reduce seed production.

Broadscale removal of dense stands may threaten bank stability. Remove in nodes perpendicular to the water course or remove the bulk of biomass then treat with herbicide. Ensure the dense rhizome mat intact.

Control priority 1 - Major environmental weed, urgent control required  
 Control priority 2 - Nuisance weed, control as soon as possible  
 Control priority 3 - Minor weed, control as resources become available

Ecosystem Management Services 1999





<b>Species Name:</b>	<b>Centaurea spp</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Thistles	<input type="checkbox"/> 2	Dryland <input checked="" type="checkbox"/> Riparian <input type="checkbox"/> Aquatic <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/> Perennial <input type="checkbox"/> Annual <input checked="" type="checkbox"/>	Tree <input type="checkbox"/> Shrub <input type="checkbox"/> Herb <input checked="" type="checkbox"/> Rush/Sedge <input type="checkbox"/> Grass <input type="checkbox"/> Climber <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind				
<b>Seeding Time:</b>	April - July				
<b>Method of Spread:</b>	Spreads mostly from seed				
<b>Best Time of Control:</b>	Spring / summer				
<b>Method of Control:</b>	Hand weeding is effective for this group of plants. Vigilance is required to ensure removal prior to seeding.				

Some people have adverse reactions to the sap and prickles of these plants. Care should be taken to minimise contact with bare skin and eyes.

<b>Species Name:</b>	<b>Chenopodium album</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Goosefoot	<input type="checkbox"/> 3	Dryland <input type="checkbox"/> Riparian <input checked="" type="checkbox"/> Aquatic <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/> Perennial <input checked="" type="checkbox"/> Annual <input type="checkbox"/>	Tree <input type="checkbox"/> Shrub <input type="checkbox"/> Herb <input checked="" type="checkbox"/> Rush/Sedge <input type="checkbox"/> Grass <input type="checkbox"/> Climber <input type="checkbox"/>
<b>Seed Form:</b>	Heavy seed				
<b>Seeding Time:</b>	April - June and Sept - Oct				
<b>Method of Spread:</b>	Spreads mostly from seed				
<b>Best Time of Control:</b>	All year.				
<b>Method of Control:</b>	Hand weeding is easy and effective prior to seeding.				

Make sure that this species is correctly identified as *Chenopodium glaucum* is a similar native species.

<b>Species Name:</b>	<b>Conyza spp</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Fleabane	<input type="checkbox"/> 3	Dryland <input checked="" type="checkbox"/> Riparian <input type="checkbox"/> Aquatic <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/> Perennial <input type="checkbox"/> Annual <input checked="" type="checkbox"/>	Tree <input type="checkbox"/> Shrub <input type="checkbox"/> Herb <input checked="" type="checkbox"/> Rush/Sedge <input type="checkbox"/> Grass <input type="checkbox"/> Climber <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind				
<b>Seeding Time:</b>	April - Dec and July - Feb				
<b>Method of Spread:</b>	Spreads mostly from seed				
<b>Best Time of Control:</b>	Oct - Mar				
<b>Method of Control:</b>	Hand weeding is effective prior to seeding. Needs to be ongoing. Ensure any seed heads present are bagged prior to removal if hand weeding has not occurred prior to this time.				

Common on roadsides and disturbed areas as a primary coloniser. This species is tolerant of salt, wind and is adaptable to variable soil types and therefore represents a long term problem. It is easy to control and a difference can easily be seen when controlled in bushland communities.

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



**Species Name:** *Cortaderia selloana*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Pampas Grass       1      *Dryland*       *Bulb/Corm*       *Tree*

**Seed Form:** Light and easily spread by wind      *Riparian*       *Perennial*       *Shrub*

**Seeding Time:** Dec - Feb      *Aquatic*       *Annual*       *Herb*

**Method of Spread:** Spreads mostly from seed      *Rush/Sedge*

**Best Time of Control:** Sept - Nov      *Grass*

**Method of Control:** Cut plumes before seed ripens to limit spread. Remove most leaf material with a heavy duty brushcutter and paint regrowth with Glyphosate 1 in 2. Thoroughly wet both sides of the leaf.

In riparian situations do not attempt to dig out these plants, due to the potential to affect bank stability. Should fire occur in a riparian zone, then treat the plants as soon as they reshoot to take advantage of easy access.

**Species Name:** *Cynodon dactylon*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Couch       1      *Dryland*       *Bulb/Corm*       *Tree*

**Seed Form:** Light seed      *Riparian*       *Perennial*       *Shrub*

**Seeding Time:** May, April      *Aquatic*       *Annual*       *Herb*

**Method of Spread:** Spreads readily from rhizome growth      *Rush/Sedge*

**Best Time of Control:** Oct - Feb and April - May      *Grass*

**Method of Control:** Hand weeding is very difficult, labour intensive and rarely effective. The most effective method is to spot/blanket spray in late spring - autumn using Fusillade or Targa at 4l per ha. Brushcutting and raking off bulk of plant material prior to treatment often improves ease of removal and spraying.

Do not spray over winter as this plant does not actively grow at this time. Flauzifop-butyl can be used on couch occurring amongst native rushes and sedges as they are tolerant of this chemical. Ensure that the population requiring treatment is not *Sporobolus virginicus*, the native salt water couch.

**Species Name:** *Cyperus spp*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:**       2      *Dryland*       *Bulb/Corm*       *Tree*

**Seed Form:** Light seed      *Riparian*       *Perennial*       *Shrub*

**Seeding Time:** May - July Oct - Jan      *Aquatic*       *Annual*       *Herb*

**Method of Spread:** Spreads readily from rhizome growth and seed      *Rush/Sedge*

**Best Time of Control:** Nov - Jan      *Grass*

**Method of Control:** Spot spraying in summer using 150ml of Roundup in 15l of water + Pulse. Note, Blactive is more acceptable than other forms of Glyphosate for use over waterlogged areas. Repeated brushcutting to prevent flowering is also effective in the long term.

Identification is frequently difficult with these species so it is important to ensure that the plant to be controlled is a weed and not native to the area. Remove seed heads as a minimum control technique until such time as identification has been achieved.

*Control priority 1 - Major environmental weed, urgent control required*

*Control priority 2 - Nuisance weed, control as soon as possible*

*Control priority 3 - Minor weed, control as resources become available*

**Ecosystem Management Services 1999**



<b>Species Name:</b>	<b><i>Cytisus proliferus</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Tree lucerne	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input checked="" type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	All year				Grass <input type="checkbox"/>
<b>Method of Control:</b>	The most effective method is to cut the plant off at ground level. Treating the stump with chemical is not usually necessary, unless the stump is cut more than 20mm above ground level. Remove all plant material from the site.				
	Kings Park recommends using Glyphosate at 1:15 on the cut stump.				

<b>Species Name:</b>	<b><i>Dipogon lignosus</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Dolichos pea	2	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>			Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand removal of small populations. Spot spraying with Glyphosate 1 in 50 or 1:100, can be effective.				
	At the moment, this plant is not extensively distributed around the waterways in the Perth Metropolitan area. It does have the potential however, to become a serious weed in this region - so works should focus where this species is present.				

<b>Species Name:</b>	<b><i>Echinochloa telmatophila</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Barnyard grass	2	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Oct - Dec		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	July - Sept				Grass <input checked="" type="checkbox"/>
<b>Method of Control:</b>	Remove small populations by hand. Hand weeding is preferred provided it will not increase erosion potential of any areas. As this plant occurs in wetlands, herbicide use is not preferred.				
	Alternatively treat with Fusillade or equivalent prior to flowering. Herbicide rates of 750ml to 2l dependent on plant size - prior to flowering.				

<b>Species Name:</b>	<b><i>Echium plantagineum</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Paterson's curse	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Nov - Jan		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	July - Oct				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weed small populations. Broader scale populations can be sprayed with Glyphosate. A rate of 75-100 ml per 15l of water is recommended by Kings Park Board staff.				

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**



<b>Species Name:</b>	<b><i>Ehrharta calycina</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Veldtgrass	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	March, April and Sept, Oct		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Aug - Dec				Grass <input checked="" type="checkbox"/>
<b>Method of Control:</b>	Hand weed localised infestations. Repeated brushcutting of larger stands of the weed, close to root base has been effective, followed by spot/blanket spraying using Fusillade at 4l per ha or Sertin/Targa. It is important to tag any native plants persisting amongst stands of Veldtgrass to protect them from brushcutting activities. Hand weed grasses close to any native plants.				

This plant represents a significant fire hazard in dense, extensive populations which generally occurs along disturbed road verges and fire access tracks.

<b>Species Name:</b>	<b><i>Eragrostis curvula</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	African love grass	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	June - Nov		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Nov - March				Grass <input checked="" type="checkbox"/>
<b>Method of Control:</b>	Hand weed small infestations prior to mulching. Kings Park have found complete foliar spraying after fire or in summer months using Glyphosate 1l in 100l water and wetter e.g. Agral 60, X77 to be effective. Repeated brushcutting can be effective combined with herbicide treatment of regrowth. This minimises herbicide required by a reducing the amount of leaf material.				

This plant represents a significant fire hazard and therefore a major threat to native vegetation. Do not set fire to on purpose but take advantage of easier access should any wildfire occur over summer.

<b>Species Name:</b>	<b><i>Erodium moschatum</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Musky crowfoot	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	June - Sept				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weeding is effective in predominantly native vegetation zones. This species is difficult to control due to the widespread nature of the populations.				

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



<b>Species Name:</b>	<b><i>Erythrina x sykesii</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Coral Tree	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input checked="" type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads from suckers				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Sept - Mar				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Inject tree with systemic herbicide at 10 - 15 cm intervals around the trunk. Treatment may be required several times. Cut and paint any suckers with Glyphosate.				
	Remove any branches which fall from the tree, as these can take root. Ensure bank stability is not threatened when removing the dead trunk.				

<b>Species Name:</b>	<b><i>Ferraria crispa</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Black flag	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Heavy seed		Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Nov - Dec		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads by bulb or corm growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Aug - Oct				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weed using gloves as this species is highly toxic. Kings Park suggests spot spraying Glyphosate 1 in 100 for control or using Ally/Brushoff and Glean at flowering time.				

<b>Species Name:</b>	<b><i>Ficus spp.</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Edible fig tree	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input checked="" type="checkbox"/>
<b>Seed Form:</b>	Heavy seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Dec - Mar		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Sept - Nov				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Small plants can be removed by hand. Mature plants can be injected with full strength Glyphosate at 15 cm intervals around the trunk. Fruit removal effectively reduces the rate of spread of this weed.				
	These plants are common in riparian zones. It is important not to disturb their root structure as generally these plants provide considerable bank stability in the absence of native plants. Removing the bulk of the branches and stems in dense areas may be appropriate.				

<b>Species Name:</b>	<b><i>Foeniculum vulgare</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Fennel	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Dec - Feb		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Aug - Sept				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weeding is effective for small plants. For large plants, cut the stems below ground and remove plant material prior to fruiting to reduce future spread. Alternatively, this weed can be controlled by applying Glyphosate 1 in 100 before or at flowering or repeated brushcutting.				

Control priority 1 - Major environmental weed, urgent control required  
 Control priority 2 - Nuisance weed, control as soon as possible  
 Control priority 3 - Minor weed, control as resources become available

Ecosystem Management Services 1999



<b>Species Name:</b>	<b><i>Freesia aff leichtlinii</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Freesia	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Oct - Nov		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads by bulb or corm growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Aug - Sept				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Small infestations can be dug out, bagged and removed from site. The sieving method outlined for Watsonia can be effective. Care needs to be taken to ensure that no corms are dropped when removing the plants from site - otherwise it will create more work in the future.				

For large infestations Kings Park Board Staff recommend applying Glyphosate 1 in 100 or Brushoff 5g per ha just prior to flowering (August).

<b>Species Name:</b>	<b><i>Fumaria capreolata</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Whiteflower fumitory	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Dec - Mar		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	May - Sept				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weed prior to seeding.				

<b>Species Name:</b>	<b><i>Gladiolus spp</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Gladiolus	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Feb-June		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads by bulb/corm growth and seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Aug - Dec				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Remove flower heads to prevent seed production. In heavy soils, handweed by digging around clump, sieving and shaking back sand. Can hand weed easily in dryland areas (Aug-Sept). Bag all the corms and dispose of carefully. It is possible to use herbicide for severe infestations including Glean, Brushoff and Glyphosate - using hand wiping technique.				

<b>Species Name:</b>	<b><i>Gomphocarpus fruticosus</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Cotton bush	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light and easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Nov - Dec		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Sept - Dec				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weed small plants prior to fruiting. Alternatively cut at or slightly below ground level and remove plant material. Selectively spraying the leaves with Glyphosate 1 in 100 is the suggested herbicide treatment.				

Some people have adverse reactions to the sap of this plant. Wear gloves and take care when handling plant material.

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



<b>Species Name:</b>	<b><i>Hesperanthena falcata</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>		<input type="checkbox"/> 1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads by bulb or corm growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Kings Park Board staff have been unable to find little information about controlling this weed. This agency recommends using Glyphosate at a rate of 1 to 100 at flowering time, but because this plant has small leaves it is difficult to target. Trialling Glean/Brushoff is also recommended.				

<b>Species Name:</b>	<b><i>Homeria flaccida</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	One leaf cape tulip	<input type="checkbox"/> 1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>			Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads by bulb or corm growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Removing these plants by hand can be effective if care is taken to remove all corms. For extensive populations, it is recommended that the plants are wiped with Glyphosate 1 in 10.  It is important to note that not all corms re-shoot in a given year so it is essential to monitor and treat re-growth annually. This plant is toxic to stock.				

<b>Species Name:</b>	<b><i>Hordeum leporinum</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Barley grass	<input type="checkbox"/> 3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Sept - Oct		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	July - August				Grass <input checked="" type="checkbox"/>
<b>Method of Control:</b>	Hand weeding is effective for small populations. Herbicide treatment using Fusillade at 2l per ha can work in bushland environments. Kings Park recommends spraying in July-Aug. It is important that hand weeding or spraying occurs before seed set.				

<b>Species Name:</b>	<b><i>Hyparrhenia hirta</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Tambookie grass	<input type="checkbox"/> 1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Nov - Mar				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weeding small plants prior to flowering is relatively easy. Brushcutting to remove most leaf material prior to herbicide treatment improves the effectiveness of the application. Fusillade at 4l per ha works best on new growth. Repeat treatments are likely to be required.  This is a WA native grass which is extending its distribution as a result of disturbance and vehicle movement.				

Control priority 1 - Major environmental weed, urgent control required

Control priority 2 - Nuisance weed, control as soon as possible

Control priority 3 - Minor weed, control as resources become available

Ecosystem Management Services 1999



<b>Species Name:</b>	<b><i>Hypochaeris radicata</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Flatweed	3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light and easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Oct - Mar		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	All year				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weeding is fast and effective prior to, or during flowering.				Climber <input type="checkbox"/>

<b>Species Name:</b>	<b><i>Ipomoea spp</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Morning glory	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>			Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Cut and remove existing growth, and then treat regrowth as it develops with Glyphosate at 300ml per 15l water with Pulse. This technique is preferred by the Kings Park Board staff.				
	Continued effort to remove the bulk of the vegetative material, taking care not to drop segments, can also be helpful in minimising the need for herbicide use.				
	This plant is becoming increasingly dominant in highly urbanised streams and should be controlled.				

<b>Species Name:</b>	<b><i>Isolepis proliifera</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Budding club rush	2	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Dec - Feb		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input checked="" type="checkbox"/>
<b>Best Time of Control:</b>	Winter				Grass <input type="checkbox"/>
<b>Method of Control:</b>	This plant occurs in homogeneous clumps in seasonally waterlogged area. It may be worth trying to cover this weed with black plastic held down with rocks to drown the plant over winter.				
	Rotary hoeing and spraying the regrowth with Glyphosate with surfactant can be effective. Kings Park Board suggests Glyphosate 1 to 20 plus Pulse. It is important to do this in summer following the frog breeding season and prior to the bird breeding season. Repeat treatments will be required.				

Control priority 1 - Major environmental weed, urgent control required  
 Control priority 2 - Nuisance weed, control as soon as possible  
 Control priority 3 - Minor weed, control as resources become available

Ecosystem Management Services 1999





<b>Species Name:</b>	<b><i>Juncus articulatus</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Articulated rush	<input type="checkbox"/> 2	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Nov - Mar		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input checked="" type="checkbox"/>
<b>Best Time of Control:</b>	Sept - Mar				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Manually weeding all plants is the preferred method for removing this species.				Climber <input type="checkbox"/>

Ensure that the plants to be controlled have been correctly identified as the weed species. If unsure of weed status then removing the flowering heads to minimise spread is helpful and will not seriously interfere with the plants until they have been correctly identified.

<b>Species Name:</b>	<b><i>Juncus capitatus</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>		<input type="checkbox"/> 3	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Dec - mar		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input checked="" type="checkbox"/>
<b>Best Time of Control:</b>	Sept - Nov				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Manually weed small plants. The preferred method for removing larger clumps involves brushcutting to remove the bulk of material and then digging the plants out and removing the base and leaves from the site. Any regrowth from sections missed can then be slashed and treated with Glyphosate applied at half strength. Several applications may be required.				Climber <input type="checkbox"/>

Ensure that the plants to be controlled have been correctly identified as weed species. If unsure of weed status then removing the flowering heads to minimise spread is helpful and will not seriously interfere with the plants until they have been correctly identified.

<b>Species Name:</b>	<b><i>Juncus microcephalus</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>		<input type="checkbox"/> 2	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Dec - Mar		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input checked="" type="checkbox"/>
<b>Best Time of Control:</b>	Sept - Dec				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Manually weed small plants. The preferred method for removing larger clumps involves brushcutting to remove the bulk of material and then digging the plants out and removing the base and leaves from the site. Any regrowth from sections missed can then be slashed and treated with Glyphosate applied at half strength. Several applications may be required.				Climber <input type="checkbox"/>

This plant is a serious weed. Ensure correct identification prior to implementing weed control as this plant is similar to native rush and sedge species. Plants occurring on river banks should not be dug out as removal may create a new erosion problem. Use extra care when using herbicides close to the water.

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



<b>Species Name:</b>	<b>Lantana camara</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Lantana	<input type="checkbox"/> 3	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>			Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weed (grub out) small communities. Spray localised populations with Glyphosate 1 in 10 covering all foliage.				Climber <input checked="" type="checkbox"/>
	Monitoring re-occurrence of this plant in areas where previous control work has been undertaken is essential.				

<b>Species Name:</b>	<b>Leptospermum laevigatum</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Victorian coastal teatree	<input type="checkbox"/> 1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input checked="" type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	April - October		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	All year				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Hand weed seedlings. For mature plants, cut stems to ground level annually until control is achieved. Remove flowering branches when possible.				Climber <input type="checkbox"/>
	Note, in some cases where this weed provides shelter this should be done only after native plants have grown sufficiently to take their place.				

<b>Species Name:</b>	<b>Lolium spp.</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Rye grass	<input type="checkbox"/> 2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	March - June		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Dec - Mar				Grass <input checked="" type="checkbox"/>
<b>Method of Control:</b>	Handweeding is preferred, except for extensive populations. Spot spraying of Sertin, Targa or similar at 4l per ha prior to flowering can be effective.				Climber <input type="checkbox"/>
	In areas where steep banks are present and this species is dominant removing the seed heads to limit spread is preferred to complete removal, in order to ensure that bank stability is protected.				

<b>Species Name:</b>	<b>Lupinus angustifolia</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Lupin	<input type="checkbox"/> 2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Heavy seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Oct - Dec		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Aug - Oct				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Handweed small populations. Alternatively, spray the plants selectively with Glyphosate 2% solution.				Climber <input type="checkbox"/>

Control priority 1 - Major environmental weed, urgent control required

Control priority 2 - Nuisance weed, control as soon as possible

Control priority 3 - Minor weed, control as resources become available

Ecosystem Management Services 1999



<b>Species Name:</b>	<b><i>Medicago spp</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Medics	3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	June - Sept				Grass <input type="checkbox"/>
<b>Method of Control:</b>	This plant may be controlled effectively with Glyphosate. Kings Park Board recommends a rate of 75-100ml in 15l of water.				

<b>Species Name:</b>	<b><i>Monopsis debilis</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>		3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>			Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>					Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Pull out small populations to prevent them from spreading. Repeated rotary hoeing/mowing to prevent flowering can be helpful.				
	Kings Park Board staff suggest Glyphosate at 75-100ml in 15l of water prior to flowering.				

<b>Species Name:</b>	<b><i>Myrsiphyllum asparagoides</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Bridal Creeper	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>	Oct - Dec		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Jul - Sept				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Remove young plants by hand as they appear. If spraying, remove the bulk of the plant material prior to spraying then treat the smaller biomass of plants approximately a fortnight later. Kings Park currently recommends using either Glyphosate 360 at a rate of 1 in 100, or 2.5 to 5g per ha in 250l of water. Repeat applications will be required for either chemical.				
	Kings Park may have more up to date control measures. It is essential to take extreme care when treating this plant as it generally occurs within close proximity of native plants, and causing the unintentional death of non-target plants is possible.				

<b>Species Name:</b>	<b><i>Narcissus tazetta</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Jonquil	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads by bulb or corm growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Winter - Spring				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Removing these plants by hand can be effective if care is taken to remove all corms. For extensive populations, it is recommended that the plants are wiped with Glyphosate 1 in 10.				
	It is important to note that not all corms re-shoot in a given year so it is essential to monitor and treat re-growth annually. This plant is toxic to stock.				

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



**Species Name:** *Nerium oleander*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Oleander       3      Dryland       Bulb/Corm       Tree

**Seed Form:** Coarse seed      Riparian       Perennial       Shrub

**Seeding Time:**      Aquatic       Annual       Herb

**Method of Spread:** Spreads from both seed and vegetative growth      Rush/Sedge

**Best Time of Control:** All year      Grass

**Method of Control:** Dig out the individual plants. Otherwise cut the stumps and paint with full strength systemic herbicide.      Climber

**Species Name:** *Olea europaea*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Olive tree       2      Dryland       Bulb/Corm       Tree

**Seed Form:** Heavy seed      Riparian       Perennial       Shrub

**Seeding Time:** Nov - Jan      Aquatic       Annual       Herb

**Method of Spread:** Spreads mostly from seed      Rush/Sedge

**Best Time of Control:**      Grass

**Method of Control:** Hand weed juvenile plants. For small plants, selectively spray foliage with full strength Glyphosate. Larger trees can be managed by either cutting the stump and painting with Glyphosate or Garlon (recommended by Kings Park Board staff), or alternatively injecting into the stem at 15 cm intervals. Follow up treatments may be required.

Encouraging fruit harvesting by residents will reduce the rate of spread of this weed.

**Species Name:** *Oxalis pes-caprae*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Soursob       2      Dryland       Bulb/Corm       Tree

**Seed Form:** Light seed      Riparian       Perennial       Shrub

**Seeding Time:** Sept      Aquatic       Annual       Herb

**Method of Spread:** Spreads by runners      Rush/Sedge

**Best Time of Control:** July - Sept      Grass

**Method of Control:** Hand weeding can be effective provided that care is taken to trace all runners from the parent plant and that no stem and root is left behind.

Apply Glyphosate 75ml in 10l in winter or before foliage starts to yellow.

**Species Name:** *Panicum capillare*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Witchgrass       3      Dryland       Bulb/Corm       Tree

**Seed Form:**      Riparian       Perennial       Shrub

**Seeding Time:**      Aquatic       Annual       Herb

**Method of Spread:** Spreads mostly from seed      Rush/Sedge

**Best Time of Control:**      Grass

**Method of Control:** As with most introduced grasses, Fusillade at 2l per ha can be effective. The herbicide should be applied prior to flowering.

This species has the potential to spread rapidly through wetland environments.

Control priority 1 - Major environmental weed, urgent control required  
 Control priority 2 - Nuisance weed, control as soon as possible  
 Control priority 3 - Minor weed, control as resources become available

Ecosystem Management Services 1999



<b>Species Name:</b>	<b><i>Paspalum spp</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Paspalum	<input type="checkbox"/> 2	Dryland <input checked="" type="checkbox"/> Riparian <input checked="" type="checkbox"/> Aquatic <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/> Perennial <input checked="" type="checkbox"/> Annual <input type="checkbox"/>	Tree <input type="checkbox"/> Shrub <input type="checkbox"/> Herb <input type="checkbox"/> Rush/Sedge <input type="checkbox"/> Grass <input checked="" type="checkbox"/> Climber <input type="checkbox"/>
<b>Seed Form:</b>	Heavy seed				
<b>Seeding Time:</b>	Dec - Jan				
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				
<b>Best Time of Control:</b>	Aug - Mar				
<b>Method of Control:</b>	Repeated brushcutting/slashing can be effective in controlling this plant - provided it occurs prior to seed development. The accepted herbicide treatment is the application of Fusillade at 4l per ha.				
	It is possible to reduce the volume of herbicide required by slashing/rotary hoeing and then treating the regrowth.				

<b>Species Name:</b>	<b><i>Pelargonium capitatum</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Rose pelargonium	<input type="checkbox"/> 1	Dryland <input checked="" type="checkbox"/> Riparian <input type="checkbox"/> Aquatic <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/> Perennial <input checked="" type="checkbox"/> Annual <input type="checkbox"/>	Tree <input type="checkbox"/> Shrub <input type="checkbox"/> Herb <input checked="" type="checkbox"/> Rush/Sedge <input type="checkbox"/> Grass <input type="checkbox"/> Climber <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind				
<b>Seeding Time:</b>	Jan - April				
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				
<b>Best Time of Control:</b>	Spring				
<b>Method of Control:</b>	Hand weed in autumn / winter, trying very hard not leave any stem or root behind as the plants will reshoot. Kings Park suggests the two herbicide treatments listed. Spot Spray with Alty/Brush 5g per ha or spray with Glyphosate 1 in 100 with wetting agent in early September.				
	This plant is an effective coloniser and it may smother any small native plants present.				

<b>Species Name:</b>	<b><i>Pennisetum clandestinum</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Kikuyu	<input type="checkbox"/> 1	Dryland <input checked="" type="checkbox"/> Riparian <input checked="" type="checkbox"/> Aquatic <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/> Perennial <input checked="" type="checkbox"/> Annual <input type="checkbox"/>	Tree <input type="checkbox"/> Shrub <input type="checkbox"/> Herb <input type="checkbox"/> Rush/Sedge <input type="checkbox"/> Grass <input checked="" type="checkbox"/> Climber <input type="checkbox"/>
<b>Seed Form:</b>	Sterile or non seed producing				
<b>Seeding Time:</b>					
<b>Method of Spread:</b>	Spreads readily from rhizome growth				
<b>Best Time of Control:</b>	Sept - Dec				
<b>Method of Control:</b>	The most effective technique recognised is the application of Fusillade at a rate of 4l per ha while the plant is actively growing.				
	Fusillade should not be applied over open water. Native rushes and sedges are not at risk when using this chemical.				

<b>Species Name:</b>	<b><i>Plantago lanceolata</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Ribwort plantain	<input type="checkbox"/> 3	Dryland <input checked="" type="checkbox"/> Riparian <input type="checkbox"/> Aquatic <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/> Perennial <input checked="" type="checkbox"/> Annual <input type="checkbox"/>	Tree <input type="checkbox"/> Shrub <input type="checkbox"/> Herb <input checked="" type="checkbox"/> Rush/Sedge <input type="checkbox"/> Grass <input type="checkbox"/> Climber <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed				
<b>Seeding Time:</b>					
<b>Method of Spread:</b>	Spreads mostly from seed				
<b>Best Time of Control:</b>	Nov - Dec				
<b>Method of Control:</b>	Pull Ribwort by hand ensuring that tap root is properly removed. Generally populations of this weed are limited and can be managed effectively using manual weed control methods. Kings Park Board recommends wiping with Glyphosate 100ml in 15l water.				

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



**Species Name:** *Populus spp* **Control Priority** 2 **Location** **Habit** **Form**

**Common Name:** Poplar  Dryland  Bulb/Corm  Tree   
 Riparian  Perennial  Shrub   
 Aquatic  Annual  Herb   
Rush/Sedge   
Grass   
Climber

**Seed Form:**

**Seeding Time:**

**Method of Spread:** Spreads from suckers

**Best Time of Control:** Oct - Feb

**Method of Control:** Experience indicates that injecting concentrated systemic herbicide at 10 - 15 cm intervals around the trunk can be effective, and reduces the number of suckers which can occur following the cut stump technique. Kings Park considers this plant difficult to control and recommends the cut stump method with Garlon 600.

**Species Name:** *Raphanus raphanistrum* **Control Priority** 3 **Location** **Habit** **Form**

**Common Name:** Wild radish  Dryland  Bulb/Corm  Tree   
 Riparian  Perennial  Shrub   
 Aquatic  Annual  Herb   
Rush/Sedge   
Grass   
Climber

**Seed Form:** Light seed

**Seeding Time:** Dec

**Method of Spread:** Spreads mostly from seed

**Best Time of Control:** Sept - Nov

**Method of Control:** Removing these species by hand is easy and can be done very quickly. Removal should occur prior to the plants flowering and seeding to reduce the rate of spread. Bagging and cutting the seeding stems, from any plants, should be undertaken prior to removal.

The alternative is to paint with Glyphosate 1 in 10.

**Species Name:** *Rhynchosyris repens* **Control Priority** 1 **Location** **Habit** **Form**

**Common Name:** Red natal grass  Dryland  Bulb/Corm  Tree   
 Riparian  Perennial  Shrub   
 Aquatic  Annual  Herb   
Rush/Sedge   
Grass   
Climber

**Seed Form:** Light and easily spread by wind

**Seeding Time:** Sept - Nov

**Method of Spread:** Spreads mostly from seed

**Best Time of Control:** June to Aug

**Method of Control:** This plant is effectively controlled using Fusillade at a rate of 4l per ha (as for most other introduced grasses).

**Species Name:** *Ricinus communis* **Control Priority** 1 **Location** **Habit** **Form**

**Common Name:** Castor Oil  Dryland  Bulb/Corm  Tree   
 Riparian  Perennial  Shrub   
 Aquatic  Annual  Herb   
Rush/Sedge   
Grass   
Climber

**Seed Form:** Heavy seed

**Seeding Time:** Nov - Jan

**Method of Spread:** Spreads mostly from seed

**Best Time of Control:** Any time but best prior to fruiting

**Method of Control:** Small populations can be removed by hand. Individual plants can be cut and painted with Glyphosate. Populations of seedlings can be sprayed with Glyphosate 1 in 80, while injecting large plants with a systemic herbicide is effective.

The seed from this plant has been shown to be viable more than 1 000 years later, so vigilance is required to remove plants prior to seeding.

*Control priority 1 - Major environmental weed, urgent control required*  
*Control priority 2 - Nuisance weed, control as soon as possible*  
*Control priority 3 - Minor weed, control as resources become available*

**Ecosystem Management Services 1999**



<b>Species Name:</b>	<b>Romulea rosea</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Guildford grass	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input checked="" type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads by bulb or corm growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input checked="" type="checkbox"/>
<b>Method of Control:</b>	In areas with homogeneous populations, Kings Park Board suggests Brushoff / Ally can give good control and can be used over some turf species. Repeated rotary hoeing and slashing prior to flowering can assist in managing populations.				

<b>Species Name:</b>	<b>Rorippa nasturtium-aquaticum</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Watercress	2	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input checked="" type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Access dependent				Grass <input type="checkbox"/>
<b>Method of Control:</b>	This aquatic weed is difficult to control because it slows water movement, increases sedimentation and reduces erosion which means implementing control can affect bed and bank stability. The recommended removal technique involves manual clearing of a channel and also clearing 5 to 10 m wide bands, 20 metres apart which are perpendicular to the stream flow. This will minimise the potential for de-stabilising the stream bed.				
	Seek expert advice and approvals from the relevant government agencies prior to implementing broad scale works.				

<b>Species Name:</b>	<b>Rubus spp</b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Blackberry	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Heavy seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input checked="" type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	Dec - April				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Brush cut and remove brambles. Hand weed removing knotty stumps and as much root as possible. Paint regrowth with Glyphosate 12ml to 1l of water. Better control is often achieved with a combination of Brushoff, Garlon or blackberry and tree killer. Biological controls using a rust fungus have been successful. Agriculture WA may be able to assist with this.				
	Brushcutting these plants can provide very difficult and using a team of goats as the first method of attack can prove very useful in terms of increasing access and removing the bulk of the vegetative material. It is important that any blackberry control takes into consideration fauna corridors in continuous strips of sufficient width to discourage predators, particularly to protect birds and bandicoots.				

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



<i>Species Name:</i>	<i>Rumex spp</i>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Dock	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<i>Seed Form:</i>	Light and easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>	March - June		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<i>Method of Spread:</i>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>	Nov - Mar				Grass <input type="checkbox"/>
<i>Method of Control:</i>	These plants are readily eradicated through hand weeding. Remove flowering heads prior to seed ripening if complete plant removal is not possible.				
	Always bag plants with seeds and dispose of carefully.				

<i>Species Name:</i>	<i>Salix spp</i>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Willow	1	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input checked="" type="checkbox"/>
<i>Seed Form:</i>	Heavy seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<i>Method of Spread:</i>	Spreads from suckers				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>	Dec - Mar				Grass <input type="checkbox"/>
<i>Method of Control:</i>	Small plants can be removed by hand. Mature plants can be injected with full strength Glyphosate at 10 - 15 cm intervals around the trunk. Any suckers which appear can be painted with systemic herbicide. It is important not to remove the parent plant until it is dead and no more suckers are being produced.				
	Removal of willows along watercourses can have a detrimental effect through loss of habitat, streamside erosion and exposure of understorey. Consideration should be given to replacing the plants to be removed two years prior to undertaking removal.				

<i>Species Name:</i>	<i>Schinus terebinthifolia</i>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Japanese pepper	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input checked="" type="checkbox"/>
<i>Seed Form:</i>	Coarse seed		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>	Sept		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<i>Method of Spread:</i>	Spreads from suckers and seed				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>	All year, but in wetlands treat in summer				Grass <input type="checkbox"/>
<i>Method of Control:</i>	Hand weed small seedlings. It is important to monitor for any new germinants to enable rapid removal from the site. Treating the large plants can be undertaken either by cutting the trunk and immediately painting the stump, or alternatively injecting systemic herbicide at 10 - 15 cm intervals around the trunk. Kings Park recommends either Glyphosate, Velpar or Garlon.				
	The seed is spread predominantly by introduced birds and there is some anecdotal evidence that many native birds are poisoned by the seeds.				

*Control priority 1 - Major environmental weed, urgent control required*

*Control priority 2 - Nuisance weed, control as soon as possible*

*Control priority 3 - Minor weed, control as resources become available*

*Ecosystem Management Services 1999*





<i>Species Name:</i>	<b><i>Solenum nigrum</i></b>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Deadly nightshade	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<i>Seed Form:</i>	Coarse seed		Riparian <input checked="" type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>	Oct - Dec		Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<i>Method of Spread:</i>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>	Sept - Oct				Grass <input type="checkbox"/>
<i>Method of Control:</i>	Hand weed small infestations. Kings Park Board recommends using Glyphosate 1 in 100. Dessicant herbicides applied to all parts of the plant can be effective on warm to hot days.				

<i>Species Name:</i>	<b><i>Stachys arvensis</i></b>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Staggerweed	3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<i>Seed Form:</i>	Heavy seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<i>Method of Spread:</i>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>					Grass <input type="checkbox"/>
<i>Method of Control:</i>	Pull out small populations to prevent them from spreading. Repeated rotary hoeing/mowing to prevent flowering can be helpful where there are no remnant native species. Kings Park Board staff suggest Glyphosate at 75-100ml in 15l of water prior to flowering.				

<i>Species Name:</i>	<b><i>Stenotaphrum secundatum</i></b>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Buffalo grass	1	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<i>Seed Form:</i>	Sterile or non seed producing		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<i>Method of Spread:</i>	Spreads readily from rhizome growth				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>	Aug - Sept				Grass <input checked="" type="checkbox"/>
<i>Method of Control:</i>	Hand weeding is very difficult, labour intensive and rarely successful. The most effective method is to implement a minimum of two spot/blanket treatments in Aug-Oct and April-May using Fusillade or Targa at 4l per ha. Brushcutting often improves ease of removal and spraying. This process typically requires more than two treatments. Can implement spraying amongst native rushes and sedges which have been demonstrated to tolerate flauzifop-butyl.				

<i>Species Name:</i>	<b><i>Taraxacum officinale</i></b>	<i>Control Priority</i>	<i>Location</i>	<i>Habit</i>	<i>Form</i>
<i>Common Name:</i>	Dandelion	2	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<i>Seed Form:</i>	Light, easily spread by wind		Riparian <input type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<i>Seeding Time:</i>	All year round		Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<i>Method of Spread:</i>	Spreads mostly from seed				Rush/Sedge <input type="checkbox"/>
<i>Best Time of Control:</i>	Sept - Nov				Grass <input type="checkbox"/>
<i>Method of Control:</i>	Hand weeding is the most effective means of control ensuring that if seed heads are present , they are carefully bagged prior to removal of the plant. Wiping with Glyphosate is also effective.				

*Control priority 1 - Major environmental weed, urgent control required*

*Control priority 2 - Nuisance weed, control as soon as possible*

*Control priority 3 - Minor weed, control as resources become available*

**Ecosystem Management Services 1999**



**Species Name:** *Thunbergia alata*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Black-eyed Susan       2      Dryland       Bulb/Corm       Tree

**Seed Form:** Coarse seed      Riparian       Perennial       Shrub

**Seeding Time:**      Aquatic       Annual       Herb

**Method of Spread:** Spreads from both seed and vegetative growth      Rush/Sedge

**Best Time of Control:**      Grass

**Method of Control:** Remove small plants manually. Spot spraying with Glyphosate at a rate of 1 in 50 can be effective.      Climber

This plant poses a serious threat to the State's waterways and any small populations should be worked on quickly to reduce the potential spread.

**Species Name:** *Trifolium spp.*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Clovers       3      Dryland       Bulb/Corm       Tree

**Seed Form:** Heavy seed      Riparian       Perennial       Shrub

**Seeding Time:**      Aquatic       Annual       Herb

**Method of Spread:** Spreads mostly from seed      Rush/Sedge

**Best Time of Control:**      Grass

**Method of Control:** Hand weed small populations. Spraying populations with Glyphosate at 75 - 100 ml in 15l of water is recommended by Kings Park Board. Repeated rotary hoeing with follow up spraying can be effective in pasture situations.      Climber

**Species Name:** *Tropaeolum majus*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Nasturtium       3      Dryland       Bulb/Corm       Tree

**Seed Form:** Heavy seed      Riparian       Perennial       Shrub

**Seeding Time:** Nov - Jan      Aquatic       Annual       Herb

**Method of Spread:** Spreads mostly from seed      Rush/Sedge

**Best Time of Control:** Aug / Sept      Grass

**Method of Control:** Removing this species by hand is effective. Selectively applying Glyphosate 1 in 100 can be effective.      Climber

Awareness campaigns about the implications of dumping garden waste in reserves need to be upgraded and implemented intensively to discourage such activities.

Control priority 1 - Major environmental weed, urgent control required  
 Control priority 2 - Nuisance weed, control as soon as possible  
 Control priority 3 - Minor weed, control as resources become available

Ecosystem Management Services 1999



<b>Species Name:</b>	<b><i>Typha orientalis</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Bulrush	<input type="checkbox"/> 1	Dryland <input type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light, easily spread by wind		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads readily from rhizome growth and seed				Rush/Sedge <input checked="" type="checkbox"/>
<b>Best Time of Control:</b>	Winter				Grass <input type="checkbox"/>
<b>Method of Control:</b>	Remove seed heads prior to ripening in September - December. Cut stems below water level in May, if sufficient water is present, monitor regrowth and continue to cut until September to drown the plants.				Climber <input type="checkbox"/>

For populations occurring in waterlogged areas only use Glyphosate BioActive 1 to 10 in spring, after slashing plants first and wipe new growth when plants are 1m tall. Take care when using herbicide over water.

The native cumbungi, *Typha domingensis*, looks similar to Bulrush and it is important to ensure that the population being controlled is in fact the weed species.

<b>Species Name:</b>	<b><i>Ursinia anthemoides</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Ursinia	<input type="checkbox"/> 3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Light seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>					Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Pull out small populations to prevent them from spreading. Repeated rotary hoeing/mowing to prevent flowering can be helpful.				Climber <input type="checkbox"/>

Kings Park Board staff suggest Glyphosate at 75-100ml in 15l of water prior to flowering.

<b>Species Name:</b>	<b><i>Vicia sativa</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Vetch	<input type="checkbox"/> 3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Heavy seed		Riparian <input type="checkbox"/>	Perennial <input type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input checked="" type="checkbox"/>	Herb <input checked="" type="checkbox"/>
<b>Method of Spread:</b>	Spreads from both seed and vegetative growth				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>					Grass <input type="checkbox"/>
<b>Method of Control:</b>	Kings Park recommends trying Glyphosate 75ml in 15 l when the plants are actively growing. Hand weeding small populations is possible and effective.				Climber <input type="checkbox"/>

<b>Species Name:</b>	<b><i>Vinca major</i></b>	<b>Control Priority</b>	<b>Location</b>	<b>Habit</b>	<b>Form</b>
<b>Common Name:</b>	Periwinkle	<input type="checkbox"/> 3	Dryland <input checked="" type="checkbox"/>	Bulb/Corm <input type="checkbox"/>	Tree <input type="checkbox"/>
<b>Seed Form:</b>	Coarse seed		Riparian <input checked="" type="checkbox"/>	Perennial <input checked="" type="checkbox"/>	Shrub <input type="checkbox"/>
<b>Seeding Time:</b>			Aquatic <input type="checkbox"/>	Annual <input type="checkbox"/>	Herb <input type="checkbox"/>
<b>Method of Spread:</b>	Spreads by runners				Rush/Sedge <input type="checkbox"/>
<b>Best Time of Control:</b>	June - Aug				Grass <input type="checkbox"/>
<b>Method of Control:</b>	It is generally recommended that this weed is managed by applying Glyphosate at 1 in 10 with surfactant.				Climber <input checked="" type="checkbox"/>

Applications will need to be repeated several times at intervals of one month.

**Control priority 1 - Major environmental weed, urgent control required**

**Control priority 2 - Nuisance weed, control as soon as possible**

**Control priority 3 - Minor weed, control as resources become available**

**Ecosystem Management Services 1999**



**Species Name:** *Watsonia bulbiflora*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Watsonia       1      Dryland       Bulb/Corm       Tree

**Seed Form:** Light and easily spread by wind and wat      Riparian       Perennial       Shrub

**Seeding Time:** March - May      Aquatic       Annual       Herb

**Method of Spread:** Spreads by bulb/corm growth      Rush/Sedge

**Best Time of Control:**

**Method of Control:** Remove corms by carefully digging a large area around each plant, putting the sand onto flywire, sieving and collecting all the corms. Flowers should also be harvested to prevent the production of seed and subsequent spread. The collected corms should be disposed of carefully.

Broadscale removal of dense stands may threaten bank stability. Remove in nodes along the waterway.  
 Selectively spray a combination of herbicides between July to August using Glean and Ally/Brushoff and subsequently painting leaf with Glyphosate in September to November can be effective. Remove the bulk of dead biomass leaving the rhizome mats in tact.

**Species Name:** *Zantedeschia aethiopica*      **Control Priority**      **Location**      **Habit**      **Form**

**Common Name:** Arum lily       1      Dryland       Bulb/Corm       Tree

**Seed Form:** Coarse seed      Riparian       Perennial       Shrub

**Seeding Time:** Dec      Aquatic       Annual       Herb

**Method of Spread:** Spreads from both seed and vegetative growth      Rush/Sedge

**Best Time of Control:** April - Nov      Grass

**Method of Control:** Entire plants can be removed by digging - make sure to remove all of the rhizome. Spot spray from April to November using Glyphosate 1 in 100 or Glean Ally/Brushoff 1 in 50 (20g per ha). Respraying is likely to be required 8 weeks later.

In wetland environments Roundup Biactive should be used to minimise fauna losses.

*Control priority 1 - Major environmental weed, urgent control required*  
*Control priority 2 - Nuisance weed, control as soon as possible*  
*Control priority 3 - Minor weed, control as resources become available*





# Appendix 3

## Suggested species for revegetation works



### Appendix 3: Suggested species for revegetation works

Species	CommonName	Location								Habitat		
		Roley Pool	Wright Brook	Breera Brook	Bannister Creek	Bennett Brook	Ellen Brook	Southern Wood Creek	Upper Canning	Dryland	Bank	Emergent
<b><u>1. Spreading tree</u></b>												
<i>Banksia attenuata</i>	Slender banksia	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Banksia littoralis</i>	Swamp banksia	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Banksia menziesii</i>	Firewood banksia	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Casuarina obesa</i>	Saltwater sheoak	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Corymbia calophylla</i>	Marri	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Eucalyptus marginata</i>	Jarrah	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Eucalyptus rudis</i>	Flooded gum	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Eucalyptus wandoo</i>	Wandoo	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Paraserianthes lophantha</i>	Native albizia	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<b><u>2. Compact tree</u></b>												
<i>Eucalyptus todtiana</i>	Coastal blackbutt	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Melaleuca cuticularis</i>	Saltwater paperbark	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Melaleuca preissiana</i>	Modong	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Melaleuca raphiophylla</i>	Swamp paperbark	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Nuytsia floribunda</i>	Christmas tree	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<b><u>3. Large shrub</u></b>												
<i>Acacia saligna</i>	Coojong	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Agonis linearifolia</i>	Swamp peppermint	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Dryandra sessilis</i>	Parrot bush	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Grevillea diversifolia</i>	Variable leaved grevillea	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Melaleuca incana</i>	Grey honeymyrtle	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Melaleuca teretifolia</i>		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>

This information is site specific to the sections of assessed foreshore. Please seek expert advice if placing these species outside of the surveyed sections.

Copyright Ecosystem Management Services 1999

Species	CommonName	Location								Habitat		
		Roley Pool	Wright Brook	Breera Brook	Bannister Creek	Bennett Brook	Ellen Brook	Southern Wood Creek	Upper Canning	Dryland	Bank	Emergent
<i>Melaleuca viminea</i>	Mohan	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Oxylobium lineare</i>	River pea	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Viminaria juncea</i>	Swishbush	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<b>4. Medium shrub</b>												
<i>Acacia pulchella</i>	Prickly moses	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Astartea fascicularis</i>	Common Astartea	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Darwinia citriodora</i>	Lemon scented darwinia	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Hakea varia</i>	Harsh hakea	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Hibbertia spp</i>	Native buttercups	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Jacksonia furcellata</i>	Grey stinkwood	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Jacksonia sternbergiana</i>	Green stinkwood	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Kunzea ericifolia</i>	Spearwood	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Lasiopetalum bracteatum</i>	Helena Velvet Bush	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Melaleuca lateritia</i>	Robin Red-breast bush	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Melaleuca viminea</i>	Mohan	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Pericalymma ellipticum</i>	Swamp teatree	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Pteridium esculentum</i>	Bracken fern	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Regelia ciliata</i>	Regelia	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Thomasia macrocarpa</i>		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<b>5. Low shrub</b>												
<i>Acacia alata</i>	Winged wattle	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Acanthocarpus preissii</i>		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Bossiaea spp</i>		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Corynotheca micrantha</i>	Sand lily	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Gompholobium tomentosum</i>	Hairy yellow pea	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

This information is site specific to the sections of assessed foreshore. Please seek expert advice if placing these species outside of the surveyed sections.

Copyright Ecosystem Management Services 1999



Species	CommonName	Location								Habitat		
		Roley Pool	Wright Brook	Breera Brook	Bannister Creek	Bennett Brook	Ellen Brook	Southern Wood Creek	Upper Canning	Dryland	Bank	Emergent
<i>Hakea prostrata</i>	Harsh Hakea	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Hypocalymma angustifolium</i>	White myrtle	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Hypocalymma robustum</i>	Swan River myrtle	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Leucopogon spp</i>		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Macrozamia riedlei</i>	Zamia	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Verticordia spp</i>	Featherflowers	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<b><u>6. Ground cover</u></b>												
<i>Centella cordifolia</i>	Centella	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Conostylis candicans</i>	Grey cottonhead	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Cotula coronopifolia</i>	Waterbuttons	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Dryandra nivea</i>	Couch honeypots	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Hemarthria uncinata</i>	Mat grass	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Hemiandra pungens</i>	Snake bush	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Patersonia occidentalis</i>	Western iris	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Sporobolus virginicus</i>	Saltwater couch	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<b><u>7. Climber</u></b>												
<i>Clematis pubescens</i>	Common clematis	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Hardenbergia comptoniana</i>	Native wisteria	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Kennedia coccinea</i>	Coral creeper	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<i>Kennedia prostrata</i>	Running postman	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<b><u>8. Rush or Sedge</u></b>												
<i>Juncus subsecundus</i>	Finger rush	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Baumea articulata</i>	Jointed twig sedge	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Baumea juncea</i>	Bare twig rush	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Baumea preissii</i>	Broad twig sedge	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>

This information is site specific to the sections of assessed foreshore. Please seek expert advice if placing these species outside of the surveyed sections.

Copyright Ecosystem Management Services 1999

Species	CommonName	Location								Habitat		
		Roley Pool	Wright Brook	Breera Brook	Bannister Creek	ennett Brook	Ellen Brook	Southern Wood Creek	Upper Canning	Dryland	Bank	Emergent
<i>Baumea rubiginosa</i>	River twig	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Bolboschoenus caldwellii</i>	Marsh club rush	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Carex appressa</i>	Tall sedge	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Carex divisa</i>	Divided sedge	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Carex fascicularis</i>	Tassel sedge	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Carex tereticaulis</i>	Tube sedge	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Centrolepis spp</i>		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Eleocharis acuta</i>	Spike sedge	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Isolepis nodosa</i>	Knotted Club sedge	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Isolepis setiformis</i>	Tufted sedge	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Juncus holoschoenus</i>	Joint-leaf rush	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Juncus kraussii</i>	Shore rush	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Juncus pallidus</i>	Pale rush	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Juncus pauciflorus</i>	Slender rush	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Lepidosperma effusum</i>	Spreading sword sedge	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Lepidosperma longitudinale</i>	Pithy sword sedge	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Lepidosperma tetraquetrum</i>	Angle sword sedge	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
<i>Restio spp</i>		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
<i>Schoenoplectus validus</i>	Lake Club Sedge	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>

*This information is site specific to the sections of assessed foreshore. Please seek expert advice if placing these species outside of the surveyed sections.*

**Copyright Ecosystem Management Services 1999**







# Appendix 4

## Suggested soft engineering works



### Appendix 4: Suggested soft engineering works



# Appendix 5

## Condition mapping symbols





**Weeds**

Symbol	Common name	Scientific name
	Weed wattles	<i>Acacia spp.</i>
	Giant reed	<i>Arundo donax</i>
	Canna lily	<i>Canna spp.</i>
	Pampas grass	<i>Cortaderia selloana</i>
	Perennial veldtgrass	<i>Ehrharta calycina</i>
	African lovegrass	<i>Eragrostis curvula</i>
	Coral tree	<i>Erythrina x sykesii</i>
	Edible fig tree	<i>Ficus spp.</i>
	Cotton bush	<i>Gomphocarpus fruticosus</i>
	One leaf cape tulip	<i>Homeria flaccida</i>
	Morning glory	<i>Ipomoea spp.</i>
	Lantana	<i>Juncus microcephalus</i>
	Bridal creeper	<i>Lantana camara</i>
	Paspalum	<i>Myrsiphyllum asparagoides</i>
	Castor oil bush	<i>Paspalum spp.</i>
	Blackberry	<i>Ricinus communis</i>
	Willow	<i>Rubus fruticosus</i>
	Japanese pepper	<i>Salix spp.</i>
	Deadly nightshade	<i>Schinus terebinthifolia</i>
	Nasturtium	<i>Solanum nigrum</i>
	Bulrush	<i>Tropeolum spp.</i>
	Vetch	<i>Typha orientalis</i>
	Watsonia	<i>Vicia sativa</i>
	Arum lily	<i>Watsonia bulbifera</i>
		<i>Zantedeschia aethiopica</i>

**Native Species**

Symbol	Common name	Scientific name
Al	Swamp peppermint	<i>Agonis linearifolia</i>
As	Coojong	<i>Acacia saligna</i>
Ba	Slender banksia	<i>Banksia attenuata</i>
Bj	Bare twigrush	<i>Baumea juncea</i>
Ca	Tall sedge	<i>Carex appressa</i>
Cc	Marri	<i>Corymbia calophylla</i>
Er	Flooded gum	<i>Eucalyptus rudis</i>
Hc	Native wisteria	<i>Hardenbergia comptoniana</i>
Jp	Pale rush	<i>Juncus pallidus</i>
Js	Green stinkwood	<i>Jacksonia sternbergiana</i>
Kp	Running postman	<i>Kennedia prostrata</i>
Li	Pithy sword-sedge	<i>Lepidosperma longitudinale</i>
Lt	Angle sword-sedge	<i>Lepidosperma tetraquetrum</i>
Mr	Swamp paperbark	<i>Melaleuca rhaphiophylla</i>
Ol	Narrow-leaved Oxylobium	<i>Oxylobium lineare</i>
Pe	Bracken fern	<i>Pteridium esculentum</i>
Vj	Swishbush	<i>Viminaria juncea</i>

Cadastral and Streetsmart data supplied by the Dept. of Land Administration (1998)

**Map Legend**