

GOSN ϕ 11

Wet flat, bare ground absent in
spring, summer annuals $\approx 10\%$
Litter 2%

Sandy clay

Centropogon nutans

CORN 11






BUSHLAND PLANT SURVEY RECORDING SHEET 2 - use pencil only




From 'Bushland Plant Survey' written by B. Keighery (1994) and published by the Wildflower Society of WA (Inc.), PO Box 64 Nedlands WA 6008.




3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below), and dominant species in each layer.

Cover Class	2-10%	10-30%	30-70%	over 70%
-------------	-------	--------	--------	----------

LIFE FORM	TREES			MALLEES	
	over 30m	10-30m	under 10m	over 8m	under 8m
					
COVER CLASS (%)					
DOMINANT SPECIES					

LIFE FORM	SHRUBS		SHRUBS
	over 2m	2m - 1m	under 1m
		<i>1m - 1.5</i> 	<i>< 0.5</i> 
COVER CLASS (%)		<i>2-10%</i>	<i>30-70</i>
DOMINANT SPECIES		<i>mixed.</i>	<i>vet lindeneyi</i>

LIFE FORM	GRASSES	HERBS	SEDGES	OTHER
				
COVER CLASS (%)		<i>10-30%</i>	<i>30-70%</i>	
DOMINANT SPECIES		<i>Borja Scirp Drosera zig</i>	<i>Lox pub Schoenus</i>	

4. VEGETATION CONDITION

1	'PRISTINE'	COMMENTS <i>Burnt, not all 2 years.</i>
2	EXCELLENT <input checked="" type="checkbox"/>	
3	VERY GOOD	
4	GOOD	
5	DEGRADED	

