
















BUSHLAND PLANT SURVEY RECORDING SHEET 2 (Muir)- use pencil only

From 'Bushland Plant Survey' written by B. Keighery (1994) and published by the Wildflower Society of WA Inc., PO Box 64 Nedlands WA 6008.

3. VEGETATION STRUCTURE AND COVER

For each layer record - appropriate life form, cover class (see below) and dominant species in each layer.

Cover Class 2-10% 10-30% 30 - 70% over 70%

TREES				MALLEES	
over 30m		15 - 30m	5 - 15m	over 9m	under 9m
			under 5m		30m
LIFE FORM			12m 		
COVER CLASS (%)		2-10	10-30		
DOMINANT SPECIES		Marr	Marr		
SHRUBS			SHRUBS		
over 2m		2m - 1.5m	1.5 - 1m	1 - 0.5m	under 0.5m
LIFE FORM	 				
COVER CLASS (%)	2-10			30-70	10-30
DOMINANT SPECIES	Jack tree			Dox ptyocida Alloea hum	South horn Daisy horn
GRASSES		HERBS	SEDGES	over 0.5m	under 0.5m
LIFE FORM					
COVER CLASS (%)	10-30				
DOMINANT SPECIES	Buzza murex				

4. VEGETATION CONDITION

1	'PRISTINE'	COMMENTS Die back front south off
2	EXCELLENT	
3	VERY GOOD ✓	
4	GOOD	
5	DEGRADED	

